



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 17, 2024 – 02:11 PM EST

PDB ID : 4RGW  
Title : Crystal Structure of a TAF1-TAF7 Complex in Human Transcription Factor IID  
Authors : Wang, H.; Curran, E.C.; Hinds, T.R.; Wang, E.H.; Zheng, N.  
Deposited on : 2014-09-30  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	638	 2% 56% 7% 36%
2	B	350	 30% 5% 65%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4500 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription initiation factor TFIID subunit 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	406	3290	2090	580	596	2	22	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	599	SER	-	expression tag	UNP P21675

- Molecule 2 is a protein called Transcription initiation factor TFIID subunit 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	123	998	638	184	172	4	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	SER	-	expression tag	UNP Q15545

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	155	Total O 155 155	0	0
4	B	45	Total O 45 45	0	0



GLU  
ASP  
ARG  
GLU  
LYS  
GLU  
GLN  
LEU  
SER  
SER  
LEU  
GLN  
GLU  
LEU  
GLU  
SER  
SER  
LEU  
LEU  
GLU  
LYS

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.22Å 94.54Å 101.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.27 – 2.30 47.27 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.27-2.30) 99.8 (47.27-2.30)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.25 (at 2.29Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.186 , 0.234 0.190 , 0.236	Depositor DCC
$R_{free}$ test set	1813 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.1	Xtrriage
Anisotropy	0.466	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 28.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4500	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, TPO, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/3349	0.55	0/4506
2	B	0.41	0/1017	0.59	1/1370 (0.1%)
All	All	0.41	0/4366	0.56	1/5876 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	143	VAL	CB-CA-C	-5.24	101.44	111.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1104	LEU	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3290	0	3278	19	0
2	B	998	0	1055	8	0
3	A	12	0	16	1	0
4	A	155	0	0	0	0
4	B	45	0	0	0	0
All	All	4500	0	4349	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:819:ARG:HH11	1:A:819:ARG:HG3	1.56	0.69
1:A:835:LEU:HB3	1:A:846:ILE:HG21	1.77	0.66
1:A:678:SER:N	1:A:679:GLY:HA2	2.16	0.61
1:A:819:ARG:HG3	1:A:819:ARG:NH1	2.16	0.61
2:B:140:LEU:O	2:B:143:VAL:HG22	2.06	0.56
1:A:829:GLN:OE1	1:A:866:ARG:HG3	2.07	0.54
1:A:970:GLU:HG2	1:A:972:THR:O	2.08	0.53
1:A:943:LYS:HZ2	1:A:943:LYS:N	2.08	0.52
1:A:796:GLU:HG3	2:B:29:ARG:NH1	2.26	0.50
1:A:1104:LEU:O	2:B:141:LYS:HG3	2.12	0.49
2:B:75:GLU:HG2	2:B:87:LYS:HE3	1.95	0.49
1:A:865:LYS:O	1:A:868:LYS:HG2	2.15	0.47
1:A:968:LEU:HB2	1:A:989:ILE:HG13	1.98	0.45
1:A:911:SER:HA	3:A:1302:GOL:H11	1.98	0.45
2:B:53:HIS:CG	2:B:105:PRO:HG2	2.51	0.45
1:A:711:LEU:HD22	1:A:768:LEU:HD22	1.99	0.43
1:A:820:ALA:O	1:A:824:ILE:HG13	2.18	0.43
2:B:105:PRO:HA	2:B:106:PRO:HD2	1.88	0.42
1:A:647:GLY:HA3	1:A:789:ARG:HG2	2.03	0.41
1:A:660:LEU:HD22	1:A:793:TYR:HB3	2.02	0.41
2:B:21:PRO:HA	2:B:22:PRO:HD3	1.92	0.41
1:A:722:LYS:HD3	1:A:722:LYS:HA	1.86	0.40
1:A:718:ALA:HA	2:B:85:PHE:O	2.21	0.40
1:A:1084:LYS:HD2	1:A:1084:LYS:HA	1.66	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	398/638 (62%)	391 (98%)	7 (2%)	0	100	100
2	B	119/350 (34%)	117 (98%)	2 (2%)	0	100	100
All	All	517/988 (52%)	508 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	355/558 (64%)	331 (93%)	24 (7%)	13	18
2	B	113/323 (35%)	104 (92%)	9 (8%)	10	13
All	All	468/881 (53%)	435 (93%)	33 (7%)	12	17

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	617	VAL
1	A	619	LEU
1	A	660	LEU
1	A	666	LYS
1	A	669	LYS
1	A	670	MET
1	A	675	ARG
1	A	699	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	701	LEU
1	A	748	THR
1	A	752	LEU
1	A	797	LEU
1	A	852	LYS
1	A	866	ARG
1	A	868	LYS
1	A	869	LEU
1	A	878	MET
1	A	943	LYS
1	A	944	ILE
1	A	963	MET
1	A	1084	LYS
1	A	1093	LEU
1	A	1104	LEU
1	A	1108	THR
2	B	23	GLU
2	B	73	VAL
2	B	99	VAL
2	B	100	ASP
2	B	129	LYS
2	B	138	LEU
2	B	141	LYS
2	B	143	VAL
2	B	154	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
1	TPO	A	1106	1	8,10,11	1.17	0	10,14,16	1.79	1 (10%)
1	SEP	A	1105	1	8,9,10	1.68	1 (12%)	7,12,14	1.17	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	1106	1	-	0/9/11/13	-
1	SEP	A	1105	1	-	3/6/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1105	SEP	P-O1P	3.67	1.61	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1106	TPO	P-OG1-CB	-5.08	109.52	123.33
1	A	1105	SEP	OG-P-O1P	2.27	112.57	106.44

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1105	SEP	CB-OG-P-O2P
1	A	1105	SEP	CB-OG-P-O3P
1	A	1105	SEP	N-CA-CB-OG

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	A	1301	-	5,5,5	0.38	0	5,5,5	0.43	0
3	GOL	A	1302	-	5,5,5	0.43	0	5,5,5	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1301	-	-	0/4/4/4	-
3	GOL	A	1302	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1302	GOL	O1-C1-C2-C3
3	A	1302	GOL	C1-C2-C3-O3
3	A	1302	GOL	O1-C1-C2-O2
3	A	1302	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1302	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	404/638 (63%)	-0.39	13 (3%) 50 52	5, 14, 46, 84	0
2	B	123/350 (35%)	-0.70	3 (2%) 59 61	4, 10, 29, 44	0
All	All	527/988 (53%)	-0.46	16 (3%) 52 54	4, 13, 43, 84	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	880	SER	6.1
1	A	747	HIS	4.9
1	A	943	LYS	4.2
1	A	879	ASP	4.1
1	A	876	THR	3.5
1	A	877	GLY	3.5
1	A	927	LYS	3.1
2	B	106	PRO	2.7
1	A	992	LYS	2.7
1	A	875	ARG	2.6
1	A	878	MET	2.5
1	A	679	GLY	2.4
2	B	128	LYS	2.3
1	A	1108	THR	2.2
1	A	929	PHE	2.2
2	B	100	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	SEP	A	1105	10/11	0.77	0.13	20,42,52,68	0
1	TPO	A	1106	11/12	0.92	0.08	20,24,37,38	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	1301	6/6	0.94	0.09	12,13,16,19	0
3	GOL	A	1302	6/6	0.94	0.08	13,15,15,18	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.