



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2024 – 07:09 PM EDT

PDB ID : 4RI4
Title : Crystal structure of PTPN3 (PTPH1) Y676I mutant in complex with vanadate
Authors : Chen, K.-E.; Meng, T.C.; Wang, A.H.-J.
Deposited on : 2014-10-05
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	
1	B	306	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9777 atoms, of which 4636 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	283	4592	1452	2304	387	431	18	12	9	0
1	B	281	4640	1466	2332	392	431	19	25	14	0

There are 50 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	expression tag	UNP P26045
A	-22	HIS	-	expression tag	UNP P26045
A	-21	HIS	-	expression tag	UNP P26045
A	-20	HIS	-	expression tag	UNP P26045
A	-19	HIS	-	expression tag	UNP P26045
A	-18	HIS	-	expression tag	UNP P26045
A	-17	HIS	-	expression tag	UNP P26045
A	-16	SER	-	expression tag	UNP P26045
A	-15	SER	-	expression tag	UNP P26045
A	-14	GLY	-	expression tag	UNP P26045
A	-13	VAL	-	expression tag	UNP P26045
A	-12	ASP	-	expression tag	UNP P26045
A	-11	LEU	-	expression tag	UNP P26045
A	-10	GLY	-	expression tag	UNP P26045
A	-9	THR	-	expression tag	UNP P26045
A	-8	GLU	-	expression tag	UNP P26045
A	-7	ASN	-	expression tag	UNP P26045
A	-6	LEU	-	expression tag	UNP P26045
A	-5	TYR	-	expression tag	UNP P26045
A	-4	PHE	-	expression tag	UNP P26045
A	-3	GLN	-	expression tag	UNP P26045
A	-2	SER	-	expression tag	UNP P26045
A	-1	ASN	-	expression tag	UNP P26045
A	0	ALA	-	expression tag	UNP P26045
A	676	ILE	TYR	engineered mutation	UNP P26045

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-23	MET	-	expression tag	UNP P26045
B	-22	HIS	-	expression tag	UNP P26045
B	-21	HIS	-	expression tag	UNP P26045
B	-20	HIS	-	expression tag	UNP P26045
B	-19	HIS	-	expression tag	UNP P26045
B	-18	HIS	-	expression tag	UNP P26045
B	-17	HIS	-	expression tag	UNP P26045
B	-16	SER	-	expression tag	UNP P26045
B	-15	SER	-	expression tag	UNP P26045
B	-14	GLY	-	expression tag	UNP P26045
B	-13	VAL	-	expression tag	UNP P26045
B	-12	ASP	-	expression tag	UNP P26045
B	-11	LEU	-	expression tag	UNP P26045
B	-10	GLY	-	expression tag	UNP P26045
B	-9	THR	-	expression tag	UNP P26045
B	-8	GLU	-	expression tag	UNP P26045
B	-7	ASN	-	expression tag	UNP P26045
B	-6	LEU	-	expression tag	UNP P26045
B	-5	TYR	-	expression tag	UNP P26045
B	-4	PHE	-	expression tag	UNP P26045
B	-3	GLN	-	expression tag	UNP P26045
B	-2	SER	-	expression tag	UNP P26045
B	-1	ASN	-	expression tag	UNP P26045
B	0	ALA	-	expression tag	UNP P26045
B	676	ILE	TYR	engineered mutation	UNP P26045

- Molecule 2 is VANADATE ION (three-letter code: VO4) (formula: O₄V).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	V	0	0
			5	4	1		
2	B	1	Total	O	V	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	277	Total	O	0	0
			277	277		
3	B	258	Total	O	0	0
			258	258		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 3



- Molecule 1: Tyrosine-protein phosphatase non-receptor type 3



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	94.88Å 99.41Å 134.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.00 – 1.60 24.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.7 (24.00-1.60) 98.7 (24.00-1.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 1.60Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.163 , 0.189 0.165 , 0.191	Depositor DCC
R_{free} test set	4143 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	20.6	Xtrriage
Anisotropy	0.299	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 48.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.035 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9777	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: VO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/2363	0.69	0/3214
1	B	0.50	0/2399	0.66	0/3261
All	All	0.52	0/4762	0.67	0/6475

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2288	2304	2298	16	0
1	B	2308	2332	2326	8	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	277	0	0	3	0
3	B	258	0	0	0	0
All	All	5141	4636	4624	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:-2:SER:HA	1:A:867:PRO:CD	2.26	0.66
1:A:833:VAL:HG23	1:A:836:GLU:HB3	1.79	0.64
1:B:834:ASP:O	1:B:835:SER:OG	2.16	0.58
1:A:795:THR:HG23	1:A:797:GLU:H	1.67	0.58
1:A:730:GLN:NE2	1:A:734:ASP:OD1	2.37	0.54
1:A:655:ARG:CZ	3:A:1285:HOH:O	2.55	0.53
1:A:655:ARG:NH1	3:A:1285:HOH:O	2.44	0.50
1:B:640:GLY:HA3	1:B:646:VAL:HG23	1.95	0.47
1:A:833:VAL:CG2	1:A:836:GLU:HB3	2.44	0.47
1:A:881:ARG:O	1:A:884[B]:MET:HG2	2.13	0.47
1:B:874:VAL:HA	1:B:877:MET:HE2	1.97	0.47
1:A:825:VAL:HG22	1:A:852:LEU:CD2	2.43	0.47
1:A:809:TRP:CE2	1:A:848:ARG:HG2	2.50	0.46
1:A:709:ASN:O	1:A:709:ASN:ND2	2.47	0.46
1:B:834:ASP:O	1:B:835:SER:CB	2.64	0.46
1:B:858:ALA:O	1:B:862[B]:THR:HG23	2.17	0.44
1:B:809:TRP:CE2	1:B:848:ARG:HG2	2.54	0.43
1:A:833:VAL:HG23	1:A:836:GLU:CB	2.48	0.43
1:A:903:GLU:OE2	3:A:1217:HOH:O	2.21	0.43
1:A:862[B]:THR:HG23	1:A:868:ILE:HD11	2.01	0.42
1:A:720:PRO:HG3	1:A:728:PHE:CD2	2.55	0.42
1:B:881:ARG:O	1:B:884[B]:MET:HG2	2.21	0.41
1:B:647:LEU:HD12	1:B:888[A]:SER:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	290/306 (95%)	280 (97%)	9 (3%)	1 (0%)	41	21
1	B	293/306 (96%)	286 (98%)	6 (2%)	1 (0%)	41	21

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	583/612 (95%)	566 (97%)	15 (3%)	2 (0%)	34	21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	885	VAL
1	B	885	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/274 (96%)	260 (99%)	2 (1%)	81	70
1	B	266/274 (97%)	264 (99%)	2 (1%)	81	70
All	All	528/548 (96%)	524 (99%)	4 (1%)	86	70

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-1	ASN
1	A	834	ASP
1	B	771[A]	HIS
1	B	771[B]	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	812	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	VO4	B	1001	-	1,4,4	3.09	1 (100%)	-		
2	VO4	A	1001	-	1,4,4	6.35	1 (100%)	-		

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	VO4	O1-V	6.35	2.00	1.63
2	B	1001	VO4	O1-V	3.09	1.81	1.63

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	283/306 (92%)	0.21	24 (8%) 10 9	13, 21, 43, 62	4 (1%)
1	B	281/306 (91%)	0.21	18 (6%) 19 17	14, 22, 45, 63	5 (1%)
All	All	564/612 (92%)	0.21	42 (7%) 14 13	13, 21, 44, 63	9 (1%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	706	PRO	7.3
1	A	707	ALA	6.5
1	B	793	THR	6.4
1	A	691	GLY	5.9
1	A	692	ASN	5.9
1	A	762	PRO	5.5
1	B	795	THR	5.4
1	A	794	GLN	5.4
1	B	794	GLN	4.6
1	B	706	PRO	4.5
1	A	709	ASN	4.5
1	A	-2	SER	4.5
1	B	709	ASN	4.0
1	A	761	PRO	3.9
1	B	764	VAL	3.7
1	A	795	THR	3.6
1	A	690	GLN	3.5
1	A	839[A]	LEU	3.4
1	A	842	CYS	3.3
1	A	873	ILE	3.3
1	B	707	ALA	3.1
1	B	690	GLN	3.0
1	B	762	PRO	2.9
1	A	793	THR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	834	ASP	2.8
1	B	769	GLY	2.8
1	B	693	GLU	2.8
1	A	0	ALA	2.7
1	B	796	GLY	2.6
1	A	792	ASN	2.5
1	B	757	TYR	2.4
1	A	835	SER	2.4
1	A	764	VAL	2.3
1	B	835	SER	2.2
1	A	705	ILE	2.2
1	A	796	GLY	2.1
1	B	729	TRP	2.1
1	B	791	THR	2.1
1	A	728	PHE	2.1
1	A	-1	ASN	2.0
1	B	692	ASN	2.0
1	B	768	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	VO4	B	1001	5/5	0.98	0.12	15,17,19,19	1
2	VO4	A	1001	5/5	0.99	0.13	12,17,18,19	1

6.5 Other polymers [i](#)

There are no such residues in this entry.