



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 18, 2023 – 05:41 am GMT

PDB ID : 4U4J
Title : Crystal structure of Salmonella alpha-2-macroglobulin mutant Y1175G
Authors : Wong, S.G.; Dessen, A.
Deposited on : 2014-07-23
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1647	

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 23627 atoms, of which 11754 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative inner membrane lipoprotein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	1538	23627	7480	11754	2091	2279	23	0	0	0


There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	initiating methionine	UNP Q8ZN46
A	-1	GLY	-	expression tag	UNP Q8ZN46
A	0	SER	-	expression tag	UNP Q8ZN46
A	1	SER	-	expression tag	UNP Q8ZN46
A	2	HIS	-	expression tag	UNP Q8ZN46
A	3	HIS	-	expression tag	UNP Q8ZN46
A	4	HIS	-	expression tag	UNP Q8ZN46
A	5	HIS	-	expression tag	UNP Q8ZN46
A	6	HIS	-	expression tag	UNP Q8ZN46
A	7	HIS	-	expression tag	UNP Q8ZN46
A	8	SER	-	expression tag	UNP Q8ZN46
A	9	SER	-	expression tag	UNP Q8ZN46
A	10	GLY	-	expression tag	UNP Q8ZN46
A	11	LEU	-	expression tag	UNP Q8ZN46
A	12	VAL	-	expression tag	UNP Q8ZN46
A	13	PRO	-	expression tag	UNP Q8ZN46
A	14	ARG	-	expression tag	UNP Q8ZN46
A	15	GLY	-	expression tag	UNP Q8ZN46
A	16	SER	-	expression tag	UNP Q8ZN46
A	17	HIS	-	expression tag	UNP Q8ZN46
A	18	MET	-	expression tag	UNP Q8ZN46
A	98	ALA	LYS	engineered mutation	UNP Q8ZN46
A	99	ALA	LYS	engineered mutation	UNP Q8ZN46
A	1175	GLY	TYR	engineered mutation	UNP Q8ZN46

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative inner membrane lipoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	254.24Å 82.16Å 98.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.99 – 2.90 58.99 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (58.99-2.90) 99.9 (58.99-2.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 2.91Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.244 , 0.284 0.250 , 0.287	Depositor DCC
R_{free} test set	2340 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	87.1	Xtrriage
Anisotropy	0.461	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	23627	wwPDB-VP
Average B, all atoms (Å ²)	102.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/12113	0.43	1/16503 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	316	LEU	CA-CB-CG	7.46	132.47	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	252	LYS	Peptide
1	A	808	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11873	11754	11772	59	0
All	All	11873	11754	11772	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87:GLN:N	1:A:87:GLN:OE1	2.19	0.76
1:A:1366:GLU:O	1:A:1368:GLN:N	2.19	0.75
1:A:1368:GLN:OE1	1:A:1369:TRP:N	2.30	0.65
1:A:463:LEU:O	1:A:465:ARG:NH1	2.34	0.60
1:A:317:ASN:O	1:A:319:LYS:N	2.33	0.60
1:A:1091:GLN:NE2	1:A:1110:GLN:OE1	2.36	0.58
1:A:337:GLU:N	1:A:337:GLU:OE1	2.37	0.57
1:A:1181:GLU:OE1	1:A:1416:THR:OG1	2.22	0.57
1:A:829:ARG:O	1:A:831:ALA:N	2.38	0.57
1:A:1294:GLN:OE1	1:A:1328:GLN:NE2	2.38	0.57
1:A:1566:GLU:OE1	1:A:1566:GLU:N	2.40	0.55
1:A:84:ASP:HB2	1:A:136:LYS:HE3	1.89	0.55
1:A:922:ASP:OD2	1:A:924:TYR:N	2.40	0.54
1:A:1241:GLU:OE2	1:A:1278:TYR:OH	2.19	0.54
1:A:349:LYS:O	1:A:351:GLU:N	2.41	0.53
1:A:663:ARG:NH2	1:A:710:TRP:O	2.40	0.53
1:A:784:SER:O	1:A:786:GLY:N	2.41	0.53
1:A:555:LEU:O	1:A:557:LYS:N	2.42	0.53
1:A:1105:ASP:OD1	1:A:1106:ALA:N	2.42	0.53
1:A:1181:GLU:O	1:A:1184:THR:N	2.42	0.52
1:A:1583:GLU:OE2	1:A:1585:ARG:NE	2.42	0.52
1:A:137:ALA:O	1:A:140:ASN:N	2.43	0.52
1:A:758:ASN:O	1:A:758:ASN:ND2	2.42	0.51
1:A:555:LEU:HD23	1:A:555:LEU:C	2.32	0.50
1:A:320:GLY:O	1:A:321:GLN:NE2	2.45	0.49
1:A:1227:ARG:NH1	1:A:1241:GLU:OE1	2.46	0.49
1:A:1347:ASP:OD2	1:A:1350:ARG:NH1	2.45	0.49
1:A:136:LYS:HE2	1:A:138:LEU:HA	1.96	0.48
1:A:1289:TYR:OH	1:A:1578:ASP:OD2	2.22	0.48
1:A:983:ASP:OD1	1:A:1109:LYS:NZ	2.47	0.47
1:A:1494:ASN:O	1:A:1496:LEU:N	2.47	0.47
1:A:204:SER:OG	1:A:319:LYS:O	2.27	0.46
1:A:103:VAL:HG22	1:A:104:ASP:H	1.80	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1251:ASP:OD1	1:A:1310:ARG:NH1	2.50	0.45
1:A:486:GLN:O	1:A:488:THR:N	2.50	0.45
1:A:134:ALA:O	1:A:136:LYS:N	2.51	0.44
1:A:557:LYS:HG2	1:A:558:GLY:H	1.82	0.44
1:A:1610:VAL:HG23	1:A:1611:THR:N	2.33	0.44
1:A:251:ILE:HG23	1:A:253:PRO:HD2	2.00	0.44
1:A:134:ALA:C	1:A:136:LYS:H	2.20	0.43
1:A:84:ASP:HB2	1:A:136:LYS:HD3	2.01	0.43
1:A:1364:GLN:OE1	1:A:1364:GLN:N	2.49	0.42
1:A:604:ASP:N	1:A:604:ASP:OD1	2.52	0.42
1:A:420:LYS:HB2	1:A:421:PRO:HD2	2.01	0.42
1:A:136:LYS:HZ3	1:A:137:ALA:CA	2.33	0.42
1:A:957:VAL:O	1:A:958:ASN:CB	2.68	0.42
1:A:193:ARG:NH2	1:A:256:GLN:OE1	2.53	0.41
1:A:252:LYS:HG3	1:A:255:GLN:HB2	2.01	0.41
1:A:1094:ILE:HB	1:A:1107:GLN:HB2	2.01	0.41
1:A:1288:ARG:NH2	1:A:1289:TYR:OH	2.54	0.41
1:A:810:TRP:CD1	1:A:810:TRP:N	2.89	0.41
1:A:1242:GLU:OE1	1:A:1373:TYR:OH	2.31	0.41
1:A:173:ILE:CD1	1:A:175:GLU:HB2	2.51	0.41
1:A:477:GLU:O	1:A:479:MET:N	2.54	0.41
1:A:1368:GLN:N	1:A:1368:GLN:CD	2.73	0.41
1:A:832:THR:HA	1:A:833:PRO:HD2	2.00	0.40
1:A:1333:LEU:HB3	1:A:1334:PRO:HD3	2.03	0.40
1:A:135:VAL:CG2	1:A:143:PHE:HB2	2.51	0.40
1:A:136:LYS:HE2	1:A:137:ALA:C	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1526/1647 (93%)	1375 (90%)	106 (7%)	45 (3%)	4	18

All (45) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	321	GLN
1	A	556	ASP
1	A	587	SER
1	A	651	LYS
1	A	758	ASN
1	A	882	LYS
1	A	957	VAL
1	A	1098	ASN
1	A	1367	ARG
1	A	1368	GLN
1	A	1370	ILE
1	A	134	ALA
1	A	137	ALA
1	A	142	THR
1	A	231	ARG
1	A	829	ARG
1	A	830	SER
1	A	958	ASN
1	A	983	ASP
1	A	1347	ASP
1	A	1635	ALA
1	A	135	VAL
1	A	487	LYS
1	A	588	GLY
1	A	876	LYS
1	A	1369	TRP
1	A	1411	GLN
1	A	90	SER
1	A	307	ALA
1	A	319	LYS
1	A	350	GLU
1	A	832	THR
1	A	1104	LEU
1	A	174	ALA
1	A	253	PRO
1	A	831	ALA
1	A	881	PRO
1	A	984	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1178	GLY
1	A	1206	ASP
1	A	1394	PRO
1	A	929	GLU
1	A	1161	PRO
1	A	103	VAL
1	A	824	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1259/1347 (94%)	1227 (98%)	32 (2%)	47 78

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	GLN
1	A	53	GLN
1	A	70	ASP
1	A	87	GLN
1	A	91	ARG
1	A	136	LYS
1	A	142	THR
1	A	154	ARG
1	A	227	VAL
1	A	248	LEU
1	A	256	GLN
1	A	311	ILE
1	A	319	LYS
1	A	331	GLN
1	A	334	VAL
1	A	350	GLU
1	A	439	ARG
1	A	555	LEU
1	A	586	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	663	ARG
1	A	753	LYS
1	A	766	LYS
1	A	826	ASP
1	A	876	LYS
1	A	926	GLN
1	A	1179	CYS
1	A	1303	TYR
1	A	1321	GLU
1	A	1367	ARG
1	A	1368	GLN
1	A	1378	ARG
1	A	1400	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	758	ASN
1	A	956	HIS
1	A	1091	GLN
1	A	1328	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1538/1647 (93%)	0.62	101 (6%) 18 14	51, 90, 137, 213	0

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	588	GLY	7.8
1	A	928	ILE	7.3
1	A	70	ASP	6.0
1	A	57	LYS	5.7
1	A	85	PRO	5.3
1	A	929	GLU	5.3
1	A	559	GLY	5.0
1	A	937	LEU	4.7
1	A	83	LEU	4.5
1	A	589	GLY	4.4
1	A	649	GLY	4.2
1	A	930	GLY	4.1
1	A	587	SER	4.0
1	A	194	VAL	3.9
1	A	1176	PRO	3.9
1	A	118	LEU	3.8
1	A	223	MET	3.7
1	A	1570	LEU	3.6
1	A	343	ALA	3.6
1	A	325	GLN	3.5
1	A	622	ARG	3.4
1	A	650	GLU	3.4
1	A	768	HIS	3.4
1	A	314	VAL	3.4
1	A	758	ASN	3.4
1	A	851	LEU	3.4
1	A	87	GLN	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	519	LEU	3.3
1	A	761	PRO	3.3
1	A	225	ASP	3.2
1	A	75	LEU	3.2
1	A	1101	GLY	3.2
1	A	108	GLU	3.2
1	A	1178	GLY	3.1
1	A	228	TYR	3.0
1	A	551	VAL	3.0
1	A	764	THR	3.0
1	A	1175	GLY	3.0
1	A	596	VAL	3.0
1	A	762	GLY	3.0
1	A	1567	VAL	2.9
1	A	548	LEU	2.8
1	A	1364	GLN	2.7
1	A	498	PHE	2.6
1	A	1099	LEU	2.5
1	A	247	PRO	2.5
1	A	311	ILE	2.5
1	A	802	LEU	2.5
1	A	234	LEU	2.5
1	A	988	LEU	2.5
1	A	202	PHE	2.5
1	A	262	ALA	2.5
1	A	248	LEU	2.5
1	A	349	LYS	2.4
1	A	172	LYS	2.4
1	A	926	GLN	2.4
1	A	517	GLY	2.4
1	A	808	LYS	2.4
1	A	174	ALA	2.4
1	A	515	LEU	2.4
1	A	89	PHE	2.4
1	A	91	ARG	2.4
1	A	353	THR	2.3
1	A	344	LEU	2.3
1	A	767	LEU	2.3
1	A	927	VAL	2.3
1	A	281	LEU	2.3
1	A	638	ALA	2.3
1	A	1152	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	817	LEU	2.3
1	A	625	THR	2.2
1	A	358	LEU	2.2
1	A	924	TYR	2.2
1	A	116	LEU	2.2
1	A	674	GLU	2.2
1	A	1630	TRP	2.2
1	A	1457	LEU	2.2
1	A	934	LEU	2.2
1	A	1177	TYR	2.2
1	A	191	PHE	2.1
1	A	312	GLU	2.1
1	A	226	LEU	2.1
1	A	735	TRP	2.1
1	A	875	VAL	2.1
1	A	1367	ARG	2.1
1	A	304	ASN	2.1
1	A	810	TRP	2.1
1	A	54	SER	2.1
1	A	1103	THR	2.1
1	A	113	LEU	2.1
1	A	261	VAL	2.1
1	A	254	LEU	2.1
1	A	1509	ARG	2.1
1	A	400	LEU	2.1
1	A	322	THR	2.1
1	A	766	LYS	2.0
1	A	931	GLN	2.0
1	A	616	LYS	2.0
1	A	1546	LEU	2.0
1	A	107	TRP	2.0
1	A	192	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.