



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 4, 2024 – 03:36 pm GMT

PDB ID : 4UMC
Title : Structural analysis of substrate-mimicking inhibitors in complex with *Neisseria meningitidis* 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase - the importance of accommodating the active site water
Authors : Heyes, L.C.; Reichau, S.; Cross, P.J.; Parker, E.J.
Deposited on : 2014-05-16
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	91% 5%
1	B	351	88% 7% 5%
1	C	351	90% 5% 5%
1	D	351	90% 5% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	B	1353	-	-	-	X
4	PEQ	D	1352	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10499 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	334	Total 2562	C 1599	N 463	O 486	S 14	0	2	0
1	B	334	Total 2551	C 1596	N 457	O 483	S 15	0	2	0
1	C	335	Total 2530	C 1581	N 455	O 480	S 14	0	1	0
1	D	333	Total 2529	C 1581	N 455	O 479	S 14	0	2	0

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Mn 1	0	0
2	B	1	Total 1	Mn 1	0	0
2	C	1	Total 1	Mn 1	0	0
2	D	1	Total 1	Mn 1	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		
3	D	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is L-PHOSPHOLACTATE (three-letter code: PEQ) (formula: $C_3H_7O_6P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			10	3	6	1		
4	B	1	Total	C	O	P	0	0
			10	3	6	1		
4	C	1	Total	C	O	P	0	0
			10	3	6	1		
4	D	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	96	Total	O	0	0
			96	96		
5	B	65	Total	O	0	0
			65	65		
5	C	57	Total	O	0	0
			57	57		
5	D	45	Total	O	0	0
			45	45		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE

Chain A: 




- Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE

Chain B: 



- Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE

Chain C: 



- Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE

Chain D: 



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.42Å 137.01Å 76.48Å 90.00° 96.48° 90.00°	Depositor
Resolution (Å)	75.99 – 2.34 40.33 – 2.34	Depositor EDS
% Data completeness (in resolution range)	99.9 (75.99-2.34) 99.9 (40.33-2.34)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.203 , 0.230 0.209 , 0.233	Depositor DCC
R_{free} test set	3205 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	33.6	Xtrriage
Anisotropy	0.482	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 24.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.027 for l,-k,h	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10499	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEQ, PO4, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	0/2610	0.71	0/3536
1	B	0.58	0/2599	0.70	1/3521 (0.0%)
1	C	0.60	0/2578	0.70	3/3499 (0.1%)
1	D	0.56	0/2577	0.68	0/3496
All	All	0.58	0/10364	0.70	4/14052 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	236	ARG	NE-CZ-NH1	5.70	123.15	120.30
1	C	82	ARG	NE-CZ-NH2	-5.63	117.49	120.30
1	C	256	ARG	NE-CZ-NH2	-5.52	117.54	120.30
1	C	256	ARG	NE-CZ-NH1	5.28	122.94	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	280	GLN	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2562	0	2511	8	0
1	B	2551	0	2508	14	0
1	C	2530	0	2456	10	0
1	D	2529	0	2464	10	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	5	0	0	1	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
4	A	10	0	4	1	0
4	B	10	0	4	0	0
4	C	10	0	4	0	0
4	D	10	0	4	1	0
5	A	96	0	0	2	0
5	B	65	0	0	4	0
5	C	57	0	0	5	0
5	D	45	0	0	1	0
All	All	10499	0	9955	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:219:HIS:ND1	1:C:219:HIS:HE1	1.45	1.14
5:C:2045:HOH:O	1:D:148[A]:ASP:OD1	1.75	1.02
1:A:148[A]:ASP:OD2	5:A:2049:HOH:O	1.87	0.93
1:B:219:HIS:ND1	1:C:219:HIS:CE1	2.38	0.92
1:D:311:ARG:NH1	1:D:323:THR:O	2.05	0.90
1:B:236:ARG:O	1:B:271:ALA:HB3	1.86	0.75
1:B:278:THR:O	5:B:2057:HOH:O	2.08	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:2045:HOH:O	1:D:148[A]:ASP:CG	2.25	0.68
1:B:112:ASP:OD1	5:B:2025:HOH:O	2.12	0.67
1:D:166:ALA:HB2	1:D:236:ARG:HD3	1.80	0.63
1:C:17:GLU:CB	5:C:2001:HOH:O	2.53	0.56
1:C:267:ASP:HA	1:C:302:MET:HB3	1.89	0.55
1:D:17:GLU:HA	5:D:2001:HOH:O	2.07	0.54
1:D:267:ASP:HA	1:D:302:MET:HB3	1.90	0.54
1:B:219:HIS:HD2	1:B:220:SER:O	1.91	0.53
1:A:267:ASP:HA	1:A:302:MET:HB3	1.91	0.52
1:C:17:GLU:HB3	5:C:2001:HOH:O	2.08	0.52
1:C:114:HIS:HB2	1:C:116:ASP:OD1	2.10	0.52
1:B:267:ASP:HA	1:B:302:MET:HB3	1.92	0.52
1:B:292:GLU:O	1:B:349:ARG:NH1	2.34	0.49
1:A:99:LYS:NZ	4:A:1352:PEQ:O2'	2.41	0.49
1:A:219:HIS:NE2	1:D:219:HIS:ND1	2.62	0.45
1:C:151:THR:N	1:C:152:PRO:CD	2.80	0.45
1:C:267:ASP:OD2	1:C:270:HIS:ND1	2.38	0.45
1:B:151:THR:N	1:B:152:PRO:CD	2.80	0.45
1:B:210:HIS:CD2	5:B:2046:HOH:O	2.70	0.45
1:A:101:ARG:NE	3:A:1351:PO4:O1	2.22	0.44
1:C:17:GLU:N	5:C:2001:HOH:O	2.48	0.44
1:D:114:HIS:HB2	1:D:116:ASP:OD1	2.17	0.44
1:D:151:THR:N	1:D:152:PRO:CD	2.81	0.43
1:A:151:THR:N	1:A:152:PRO:CD	2.82	0.43
1:D:236:ARG:HD2	4:D:1352:PEQ:O1P	2.19	0.42
1:B:31:SER:HB2	5:B:2003:HOH:O	2.19	0.42
1:A:347:MET:HB2	5:A:2096:HOH:O	2.21	0.41
1:B:348:ALA:C	1:B:350:ALA:H	2.24	0.41
1:B:237:GLY:HA3	1:B:271:ALA:HB3	2.03	0.41
1:A:236:ARG:C	1:A:236:ARG:HD3	2.42	0.40
1:B:285:GLN:HA	1:B:338:LEU:HD21	2.03	0.40
1:C:285:GLN:HA	1:C:338:LEU:HD21	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/351 (95%)	327 (98%)	7 (2%)	0	100	100
1	B	334/351 (95%)	327 (98%)	7 (2%)	0	100	100
1	C	334/351 (95%)	327 (98%)	7 (2%)	0	100	100
1	D	333/351 (95%)	325 (98%)	8 (2%)	0	100	100
All	All	1335/1404 (95%)	1306 (98%)	29 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/289 (92%)	262 (98%)	5 (2%)	57	68
1	B	266/289 (92%)	260 (98%)	6 (2%)	50	61
1	C	260/289 (90%)	253 (97%)	7 (3%)	44	55
1	D	261/289 (90%)	255 (98%)	6 (2%)	50	61
All	All	1054/1156 (91%)	1030 (98%)	24 (2%)	50	61

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	189	ASN
1	A	233	VAL
1	A	236	ARG
1	A	269	SER
1	A	306	HIS
1	B	43	GLN
1	B	189	ASN
1	B	236	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	269	SER
1	B	306	HIS
1	B	323	THR
1	C	17	GLU
1	C	189	ASN
1	C	224	HIS
1	C	233	VAL
1	C	236	ARG
1	C	269	SER
1	C	306	HIS
1	D	189	ASN
1	D	196	LYS
1	D	233	VAL
1	D	236	ARG
1	D	269	SER
1	D	306	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	C	1352	-	4,4,4	0.64	0	6,6,6	1.42	1 (16%)
3	PO4	D	1351	-	4,4,4	0.78	0	6,6,6	1.09	1 (16%)
4	PEQ	B	1352	2	8,9,9	1.81	2 (25%)	11,13,13	1.69	2 (18%)
4	PEQ	C	1351	2	8,9,9	2.16	2 (25%)	11,13,13	2.11	3 (27%)
4	PEQ	D	1352	2	8,9,9	1.98	3 (37%)	11,13,13	2.11	4 (36%)
3	PO4	A	1351	-	4,4,4	0.79	0	6,6,6	1.11	0
4	PEQ	A	1352	2	8,9,9	2.66	4 (50%)	11,13,13	1.71	3 (27%)
3	PO4	B	1353	-	4,4,4	0.57	0	6,6,6	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEQ	A	1352	2	-	5/9/9/9	-
4	PEQ	C	1351	2	-	6/9/9/9	-
4	PEQ	D	1352	2	-	6/9/9/9	-
4	PEQ	B	1352	2	-	4/9/9/9	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1352	PEQ	P-O2	4.78	1.68	1.59
4	C	1351	PEQ	P-O1P	4.68	1.65	1.50
4	A	1352	PEQ	C2-C1	3.57	1.57	1.52
4	B	1352	PEQ	P-O1P	3.42	1.61	1.50
4	D	1352	PEQ	P-O2P	3.39	1.67	1.54
4	A	1352	PEQ	P-O1P	2.95	1.60	1.50
4	C	1351	PEQ	P-O2P	2.87	1.65	1.54
4	D	1352	PEQ	P-O1P	2.43	1.58	1.50
4	A	1352	PEQ	P-O2P	2.36	1.63	1.54
4	D	1352	PEQ	P-O3P	-2.17	1.46	1.54
4	B	1352	PEQ	P-O2P	2.16	1.63	1.54

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	1351	PEQ	O2-C2-C1	-5.41	101.06	111.04
4	D	1352	PEQ	O2-C2-C1	-4.51	102.72	111.04
4	A	1352	PEQ	C3-C2-C1	3.12	114.42	109.28
4	B	1352	PEQ	O2-P-O1P	-3.11	97.37	109.39
4	D	1352	PEQ	O3P-P-O2P	3.07	119.36	107.64
4	D	1352	PEQ	C3-C2-C1	2.77	113.84	109.28
4	B	1352	PEQ	O2-C2-C1	-2.76	105.95	111.04
4	A	1352	PEQ	O3P-P-O2P	2.63	117.67	107.64
4	C	1351	PEQ	O3P-P-O2P	2.53	117.32	107.64
3	C	1352	PO4	O4-P-O2	2.33	115.45	107.97
4	A	1352	PEQ	O1-C1-C2	-2.21	114.95	121.88
4	C	1351	PEQ	O1-C1-C2	-2.19	115.00	121.88
4	D	1352	PEQ	O1-C1-C2	-2.17	115.07	121.88
3	D	1351	PO4	O4-P-O3	2.00	114.40	107.97

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1352	PEQ	O2'-C1-C2-O2
4	B	1352	PEQ	C3-C2-O2-P
4	C	1351	PEQ	O2'-C1-C2-O2
4	C	1351	PEQ	C2-O2-P-O2P
4	D	1352	PEQ	O2'-C1-C2-O2
4	D	1352	PEQ	C1-C2-O2-P
4	A	1352	PEQ	O1-C1-C2-C3
4	A	1352	PEQ	O1-C1-C2-O2
4	A	1352	PEQ	O2'-C1-C2-C3
4	B	1352	PEQ	O1-C1-C2-C3
4	C	1351	PEQ	O1-C1-C2-C3
4	C	1351	PEQ	O1-C1-C2-O2
4	C	1351	PEQ	O2'-C1-C2-C3
4	D	1352	PEQ	O1-C1-C2-C3
4	D	1352	PEQ	O1-C1-C2-O2
4	D	1352	PEQ	O2'-C1-C2-C3
4	A	1352	PEQ	C2-O2-P-O2P
4	B	1352	PEQ	O2'-C1-C2-C3
4	C	1351	PEQ	C1-C2-O2-P
4	B	1352	PEQ	O2'-C1-C2-O2
4	D	1352	PEQ	C3-C2-O2-P

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	1352	PEQ	1	0
3	A	1351	PO4	1	0
4	A	1352	PEQ	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	334/351 (95%)	-0.32	1 (0%) 94 97	23, 33, 51, 69	0
1	B	334/351 (95%)	-0.26	2 (0%) 89 93	24, 36, 50, 77	0
1	C	335/351 (95%)	-0.20	2 (0%) 89 93	23, 37, 53, 66	0
1	D	333/351 (94%)	-0.19	0 100 100	29, 42, 56, 73	0
All	All	1336/1404 (95%)	-0.24	5 (0%) 92 96	23, 37, 54, 77	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	15	VAL	3.3
1	B	293	GLN	2.9
1	B	219	HIS	2.1
1	C	148[A]	ASP	2.1
1	A	292	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	B	1353	5/5	0.68	0.42	80,96,106,106	0
3	PO4	C	1352	5/5	0.83	0.26	77,80,86,88	0
3	PO4	D	1351	5/5	0.89	0.20	58,59,60,65	0
3	PO4	A	1351	5/5	0.90	0.18	69,69,82,83	0
2	MN	C	1350	1/1	0.96	0.10	39,39,39,39	0
2	MN	D	1350	1/1	0.97	0.06	40,40,40,40	0
4	PEQ	A	1352	10/10	0.97	0.13	32,36,40,40	0
4	PEQ	B	1352	10/10	0.97	0.13	31,41,46,46	0
4	PEQ	C	1351	10/10	0.97	0.15	36,44,46,47	0
4	PEQ	D	1352	10/10	0.97	0.10	33,39,43,43	0
2	MN	A	1350	1/1	0.99	0.09	28,28,28,28	0
2	MN	B	1351	1/1	0.99	0.07	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.