



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 07:02 AM EDT

PDB ID : 4YEP
Title : L4b Domain of Human Laminin alpha-2
Authors : Toot, M.; Gat, Y.; Fass, D.
Deposited on : 2015-02-24
Resolution : 1.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1223 (1.22-1.18)
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	185	 3% 76% 24%
1	B	185	 5% 86% 13%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	A	1409	-	-	X	-
2	EDO	A	1410	-	-	X	-
2	EDO	A	1411	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7323 atoms, of which 3499 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Laminin subunit alpha-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	185	3507	1121	1804	286	286	10	0	28	0
1	B	185	3186	1030	1605	261	281	9	0	10	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1178	GLY	-	expression tag	UNP P24043
A	1179	ALA	-	expression tag	UNP P24043
A	1180	PHE	-	expression tag	UNP P24043
B	1178	GLY	-	expression tag	UNP P24043
B	1179	ALA	-	expression tag	UNP P24043
B	1180	PHE	-	expression tag	UNP P24043

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	261	Total	O	0	0
			261	261		
3	B	219	Total	O	0	0
			219	219		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Laminin subunit alpha-2



- Molecule 1: Laminin subunit alpha-2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	57.71Å 131.51Å 44.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.85 – 1.19 52.85 – 1.19	Depositor EDS
% Data completeness (in resolution range)	99.9 (52.85-1.19) 96.3 (52.85-1.19)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 1.19Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.127 , 0.162 0.128 , 0.163	Depositor DCC
R_{free} test set	7756 reflections (7.04%)	wwPDB-VP
Wilson B-factor (Å ²)	9.1	Xtrriage
Anisotropy	0.168	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 49.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	7323	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.07	4/1837 (0.2%)	0.97	5/2479 (0.2%)
1	B	1.09	5/1652 (0.3%)	1.03	6/2234 (0.3%)
All	All	1.08	9/3489 (0.3%)	1.00	11/4713 (0.2%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1352	SER	CB-OG	-10.64	1.28	1.42
1	B	1352	SER	CB-OG	-9.09	1.30	1.42
1	B	1327	GLU	CD-OE2	-7.43	1.17	1.25
1	A	1350	ARG	CD-NE	-6.36	1.35	1.46
1	B	1342	LYS	CE-NZ	6.22	1.64	1.49
1	B	1192	GLN	CG-CD	6.03	1.65	1.51
1	A	1351	GLN	CB-CG	-5.70	1.37	1.52
1	A	1191	GLU	CD-OE1	5.64	1.31	1.25
1	B	1327	GLU	CD-OE1	5.42	1.31	1.25

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1301	ARG	NE-CZ-NH2	-9.40	115.60	120.30
1	A	1350	ARG	NE-CZ-NH1	8.37	124.49	120.30
1	B	1183	ARG	NE-CZ-NH2	-8.33	116.13	120.30
1	A	1326	ARG	NE-CZ-NH2	-7.87	116.37	120.30
1	A	1289	ARG	NE-CZ-NH2	-7.14	116.73	120.30
1	B	1301	ARG	NE-CZ-NH1	7.06	123.83	120.30
1	B	1188	LEU	CB-CG-CD1	6.75	122.47	111.00
1	A	1352	SER	N-CA-CB	-6.05	101.43	110.50
1	B	1335	ASP	CB-CG-OD2	5.59	123.33	118.30
1	A	1350	ARG	NE-CZ-NH2	-5.29	117.65	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1199	ASP	CB-CG-OD1	5.18	122.96	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1703	1804	1804	48	2
1	B	1581	1605	1604	14	1
2	A	44	66	66	30	1
2	B	16	24	24	4	0
3	A	261	0	0	18	7
3	B	219	0	0	8	4
All	All	3824	3499	3498	77	10

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1188:LEU:N	2:A:1411:EDO:O2	1.90	1.05
1:A:1347[B]:ASN:OD1	3:A:1743:HOH:O	1.75	1.04
2:A:1409:EDO:H12	2:A:1411:EDO:H21	1.52	0.91
1:B:1290:HIS:ND1	3:B:1716:HOH:O	2.07	0.86
1:A:1277[B]:ARG:NH2	3:A:1501:HOH:O	2.11	0.81
1:A:1269:THR:HB	3:A:1760:HOH:O	1.79	0.81
1:B:1208:LYS:NZ	1:B:1226[B]:GLU:OE1	2.13	0.80
2:A:1409:EDO:H12	2:A:1411:EDO:C1	2.13	0.79
2:A:1409:EDO:H12	2:A:1411:EDO:C2	2.14	0.76
2:A:1409:EDO:C1	2:A:1411:EDO:H21	2.15	0.76
1:B:1194[B]:ILE:HD12	1:B:1210:ILE:HG22	1.68	0.76
2:A:1409:EDO:H12	2:A:1411:EDO:H12	1.67	0.76
2:A:1409:EDO:C2	2:A:1411:EDO:H21	2.20	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1188:LEU:H	2:A:1411:EDO:C2	2.04	0.71
1:A:1208:LYS:O	3:A:1759:HOH:O	2.07	0.71
1:A:1300:THR:OG1	3:A:1752:HOH:O	2.09	0.70
1:A:1188:LEU:HG	2:A:1411:EDO:O2	1.92	0.69
1:A:1187:THR:OG1	3:A:1756:HOH:O	2.12	0.68
1:A:1357:ILE:HB	2:A:1409:EDO:H21	1.76	0.66
2:A:1402:EDO:O2	3:A:1744:HOH:O	2.14	0.65
1:B:1301:ARG:NH1	3:B:1501:HOH:O	2.27	0.65
1:A:1277[B]:ARG:NH1	1:A:1337:HIS:ND1	2.46	0.64
1:A:1277[B]:ARG:NH2	3:A:1502:HOH:O	2.13	0.62
2:B:1401:EDO:C1	3:B:1718:HOH:O	2.46	0.62
1:B:1285:ARG:HD3	3:B:1717:HOH:O	1.99	0.62
1:B:1269:THR:HA	3:B:1716:HOH:O	2.01	0.60
1:A:1309[B]:LYS:HG3	1:A:1310:GLU:HG2	1.83	0.60
2:A:1409:EDO:C1	2:A:1411:EDO:C1	2.78	0.59
2:A:1409:EDO:H21	2:A:1411:EDO:H21	1.83	0.59
1:A:1252[C]:LYS:NZ	3:A:1504:HOH:O	2.36	0.59
1:A:1291[C]:MET:SD	3:A:1578:HOH:O	2.56	0.59
1:A:1222[B]:ASP:OD1	1:A:1225[B]:ARG:NE	2.36	0.58
1:A:1357:ILE:H	2:A:1409:EDO:C2	2.16	0.58
1:A:1357:ILE:H	2:A:1409:EDO:H21	1.68	0.57
1:A:1291[B]:MET:SD	1:A:1304[B]:ILE:CD1	2.92	0.57
1:A:1291[B]:MET:SD	1:A:1304[B]:ILE:HD11	2.44	0.56
1:A:1357:ILE:O	2:A:1411:EDO:H22	2.05	0.56
1:A:1269:THR:O	2:A:1401:EDO:O2	2.26	0.54
2:A:1409:EDO:C1	2:A:1411:EDO:H12	2.35	0.53
2:A:1410:EDO:C1	2:B:1404:EDO:HO2	2.23	0.52
1:A:1357:ILE:CB	2:A:1409:EDO:H21	2.40	0.52
1:A:1187:THR:HB	2:A:1411:EDO:O1	2.10	0.52
1:A:1189[A]:LYS:NZ	3:A:1703:HOH:O	2.14	0.51
2:B:1401:EDO:H12	3:B:1718:HOH:O	2.09	0.51
1:A:1309[B]:LYS:HG2	3:A:1688:HOH:O	2.10	0.50
1:B:1200[B]:GLU:OE2	1:B:1271:ASN:ND2	2.38	0.50
1:B:1221[A]:MET:HB3	1:B:1228:LEU:HD21	1.95	0.49
1:A:1204[B]:HIS:ND1	1:A:1230:LEU:HD21	2.27	0.49
1:A:1293:ALA:HA	2:A:1401:EDO:H22	1.95	0.49
2:A:1410:EDO:O1	2:B:1404:EDO:O2	2.19	0.49
1:A:1355:SER:HA	2:A:1409:EDO:O2	2.13	0.48
1:A:1299:LEU:HD23	2:A:1407:EDO:H21	1.96	0.48
1:B:1194[B]:ILE:CG2	3:B:1705:HOH:O	2.62	0.48
1:B:1272:PRO:HA	3:B:1716:HOH:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1265[B]:THR:HG22	3:A:1691:HOH:O	2.13	0.47
1:A:1266:GLY:HA2	1:A:1347[B]:ASN:ND2	2.32	0.45
1:A:1205[A]:THR:HG22	3:A:1758:HOH:O	2.16	0.45
1:A:1214[A]:HIS:CE1	3:A:1505:HOH:O	2.70	0.44
1:B:1226[A]:GLU:H	1:B:1226[A]:GLU:CD	2.20	0.44
1:A:1212:PHE:CE1	2:A:1403:EDO:H21	2.52	0.44
1:A:1252[B]:LYS:HD3	1:A:1303:GLU:CD	2.38	0.43
1:A:1212:PHE:HE1	2:A:1403:EDO:H21	1.83	0.43
1:B:1231:GLU:HB2	1:B:1232[B]:PRO:HA	2.01	0.43
1:A:1229:HIS:CE1	1:A:1230:LEU:HD12	2.54	0.42
1:B:1317:ASP:OD2	1:B:1319:ARG:HD3	2.18	0.42
1:A:1252[C]:LYS:HD3	1:A:1303:GLU:CD	2.39	0.42
1:A:1356:GLU:N	2:A:1409:EDO:H11	2.34	0.42
1:A:1307[B]:THR:HG22	1:A:1308:GLU:N	2.35	0.42
1:A:1214[A]:HIS:CD2	3:A:1505:HOH:O	2.73	0.42
1:A:1307[B]:THR:HG22	1:A:1308:GLU:H	1.84	0.41
1:A:1357:ILE:H	2:A:1409:EDO:C1	2.33	0.41
1:B:1200[B]:GLU:CD	1:B:1232[B]:PRO:HG2	2.40	0.41
1:A:1252[B]:LYS:NZ	3:A:1507:HOH:O	2.54	0.41
1:A:1277[B]:ARG:NH2	1:A:1284:ALA:CB	2.84	0.40
1:A:1211[A]:VAL:HG13	1:A:1213:GLN:HG3	2.04	0.40
2:A:1410:EDO:H21	3:A:1535:HOH:O	2.20	0.40

All (10) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1504:HOH:O	3:A:1592:HOH:O[2_555]	1.96	0.24
3:A:1509:HOH:O	3:A:1628:HOH:O[2_555]	1.99	0.21
3:A:1505:HOH:O	3:A:1639:HOH:O[1_554]	2.08	0.12
2:A:1408:EDO:O2	2:A:1410:EDO:O2[2_555]	2.09	0.11
1:A:1305[B]:GLU:OE2	3:B:1608:HOH:O[1_556]	2.11	0.09
1:A:1222[B]:ASP:OD2	1:B:1231:GLU:OE2[3_454]	2.14	0.06
3:A:1516:HOH:O	3:B:1533:HOH:O[3_455]	2.15	0.05
3:A:1598:HOH:O	3:A:1609:HOH:O[1_556]	2.16	0.04
3:A:1610:HOH:O	3:B:1581:HOH:O[3_454]	2.18	0.02
3:A:1717:HOH:O	3:B:1531:HOH:O[1_556]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/185 (115%)	209 (98%)	4 (2%)	0	100	100
1	B	193/185 (104%)	189 (98%)	4 (2%)	0	100	100
All	All	406/370 (110%)	398 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/162 (118%)	191 (100%)	1 (0%)	88	67
1	B	172/162 (106%)	172 (100%)	0	100	100
All	All	364/324 (112%)	363 (100%)	1 (0%)	92	79

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1316	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	B	1402	-	3,3,3	1.02	0	2,2,2	1.11	0
2	EDO	A	1411	-	3,3,3	0.76	0	2,2,2	2.13	1 (50%)
2	EDO	A	1405	-	3,3,3	0.66	0	2,2,2	0.55	0
2	EDO	A	1401	-	3,3,3	0.80	0	2,2,2	1.21	0
2	EDO	B	1401	-	3,3,3	0.39	0	2,2,2	0.34	0
2	EDO	A	1404	-	3,3,3	0.44	0	2,2,2	0.79	0
2	EDO	A	1406	-	3,3,3	0.97	0	2,2,2	0.67	0
2	EDO	B	1404	-	3,3,3	0.26	0	2,2,2	0.92	0
2	EDO	A	1402	-	3,3,3	0.67	0	2,2,2	0.44	0
2	EDO	A	1407	-	3,3,3	0.93	0	2,2,2	0.43	0
2	EDO	A	1403	-	3,3,3	0.56	0	2,2,2	0.63	0
2	EDO	A	1408	-	3,3,3	0.46	0	2,2,2	0.47	0
2	EDO	B	1403	-	3,3,3	0.45	0	2,2,2	0.28	0
2	EDO	A	1409	-	3,3,3	0.61	0	2,2,2	0.72	0
2	EDO	A	1410	-	3,3,3	0.47	0	2,2,2	0.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	1402	-	-	0/1/1/1	-
2	EDO	A	1411	-	-	0/1/1/1	-
2	EDO	A	1405	-	-	0/1/1/1	-
2	EDO	A	1401	-	-	0/1/1/1	-
2	EDO	B	1401	-	-	1/1/1/1	-
2	EDO	A	1404	-	-	0/1/1/1	-
2	EDO	A	1406	-	-	0/1/1/1	-
2	EDO	B	1404	-	-	1/1/1/1	-
2	EDO	A	1402	-	-	0/1/1/1	-
2	EDO	A	1407	-	-	0/1/1/1	-
2	EDO	A	1403	-	-	0/1/1/1	-
2	EDO	A	1408	-	-	0/1/1/1	-
2	EDO	B	1403	-	-	1/1/1/1	-
2	EDO	A	1409	-	-	1/1/1/1	-
2	EDO	A	1410	-	-	1/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1411	EDO	O2-C2-C1	2.91	132.85	111.91

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1409	EDO	O1-C1-C2-O2
2	A	1410	EDO	O1-C1-C2-O2
2	B	1401	EDO	O1-C1-C2-O2
2	B	1403	EDO	O1-C1-C2-O2
2	B	1404	EDO	O1-C1-C2-O2

There are no ring outliers.

10 monomers are involved in 33 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1411	EDO	14	0
2	A	1401	EDO	2	0
2	B	1401	EDO	2	0
2	B	1404	EDO	2	0
2	A	1402	EDO	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1407	EDO	1	0
2	A	1403	EDO	2	0
2	A	1408	EDO	0	1
2	A	1409	EDO	16	0
2	A	1410	EDO	3	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	185/185 (100%)	-0.33	5 (2%) 54 53	7, 11, 30, 55	0
1	B	185/185 (100%)	-0.35	10 (5%) 25 25	7, 11, 30, 46	0
All	All	370/370 (100%)	-0.34	15 (4%) 37 37	7, 11, 30, 55	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1267	PHE	6.5
1	B	1178	GLY	4.1
1	A	1225[A]	ARG	3.8
1	A	1178	GLY	3.6
1	B	1270	TYR	3.4
1	A	1270	TYR	3.3
1	B	1269	THR	3.3
1	A	1226	GLU	2.8
1	A	1230	LEU	2.8
1	B	1309	LYS	2.7
1	B	1268	SER	2.7
1	B	1225	ARG	2.3
1	B	1208	LYS	2.1
1	B	1266	GLY	2.1
1	B	1179	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	A	1407	4/4	0.71	0.28	23,43,67,81	0
2	EDO	B	1401	4/4	0.83	0.18	22,29,81,97	0
2	EDO	B	1403	4/4	0.85	0.14	23,28,39,39	0
2	EDO	B	1404	4/4	0.85	0.16	50,72,84,87	0
2	EDO	A	1410	4/4	0.86	0.27	22,49,83,100	0
2	EDO	A	1401	4/4	0.90	0.15	17,22,31,37	0
2	EDO	A	1403	4/4	0.93	0.15	18,24,42,42	0
2	EDO	A	1409	4/4	0.93	0.32	14,22,130,156	0
2	EDO	A	1408	4/4	0.94	0.12	19,38,99,119	0
2	EDO	B	1402	4/4	0.95	0.09	16,19,22,22	0
2	EDO	A	1411	4/4	0.95	0.23	7,38,106,109	0
2	EDO	A	1404	4/4	0.95	0.07	19,24,29,35	0
2	EDO	A	1402	4/4	0.96	0.07	12,16,28,34	0
2	EDO	A	1406	4/4	0.97	0.10	11,15,17,19	0
2	EDO	A	1405	4/4	0.98	0.06	12,16,19,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.