



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 30, 2023 – 07:20 PM JST

PDB ID : 4YKG
Title : Crystal Structure of the Alkylhydroperoxide Reductase subunit F (AhpF) with NAD⁺ from Escherichia coli
Authors : Kamariah, N.; Manimekalai, M.S.S.; Gruber, G.; Eisenhaber, F.; Eisenhaber, B.
Deposited on : 2015-03-04
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	521	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	611	-	-	-	X

2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 4305 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alkyl hydroperoxide reductase subunit F.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	521	3954	2486	681	771	16	0	1	0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	44	21	7	14	2	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
4	A	1	5	4	1	0	0
4	A	1	5	4	1	0	0
4	A	1	5	4	1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is CADMIUM ION (three-letter code: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cd	0	0
			1	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		
7	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	158	Total	O	0	0
			158	158		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alkyl hydroperoxide reductase subunit F



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	106.88Å 59.58Å 121.51Å 90.00° 111.40° 90.00°	Depositor
Resolution (Å)	27.00 – 2.40 26.40 – 2.40	Depositor EDS
% Data completeness (in resolution range)	91.7 (27.00-2.40) 91.8 (26.40-2.40)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.221 , 0.285 0.225 , 0.282	Depositor DCC
R_{free} test set	1317 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	39.2	Xtrriage
Anisotropy	0.144	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4305	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, CD, PEG, SO4, NAD, GOL, CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/4003	0.73	0/5420

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	519	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3954	0	3994	56	0
2	A	53	0	31	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	44	0	26	2	0
4	A	45	0	0	0	0
5	A	1	0	0	0	0
6	A	36	0	48	1	0
7	A	14	0	20	0	0
8	A	158	0	0	3	0
All	All	4305	0	4119	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:345[B]:CSD:SG	1:A:348:CYS:SG	2.35	1.23
1:A:345[A]:CSD:SG	1:A:348:CYS:SG	2.87	0.71
3:A:602:NAD:O2A	8:A:701:HOH:O	2.16	0.62
1:A:465:ASN:HB3	1:A:471:ILE:HD11	1.83	0.61
1:A:449:ILE:CG2	1:A:449:ILE:O	2.50	0.58
1:A:244:ARG:HD3	1:A:250:LEU:HD23	1.85	0.57
1:A:40:LEU:HB3	1:A:52:PHE:CD1	2.39	0.57
1:A:374:LEU:O	1:A:378:VAL:HG22	2.03	0.57
1:A:471:ILE:HD13	1:A:491:THR:HG21	1.87	0.56
1:A:121:PHE:O	1:A:151:HIS:HA	2.06	0.55
1:A:112:ILE:HD11	1:A:192:VAL:HG12	1.89	0.55
1:A:343:THR:HB	6:A:618:GOL:H11	1.87	0.55
1:A:32:LYS:HD3	1:A:158:THR:HG23	1.89	0.54
1:A:112:ILE:HD11	1:A:192:VAL:CG1	2.38	0.53
1:A:15:GLU:HG3	8:A:853:HOH:O	2.09	0.52
1:A:132:CYS:N	1:A:133:PRO:HD2	2.24	0.52
1:A:327:ARG:HD2	1:A:452:LEU:HD21	1.91	0.51
1:A:314:LYS:NZ	8:A:705:HOH:O	2.43	0.51
1:A:514:TYR:O	1:A:518:THR:HG23	2.11	0.50
1:A:187:THR:O	1:A:191:ILE:HG12	2.12	0.50
1:A:385:GLU:OE2	3:A:602:NAD:O2B	2.27	0.48
1:A:132:CYS:O	1:A:136:VAL:HG23	2.13	0.48
1:A:26:THR:HG22	1:A:62:LYS:HG2	1.97	0.47
1:A:481:LYS:HE3	1:A:481:LYS:HA	1.97	0.47
1:A:253:VAL:O	1:A:265:GLU:HA	2.15	0.47
1:A:328:ASN:OD1	1:A:338:ARG:NH2	2.48	0.47
1:A:143:SER:HA	1:A:149:ILE:O	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:MET:HE1	1:A:191:ILE:HG21	1.98	0.46
1:A:196:ASP:CB	1:A:199:ALA:HB2	2.46	0.46
1:A:400:LEU:HD11	1:A:406:VAL:HG11	1.98	0.46
1:A:210:ARG:O	1:A:314:LYS:NZ	2.47	0.46
1:A:24:ILE:HD12	1:A:24:ILE:N	2.31	0.45
1:A:413:GLN:OE1	1:A:432:ARG:NH1	2.43	0.45
1:A:122:GLU:HA	1:A:152:THR:O	2.16	0.45
2:A:601:FAD:H9	2:A:601:FAD:H1'1	1.79	0.45
1:A:449:ILE:O	1:A:449:ILE:HG22	2.16	0.45
1:A:345[B]:CSD:OD1	1:A:348:CYS:SG	2.74	0.45
1:A:196:ASP:CG	1:A:199:ALA:HB2	2.36	0.45
1:A:176:VAL:HG13	1:A:176:VAL:O	2.17	0.44
1:A:271:GLY:O	1:A:275:VAL:HG23	2.16	0.44
1:A:348:CYS:SG	2:A:601:FAD:C4X	3.06	0.44
1:A:496:GLN:HB2	1:A:499:ILE:HD12	1.99	0.44
1:A:365:ASN:OD1	1:A:392:ALA:HA	2.18	0.43
1:A:339:THR:OG1	1:A:340:LYS:HE2	2.19	0.43
1:A:426:GLY:HA2	1:A:442:LEU:HD12	2.02	0.42
1:A:88:THR:O	1:A:92:LEU:HD13	2.19	0.42
1:A:256:GLU:HG2	1:A:263:LYS:HE2	2.02	0.42
1:A:68:THR:OG1	1:A:69:ASN:N	2.53	0.42
1:A:28:ASP:O	1:A:29:ASP:HB2	2.20	0.42
1:A:14:LEU:HD22	1:A:17:LEU:HD11	2.01	0.42
1:A:168:VAL:O	1:A:168:VAL:HG12	2.20	0.42
1:A:313:LEU:HD22	1:A:313:LEU:N	2.34	0.41
1:A:338:ARG:O	1:A:339:THR:OG1	2.30	0.41
1:A:201:LYS:O	1:A:203:ALA:N	2.53	0.41
1:A:214:ASP:HB2	1:A:237:ARG:O	2.21	0.40
1:A:345[A]:CSD:CB	1:A:348:CYS:HG	2.28	0.40
1:A:244:ARG:CD	1:A:250:LEU:HD23	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	518/521 (99%)	484 (93%)	34 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	421/421 (100%)	402 (96%)	19 (4%)	27	44

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASP
1	A	40	LEU
1	A	83	LEU
1	A	107	SER
1	A	111	GLN
1	A	118	ASP
1	A	122	GLU
1	A	134	ASP
1	A	148	ARG
1	A	163	ILE
1	A	196	ASP
1	A	206	GLU
1	A	254	ASP
1	A	314	LYS
1	A	335	ASP
1	A	389	GLU
1	A	440	ILE
1	A	442	LEU
1	A	492	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	130	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSD	A	345[A]	-	3,7,8	1.26	0	1,8,10	0.64	0
1	CSD	A	345[B]	-	3,7,8	1.35	0	1,8,10	2.31	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	345[A]	-	-	2/2/6/8	-
1	CSD	A	345[B]	-	-	1/2/6/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	345[B]	CSD	OD1-SG-CB	2.31	109.94	105.54

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	345[A]	CSD	N-CA-CB-SG
1	A	345[A]	CSD	CA-CB-SG-OD1
1	A	345[B]	CSD	CA-CB-SG-OD1

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	345[A]	CSD	3	0
1	A	345[B]	CSD	2	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 1 is monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	A	613	-	5,5,5	0.49	0	5,5,5	0.38	0
6	GOL	A	616	-	5,5,5	0.29	0	5,5,5	0.11	0
7	PEG	A	619	-	6,6,6	0.54	0	5,5,5	0.29	0
6	GOL	A	618	-	5,5,5	0.45	0	5,5,5	0.69	0
7	PEG	A	620	-	6,6,6	0.51	0	5,5,5	0.40	0
6	GOL	A	615	-	5,5,5	0.43	0	5,5,5	0.49	0
4	SO4	A	607	-	4,4,4	0.38	0	6,6,6	0.17	0
4	SO4	A	603	-	4,4,4	0.34	0	6,6,6	0.66	0
4	SO4	A	608	-	4,4,4	0.39	0	6,6,6	0.40	0
2	FAD	A	601	-	53,58,58	0.68	0	68,89,89	0.75	3 (4%)
4	SO4	A	611	-	4,4,4	0.36	0	6,6,6	0.07	0
6	GOL	A	614	-	5,5,5	0.40	0	5,5,5	0.33	0
4	SO4	A	609	-	4,4,4	0.39	0	6,6,6	0.24	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	606	-	4,4,4	0.44	0	6,6,6	0.25	0
3	NAD	A	602	-	42,48,48	0.57	0	50,73,73	0.79	1 (2%)
6	GOL	A	617	-	5,5,5	0.49	0	5,5,5	0.36	0
4	SO4	A	610	-	4,4,4	0.35	0	6,6,6	0.08	0
4	SO4	A	604	-	4,4,4	0.45	0	6,6,6	0.34	0
4	SO4	A	605	-	4,4,4	0.31	0	6,6,6	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	613	-	-	0/4/4/4	-
6	GOL	A	615	-	-	0/4/4/4	-
6	GOL	A	614	-	-	4/4/4/4	-
6	GOL	A	616	-	-	2/4/4/4	-
3	NAD	A	602	-	-	3/26/62/62	0/5/5/5
6	GOL	A	617	-	-	2/4/4/4	-
2	FAD	A	601	-	-	3/30/50/50	0/6/6/6
6	GOL	A	618	-	-	2/4/4/4	-
7	PEG	A	619	-	-	2/4/4/4	-
7	PEG	A	620	-	-	1/4/4/4	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	FAD	O5'-C5'-C4'	-2.17	103.57	109.36
2	A	601	FAD	O4B-C1B-C2B	-2.08	103.88	106.93
3	A	602	NAD	O4B-C1B-C2B	-2.04	103.95	106.93
2	A	601	FAD	C5A-C6A-N6A	2.03	123.44	120.35

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	FAD	PA-O3P-P-O5'
6	A	614	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
6	A	614	GOL	C1-C2-C3-O3
6	A	617	GOL	O1-C1-C2-O2
6	A	617	GOL	O1-C1-C2-C3
6	A	618	GOL	C1-C2-C3-O3
6	A	614	GOL	O1-C1-C2-O2
3	A	602	NAD	O4D-C4D-C5D-O5D
3	A	602	NAD	C3D-C4D-C5D-O5D
7	A	619	PEG	O2-C3-C4-O4
6	A	616	GOL	C1-C2-C3-O3
6	A	618	GOL	O2-C2-C3-O3
7	A	619	PEG	O1-C1-C2-O2
6	A	614	GOL	O2-C2-C3-O3
6	A	616	GOL	O2-C2-C3-O3
7	A	620	PEG	O1-C1-C2-O2
3	A	602	NAD	O4B-C4B-C5B-O5B
2	A	601	FAD	P-O3P-PA-O1A
2	A	601	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	618	GOL	1	0
2	A	601	FAD	2	0
3	A	602	NAD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	520/521 (99%)	0.33	36 (6%) 16 15	18, 40, 93, 126	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	197	THR	6.2
1	A	199	ALA	5.8
1	A	117	GLY	5.1
1	A	198	GLY	4.5
1	A	200	GLU	4.2
1	A	201	LYS	4.1
1	A	164	THR	3.7
1	A	169	MET	3.6
1	A	5	ASN	3.6
1	A	163	ILE	3.4
1	A	114	HIS	3.3
1	A	521	ALA	3.2
1	A	116	ASP	3.1
1	A	131	ASN	3.1
1	A	160	GLN	3.1
1	A	196	ASP	2.9
1	A	13	TYR	2.8
1	A	99	GLY	2.8
1	A	119	PHE	2.6
1	A	59	PRO	2.6
1	A	9	GLN	2.5
1	A	103	LYS	2.5
1	A	165	ASP	2.5
1	A	73	ASN	2.5
1	A	78	PHE	2.4
1	A	202	ARG	2.4
1	A	71	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	166	ARG	2.3
1	A	12	ALA	2.3
1	A	93	ALA	2.3
1	A	143	SER	2.1
1	A	76	PRO	2.1
1	A	148	ARG	2.1
1	A	139	LEU	2.1
1	A	57	SER	2.0
1	A	27	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSD	A	345[A]	8/9	0.92	0.19	21,26,30,33	5
1	CSD	A	345[B]	8/9	0.92	0.19	22,25,30,33	5

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	A	610	5/5	0.62	0.38	131,135,136,139	0
7	PEG	A	619	7/7	0.68	0.34	65,71,74,76	0
6	GOL	A	615	6/6	0.71	0.38	64,69,73,75	0
6	GOL	A	618	6/6	0.74	0.34	46,58,61,65	0
6	GOL	A	614	6/6	0.74	0.32	57,60,62,66	0
6	GOL	A	617	6/6	0.75	0.34	53,65,69,72	0
7	PEG	A	620	7/7	0.75	0.31	59,68,74,75	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	A	611	5/5	0.78	0.44	114,118,121,122	0
6	GOL	A	613	6/6	0.80	0.24	51,57,58,58	0
6	GOL	A	616	6/6	0.81	0.29	61,66,70,73	0
4	SO4	A	607	5/5	0.85	0.27	90,90,92,94	0
4	SO4	A	609	5/5	0.88	0.38	78,79,87,88	0
4	SO4	A	608	5/5	0.89	0.32	64,72,74,80	0
4	SO4	A	606	5/5	0.93	0.31	84,87,89,95	0
4	SO4	A	604	5/5	0.94	0.15	62,63,67,69	0
3	NAD	A	602	44/44	0.95	0.12	23,35,77,96	0
2	FAD	A	601	53/53	0.97	0.11	18,25,39,40	0
5	CD	A	612	1/1	0.97	0.02	91,91,91,91	0
4	SO4	A	603	5/5	0.98	0.10	27,29,29,33	0
4	SO4	A	605	5/5	0.99	0.09	33,34,39,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.