

Full wwPDB X-ray Structure Validation Report (i)

Jun 16, 2024 – 12:34 PM EDT

PDB ID : 4ZRA

Title : CRYSTAL STRUCTURE OF MYCOBACTERIUM TUBERCULOSIS LPRG

BINDING TO TRIACYLGLYCERIDE

Authors: Martinot, A.J.; Farrow, M.; Bai, L.; Layre, E.; Cheng, T.Y.; Tsai, J.H.C.;

Iqbal, J.; Annand, J.; Sullivan, Z.; Hussain, M.; Sacchettini, J.; Moody, D.B.;

Seeliger, J.; Rubin, E.J.; TB Structural Genomics Consortium (TBSGC)

Deposited on : 2015-05-12

Resolution : 1.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.37.1buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.37.1




1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	4003 (1.86-1.82)
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)
RSRZ outliers	127900	3957 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	203	87%	7% • •
1	С	203	76%	14% 5% •



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3192 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Lipoprotein LprG.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	196	Total	С	N	О	S	0	0	0
1	A	190	1443	904	247	290	2	0	U	0
1	С	194	Total	С	N	О	S	0	0	0
1		194	1433	897	248	286	2	0	0	U

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	29	MET	-	expression tag	UNP P9WK45
A	30	HIS	-	expression tag	UNP P9WK45
A	31	HIS	-	expression tag	UNP P9WK45
A	32	HIS	-	expression tag	UNP P9WK45
A	33	HIS	-	expression tag	UNP P9WK45
A	34	HIS	-	expression tag	UNP P9WK45
A	35	HIS	-	expression tag	UNP P9WK45
С	29	MET	-	expression tag	UNP P9WK45
С	30	HIS	-	expression tag	UNP P9WK45
С	31	HIS	-	expression tag	UNP P9WK45
С	32	HIS	-	expression tag	UNP P9WK45
С	33	HIS	-	expression tag	UNP P9WK45
С	34	HIS		expression tag	UNP P9WK45
C	35	HIS	-	expression tag	UNP P9WK45

• Molecule 2 is Tripalmitoylglycerol (three-letter code: 4RF) (formula: C₅₁H₉₈O₆).





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
2	A	1	Total 57	C 51	O 6	0	0

• Molecule 3 is water.


Mo	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	165	Total O 165 165	0	0
3	С	94	Total O 94 94	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Lipoprotein LprG





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	95.68Å 71.59Å 61.89Å	Donositor
a, b, c, α , β , γ	90.00° 106.55° 90.00°	Depositor
Resolution (Å)	30.67 - 1.83	Depositor
Resolution (A)	30.65 - 1.83	EDS
% Data completeness	98.8 (30.67-1.83)	Depositor
(in resolution range)	98.8 (30.65-1.83)	EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.83 (at 1.83Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
D D.	0.204 , 0.234	Depositor
R, R_{free}	0.210 , 0.242	DCC
R_{free} test set	1764 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	27.1	Xtriage
Anisotropy	0.142	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 50.0	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3192	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 9.17% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 4RF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.98	0/1470	0.96	3/2012 (0.1%)	
1	С	0.77	0/1462	0.93	4/2000 (0.2%)	
All	All	0.88	0/2932	0.95	7/4012 (0.2%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	С	0	2
All	All	0	3

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	С	73	LEU	CA-CB-CG	7.24	131.96	115.30
1	A	40	ASP	CB-CG-OD1	5.95	123.65	118.30
1	A	122	ASP	CB-CG-OD1	5.64	123.38	118.30
1	A	167	ARG	NE-CZ-NH1	5.16	122.88	120.30
1	С	186	ALA	N-CA-C	5.11	124.81	111.00
1	С	40	ASP	CB-CG-OD1	5.08	122.88	118.30
1	С	201	ASP	CB-CG-OD2	-5.08	113.73	118.30

There are no chirality outliers.

All (3) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	228	VAL	Peptide
1	С	185	ASN	Peptide
1	С	63	THR	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1443	0	1432	11	0
1	С	1433	0	1413	30	0
2	A	57	0	98	5	0
3	A	165	0	0	1	1
3	С	94	0	0	5	0
All	All	3192	0	2943	43	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:C:174:ALA:HB1	1:C:186:ALA:HB1	1.48	0.95
1:C:64:VAL:HB	1:C:65:ASN:HA	1.59	0.83
1:C:60:MET:SD	1:C:219:MET:SD	2.78	0.81
1:C:34:HIS:NE2	3:C:301:HOH:O	2.21	0.72
1:C:223:LYS:NZ	3:C:302:HOH:O	2.24	0.71
1:A:213:SER:OG	1:A:214:GLY:N	2.28	0.67
1:C:178:ASN:HD21	1:C:186:ALA:CB	2.10	0.64
1:C:178:ASN:HD21	1:C:186:ALA:HB3	1.63	0.64
1:C:34:HIS:N	3:C:304:HOH:O	2.32	0.63
1:A:70:GLY:O	2:A:301:4RF:H69	1.99	0.62
1:C:67:LYS:O	1:C:215:ASN:ND2	2.34	0.61
1:C:64:VAL:HB	1:C:65:ASN:CA	2.30	0.60
1:C:51:GLN:HG3	1:C:201:ASP:O	2.02	0.59
1:A:228:VAL:O	1:A:230:VAL:HA	2.05	0.56
1:C:66:GLY:N	3:C:303:HOH:O	2.29	0.56
1:C:71:LEU:HD22	1:C:73:LEU:CD1	2.36	0.56

Continued on next page...



Continued from previous page...

A. 1		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ (\mathring{\rm A})$	overlap (Å)
1:C:213:SER:OG	1:C:214:GLY:N	2.34	0.56
1:A:230:VAL:HG22	1:A:231:THR:H	1.73	0.54
1:C:60:MET:HE3	1:C:80:LEU:HD21	1.89	0.53
1:A:88:THR:HG21	1:A:231:THR:HA	1.91	0.53
1:A:228:VAL:HA	1:A:229:GLN:HG3	1.92	0.53
1:C:71:LEU:HG	1:C:183:PRO:HG2	1.91	0.52
1:A:51:GLN:HG3	1:A:201:ASP:O	2.11	0.50
1:A:171:LYS:HE2	3:A:448:HOH:O	2.13	0.48
1:C:72:SER:HB2	1:C:94:THR:HG22	1.95	0.48
1:C:178:ASN:ND2	1:C:186:ALA:CB	2.78	0.47
1:A:135:VAL:HG21	2:A:301:4RF:H58	1.96	0.47
1:A:72:SER:OG	2:A:301:4RF:O23	2.10	0.46
1:C:184:PHE:O	1:C:211:ARG:NH2	2.42	0.45
1:C:60:MET:HE1	1:C:135:VAL:HG21	1.98	0.45
1:C:135:VAL:HG13	1:C:142:LEU:HG	1.98	0.44
1:C:73:LEU:HA	1:C:93:LEU:HA	1.98	0.44
1:C:60:MET:HE3	1:C:80:LEU:CD2	2.48	0.43
1:C:178:ASN:OD1	1:C:186:ALA:HB2	2.18	0.43
1:C:215:ASN:OD1	1:C:215:ASN:N	2.51	0.43
1:C:88:THR:HG22	1:C:105:VAL:HG22	2.01	0.42
2:A:301:4RF:O23	2:A:301:4RF:O42	2.38	0.42
1:C:60:MET:HG3	1:C:219:MET:HG3	2.03	0.41
1:C:62:LEU:HD11	1:C:217:VAL:CG1	2.50	0.41
1:A:230:VAL:HG13	1:A:231:THR:N	2.36	0.40
1:C:208:GLN:NE2	3:C:312:HOH:O	2.53	0.40
2:A:301:4RF:O23	2:A:301:4RF:H11	2.21	0.40
1:C:60:MET:CE	1:C:80:LEU:HD21	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
3:A:560:HOH:O	3:A:560:HOH:O[2_555]	2.10	0.10

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries



of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles		
1	A	194/203~(96%)	190 (98%)	3 (2%)	1 (0%)	29 15		
1	C	192/203~(95%)	183 (95%)	8 (4%)	1 (0%)	29 15		
All	All	386/406 (95%)	373 (97%)	11 (3%)	2 (0%)	29 15		

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	230	VAL
1	С	64	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Rotameric Outliers	
1	A	155/162~(96%)	150 (97%)	5 (3%)	39 21
1	C	153/162 (94%)	139 (91%)	14 (9%)	9 1
All	All	308/324 (95%)	289 (94%)	19 (6%)	18 4

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	ASN
1	A	93	LEU
1	A	98	SER
1	A	213	SER
1	A	229	GLN
1	С	34	HIS
1	С	67	LYS
1	С	68	ILE
1	С	72	SER
1	С	73	LEU

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	С	80	LEU
1	С	93	LEU
1	С	94	THR
1	С	98	SER
1	С	108	ASP
1	С	134	GLN
1	С	187	THR
1	С	213	SER
1	С	215	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	208	GLN
1	A	218	GLN
1	С	178	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	ol Type Cha	Chain	Res	Link	Bo	nd leng	ths	В	ond ang	les
IVIOI		Chain			Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	4RF	A	301	-	56,56,56	1.16	4 (7%)	59,59,59	1.44	6 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4RF	A	301	-	-	33/59/59/59	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(Å)
2	A	301	4RF	O21-C22	4.03	1.45	1.34
2	A	301	4RF	O18-C16	4.02	1.45	1.33
2	A	301	4RF	O40-C41	3.43	1.43	1.33
2	A	301	4RF	C15-C16	2.56	1.58	1.50

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	301	4RF	O21-C22-C24	5.24	122.80	111.50
2	A	301	4RF	O18-C16-O17	-4.22	112.94	123.59
2	A	301	4RF	O18-C16-C15	3.45	122.74	111.91
2	A	301	4RF	C39-C20-C19	-3.19	104.23	111.79
2	A	301	4RF	O40-C41-O42	-2.47	117.37	123.59
2	A	301	4RF	C45-C44-C43	-2.20	105.26	113.19

There are no chirality outliers.

All (33) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	4RF	C24-C22-O21-C20
2	A	301	4RF	O23-C22-O21-C20
2	A	301	4RF	C43-C41-O40-C39
2	A	301	4RF	C15-C16-O18-C19
2	A	301	4RF	C51-C52-C53-C54
2	A	301	4RF	O17-C16-O18-C19
2	A	301	4RF	C41-C43-C44-C45
2	A	301	4RF	C32-C33-C34-C35
2	A	301	4RF	C47-C48-C49-C50

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	301	4RF	C07-C08-C09-C10
2	A	301	4RF	C12-C13-C14-C15
2	A	301	4RF	O42-C41-O40-C39
2	A	301	4RF	C05-C06-C07-C08
2	A	301	4RF	C31-C32-C33-C34
2	A	301	4RF	C44-C45-C46-C47
2	A	301	4RF	C04-C05-C06-C07
2	A	301	4RF	C48-C49-C50-C51
2	A	301	4RF	C13-C14-C15-C16
2	A	301	4RF	C46-C47-C48-C49
2	A	301	4RF	C28-C29-C30-C31
2	A	301	4RF	C19-C20-C39-O40
2	A	301	4RF	C29-C30-C31-C32
2	A	301	4RF	C34-C35-C36-C37
2	A	301	4RF	C02-C03-C04-C05
2	A	301	4RF	C06-C07-C08-C09
2	A	301	4RF	C27-C28-C29-C30
2	A	301	4RF	C10-C11-C12-C13
2	A	301	4RF	C39-C20-O21-C22
2	A	301	4RF	C11-C12-C13-C14
2	A	301	4RF	C14-C15-C16-O18
2	A	301	4RF	C03-C04-C05-C06
2	A	301	4RF	C33-C34-C35-C36
2	A	301	4RF	C49-C50-C51-C52

There are no ring outliers.


1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	4RF	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient



equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2		$OWAB(A^2)$	Q < 0.9
1	A	196/203 (96%)	-0.38	3 (1%) 73 7	'3	16, 25, 46, 69	0
1	С	194/203~(95%)	0.01	4 (2%) 63 6	2	23, 36, 67, 93	0
All	All	390/406 (96%)	-0.19	7 (1%) 68 6	7	16, 30, 63, 93	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	229	GLN	3.9
1	С	107	PHE	3.6
1	С	70	GLY	3.5
1	С	99	ASP	2.6
1	С	71	LEU	2.5
1	A	230	VAL	2.0
1	A	36	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	4RF	A	301	57/57	0.74	0.18	33,55,81,92	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers (i)

There are no such residues in this entry.

