

Full wwPDB X-ray Structure Validation Report (i)

Jun 23, 2024 – 04:43 AM EDT

PDB ID	:	4ZUX
Title	:	SAGA DUB module Ubp8/Sgf11/Sus1/Sgf73 bound to ubiqitinated nucleo-
		some
Authors	:	Morgan, M.; Wolberger, C.
Deposited on	:	2015-05-17
Resolution	:	3.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 3.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	130704	1231 (4.04-3.60)
Ramachandran outliers	138981	$1261 \ (4.04-3.60)$
Sidechain outliers	138945	1255 (4.04-3.60)
RSRZ outliers	127900	1139 (4.04-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chair	ı	
1	А	136	% • 64%	7%	29%
1	Е	136	% • 70%	•	29%
1	K	136	65%	7%	29%
1	0	136	% • 64%	7%	29%
2	В	103	% • 75%		6% 19%
2	F	103	2% 73%		• 24%



Chain Length Quality of chain Mol .% 2L 103 74% 7% 19% 4% Р 210368% 7%• 24% 3 С 13074% 5%• 21% \mathbf{G} 3 13078% 19% • .% 3 М 13075% • • 21% 2% Q 3 13077% . . 19% .% D 1234 67% 9% • 23% .% Η 1234 72% 24% • Ν 123 4 67% 11% 23% 4 R 12366% 9% • 24% Ι 514599% • \mathbf{S} 514599% 6 J 14598% Т 6 14599% . .% U 472792% • 5% 2% Ζ . . 747293% .% • • 7472е 94% 2% j 4727• 5% 93% 6% V 8 96 • 7% 91% 6% 8 96 • 6% \mathbf{a} 91% 5% f 8 96 11%86% 6% k 96 8 88% • 10% .% W 99 9 89% 9% • 9 99 b 86% 10% • 9 99 9% g 90% •

Continued from previous page...



Conti	nued fron	n previous	page	
Mol	Chain	Length	Quality of chain	
9	1	99	2% 91%	• 6%
10	X	76	99%	
10	с	76	3% 	
10	h	76	99%	
10	m	76	100%	
11	Y	104	% 75% •	21%
11	d	104	4% 80% 6%	14%
11	i	104	80% 6%	14%
11	n	104	6% 80% ·	17%



2 Entry composition (i)

There are 12 unique types of molecules in this entry. The entry contains 49079 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	07	Total	С	Ν	0	\mathbf{S}	0	0	0
	A	91	802	506	155	138	3	0	0	0
1	Б	07	Total	С	Ν	0	S	0	0	0
	E	97	801	504	155	139	3			0
1	V	07	Total	С	Ν	0	S	0	0	0
	97	802	506	155	138	3	0	0	0	
1 O	97	Total	С	Ν	0	S	0	0	0	
		801	504	155	139	3			0	

• Molecule 1 is a protein called Histone H3.2.

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	102	ALA	GLY	conflict	UNP P84233
Е	102	ALA	GLY	conflict	UNP P84233
K	102	ALA	GLY	conflict	UNP P84233
0	102	ALA	GLY	conflict	UNP P84233

• Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
0	В	83	Total	С	Ν	0	S	0	0	0
	D	00	662	418	129	114	1	0	0	0
0	Б	79	Total	С	Ν	0	S	0	0	0
	Г	10	619	391	120	107	1	0	0	0
0	т	00	Total	С	Ν	0	S	0	0	0
		00	662	418	129	114	1	0	0	0
0	D	D 70	Total	С	Ν	0	S	0	0	0
	Ľ	10	619	391	120	107	1		0	U

• Molecule 3 is a protein called Histone H2A type 1.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	С	103	Total	С	Ν	Ο	0	0	0
5	U	105	795	501	155	139	0		0
2	С	105	Total	С	Ν	Ο	0	0	0
J	G	105	809	510	158	141			
2	М	103	Total	С	Ν	Ο	0	0	0
J	5 M		795	501	155	139	0	0	0
2	2 0	105	Total	С	Ν	Ο	0	0	0
3 Q	105	809	510	158	141	0	U	U	

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	99	ARG	GLY	conflict	UNP P06897
С	123	SER	ALA	conflict	UNP P06897
G	99	ARG	GLY	conflict	UNP P06897
G	123	SER	ALA	conflict	UNP P06897
М	99	ARG	GLY	conflict	UNP P06897
М	123	SER	ALA	conflict	UNP P06897
Q	99	ARG	GLY	conflict	UNP P06897
Q	123	SER	ALA	conflict	UNP P06897

• Molecule 4 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	Л	05	Total	С	Ν	Ο	S	0	0	0
4	D	90	745	469	134	140	2	0		0
4	Ц	03	Total	С	Ν	0	S	0	0	0
4	11	90	726	457	130	137	2			0
4	N	05	Total	С	Ν	0	S	0	0	0
4	4 IN	95	745	469	134	140	2		0	0
4	4 D	D 02	Total	С	Ν	0	S	0	0	0
4 K	93	726	457	130	137	2	U	U	U	

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	0	MET	-	initiating methionine	UNP P02281
D	29	THR	SER	conflict	UNP P02281
Н	0	MET	-	initiating methionine	UNP P02281
Н	29	THR	SER	conflict	UNP P02281
N	0	MET	-	initiating methionine	UNP P02281
N	29	THR	SER	conflict	UNP P02281
R	0	MET	-	initiating methionine	UNP P02281



Chain	Residue	Modelled	Actual	Comment	Reference
R	29	THR	SER	conflict	UNP P02281

• Molecule 5 is a DNA chain called DNA (145-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	Ι	145	Total 2952	C 1404	N 537	O 867	Р 144	0	0	0
5	S	145	Total 2952	C 1404	N 537	O 867	Р 144	0	0	0

• Molecule 6 is a DNA chain called DNA (145-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	J	145	Total	С	Ν	0	Р	0	0	0
	0 5	110	2987	1416	558	869	144	Ŭ	Ŭ	Ŭ
6	т	145	Total	С	Ν	0	Р	0	0	0
0	0 1	140	2987	1416	558	869	144	0	0	0

• Molecule 7 is a protein called Ubiquitin carboxyl-terminal hydrolase 8.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
7 11		447	Total	С	Ν	0	\mathbf{S}	0	0	0
1	U	447	3569	2264	609	661	35	0	0	0
7	7	451	Total	С	Ν	0	S	0	0	0
1		451	3600	2283	615	667	35	0	0	U
7	0	452	Total	С	Ν	0	S	0	0	0
(е	400	3617	2292	618	672	35	0	0	0
7	;	447	Total	С	Ν	0	S	0	0	0
(J	447	3566	2262	609	660	35	U	U	

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
U	0	ALA	-	expression tag	UNP P50102
U	146	ALA	CYS	engineered mutation	UNP P50102
Z	0	ALA	-	expression tag	UNP P50102
Z	146	ALA	CYS	engineered mutation	UNP P50102
e	0	ALA	-	expression tag	UNP P50102
e	146	ALA	CYS	engineered mutation	UNP P50102
j	0	ALA	-	expression tag	UNP P50102
j	146	ALA	CYS	engineered mutation	UNP P50102



Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
0 V		80	Total	С	Ν	0	\mathbf{S}	0	0	0
0	v	69	719	451	118	148	2	0	0	0
0	0	00	Total	С	Ν	0	S	0	0	0
0	a	90	730	457	119	152	2	0	0	0
0	t	95	Total	С	Ν	0	S	0	0	0
0	1	00	690	432	112	144	2	0	0	0
0	l,	86	Total	С	Ν	0	S	0	0	0
8 K	80	695	435	115	143	2			U	

• Molecule 8 is a protein called Transcription and mRNA export factor SUS1.

• Molecule 9 is a protein called SAGA-associated factor 11.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
		00	Total	С	Ν	0	S	0	0	0
9	vv	90	718	440	133	142	3	0	0	0
0	h	80	Total	С	Ν	0	S	0	0	0
9	D	89	710	434	132	141	3	0	0	0
0	C.	00	Total	С	Ν	0	S	0	0	0
9	g	90	718	440	133	142	3	0	0	0
0	1	03	Total	С	Ν	0	S	0	0	0
9 1		30	739	452	136	148	3			U

• Molecule 10 is a protein called Polyubiquitin-B.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
10 V		76	Total	С	Ν	0	S	0	0	0
10	Λ	70	601	378	105	117	1	0	0	0
10	0	76	Total	С	Ν	0	S	0	0	0
10	C	10	601	378	105	117	1	0	0	0
10	h	76	Total	С	Ν	0	S	0	0	0
10	11	70	601	378	105	117	1	0	0	0
10	m	76	Total	С	Ν	0	S	0	0	0
10	III	70	601	378	105	117	1	0	0	U

• Molecule 11 is a protein called SAGA-associated factor 73.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11 V		20	Total	С	Ν	0	S	0	0	0
	1	02	663	419	112	127	5	0	0	0
11	d	20	Total	С	Ν	0	S	0	0	0
	u	89	710	447	120	138	5	0	0	0
11	;	80	Total	С	Ν	0	S	0	0	0
11 1	1	89	710	447	120	138	5	0	0	0



Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
11	n	86	Total 693	C 437	N 117	0 134	${ m S}{ m 5}$	0	0	0

• Molecule 12 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	U	6	Total Zn 6 6	0	0
12	W	1	Total Zn 1 1	0	0
12	Y	1	Total Zn 1 1	0	0
12	Ζ	6	Total Zn 6 6	0	0
12	b	1	Total Zn 1 1	0	0
12	d	1	Total Zn 1 1	0	0
12	е	6	Total Zn 6 6	0	0
12	g	1	Total Zn 1 1	0	0
12	i	1	Total Zn 1 1	0	0
12	j	6	Total Zn 6 6	0	0
12	1	1	Total Zn 1 1	0	0
12	n	1	Total Zn 1 1	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Histone H3.2



• Molecule 2: Histone H4			
Chain F:	73%	•	24%
MET SER SER ARG GLY GLY GLY GLY GLY GLY GLY ALA ALA ARG ARG	LYS VAL LEU LEU LEU ASP ASP ASP ASP 126 D68 D68 D68	61102	
• Molecule 2: Histone H4			
Chain L:	74%	7%	19%
MET SER ARG ARG ARG CLY CLY CLY CLY CLY CLY CLY CLY CLY CLY	кама 126 134 134 847 847 173	891 892 6102	
• Molecule 2: Histone H4		_	
Chain P:	68%	7% •	24%
MET SER SER ARG CLY CLY CLY CLY CLY CLY CLY ALA ALA ALA ALA ALA ALA ALA	LYS VAL LEU LEU ARP N25 126 129 129 134 134 134	173 1985 1985 1992 1998 1998 1998 1991 1991	
• Molecule 3: Histone H2A	A type 1		
Chain C:	74%	5%•	21%
MET SER GLY ARG CLY CLYS CLYS CLYS ALA ALA ALA ALA ALA ALA ALA ALA ALA AL	T59 E64 E64 S113 S113 S113 V114 LYS LYS CLU SEU	SER LYS SER LYS SER LYS	
• Molecule 3: Histone H2A	A type 1		
Chain G:	78%		19%
MET SER GLY GLY ARG CLY CLYS GLN CLYS CLY CLYS LYS LYS LYS LYS LYS M14 ARG A14 A14 A14 A14 A14 A14 A14 A14 A14 A17 A17 A17 A17 A17 A17 A17 A17 A17 A17	R88 V114 V114 LYS LYS CLU SER SER SER SER ALA	SER	
• Molecule 3: Histone H2A	A type 1		
Chain M:	75%		21%
MET SER SER ARC CLY CLY CLYS CLY CLYS ALA ALA ALA ALA ALA ALA ALA CL23 ALA	R29 E64 E91 S113 S113 S113 K118 K118 LYS T4R	GLU SER SER LYS ALA LYS SER LYS SER LYS	
• Molecule 3: Histone H2A	A type 1		
Chain Q:	77%	•••	19%
MET SER GLY GLY ARG CLY CLY CLY CLY CLY CLY ARG ALA A14 LYS A14 LYS CLY CLY CLY CLY CLY CLY CLY CLY CLY CLY	R81 8113 V114 V114 V118 K118 LYS CLU SER SER SER SER SER SER	LIYS SER LYS LYS	
	W O PROT	R L D W I D E PDB EIN DATA BANK	

• Molecule 4: Histor	ne H2B 1.1		
Chain D:	67%	9% •	23%
MET LYS SER ALA ALA ALA ALA ALA CYS CLYS CLYS SER LYS ALA ALA	VAL THR LYS LYS GLN GLN LYS ASP ASP CLYS LYS K31 ARG K31 Y39	D48 149 177 1785 1785 1785 198 198 1910	S109 T116 T119 K122
• Molecule 4: Histor	ne H2B 1.1		
Chain H:	72%		24%
MGT ALA LYS SER SER ALA PRO PRO CLYS CLYS CLYS SER LYS SER LYS ALA	VAL THR LYS LYS THR THR GIN LYS LYS LYS LYS LYS LYS LYS T29 ARG T29 ARG T29 ARG T29 ARG T29 ARG	186 187 187 188 188 188 183 6101 193 6101 173 121	
• Molecule 4: Histor	ne H2B 1.1		
Chain N:	67%	11%	23%
MET ALA LYS SER ALA ALA ALA PRO LYS CLYS CLY SER LYS ALA	VAL LYS LYS LYS LYS LYS LYS LYS LYS VAS VAS VAS	N60 177 185 185 193 193 193 193	6101 8109 1116 1119 1119 K122
• Molecule 4: Histor	ne H2B 1.1		
Chain R:	66%	9% •	24%
MET LYS SER PRO PRO LYS CLYS SER LYS ALA ALA ALA	VAL LYS LYS THR THR THR CVS CLYS CLYS ASP ASP CLYS LYS CLYS CLYS CLYS CLYS ASP AC CLYS CLYS CLYS CLYS CLYS CLYS CLYS CLY	749 L77 193 197 197	8109 T116 T119 8120 A121 LYS
• Molecule 5: DNA	(145-MER)		
Chain I:	99%		•
A-72 G-53 C-2 T72			
• Molecule 5: DNA	(145-MER)		
Chain S:	99%		
42 - <mark>23 - 2</mark> 3			
• Molecule 6: DNA	(145-MER)		
Chain L			
Unani J:	98%		•
A-72 T-63 A-62 C-61 C-61 T72			
• Molecule 6: DNA	(145-MER)		





• Molecule 8: Transcription and mRNA export factor SUS1





• Molecule 9: SAGA-associated factor 11



Chain l: 91% • 6%	-
AET HIR E3 567 567 C92 ALA ARG ARG ARG	
• Molecule 10: Polyubiquitin-B	
Chain X: 99%	•
• Molecule 10: Polyubiquitin-B	
Chain c: 99%	•
• Molecule 10: Polyubiquitin-B	
Chain h: 99%	•
M1 13 16 16 16 16 16 16 16 16 16 16 16 16 16	
• Molecule 10: Polyubiquitin-B	
Chain m: 100%	-
There are no outlier residues recorded for this chain.	
• Molecule 11: SAGA-associated factor 73	
Chain Y: 75% · 21%	-
MET ARG SER ALA ALA ALA ALA ALA ALA CGN SER CGN CGN ALA ALA ALA CGN CGN CGN CGN CGN CGN CGN CGN CGN CGN	
• Molecule 11: SAGA-associated factor 73	
Chain d: 80% 6% 14%	-
MET ARG SER CJ CJ CJ CJ CJ CJ CJ CJ CJ CJ CJ CJ CJ	

• Molecule 11: SAGA-associated factor 73







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	178.80Å 179.17 Å 353.82 Å	Deperitor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	49.50 - 3.82	Depositor
Resolution (A)	$49.50 \ - \ 3.69$	EDS
% Data completeness	98.9 (49.50-3.82)	Depositor
(in resolution range)	99.3 (49.50 - 3.69)	EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.47 (at 3.67 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
P. P.	0.234 , 0.256	Depositor
n, n_{free}	0.232 , 0.260	DCC
R_{free} test set	1997 reflections (1.64%)	wwPDB-VP
Wilson B-factor $(Å^2)$	168.1	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.27, 121.5	EDS
L-test for $twinning^2$	$< L >=0.37, < L^2>=0.20$	Xtriage
Estimated twinning fraction	0.228 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	49079	wwPDB-VP
Average B, all atoms $(Å^2)$	197.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 10.69% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond lengths		Bond angles		
WIOI	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.27	0/814	0.50	0/1092	
1	Е	0.24	0/812	0.47	0/1088	
1	K	0.26	0/814	0.50	0/1092	
1	0	0.27	0/812	0.55	0/1088	
2	В	0.28	0/669	0.59	0/894	
2	F	0.32	0/626	0.55	0/837	
2	L	0.30	0/669	0.57	0/894	
2	Р	0.28	0/626	0.59	0/837	
3	С	0.25	0/805	0.53	0/1088	
3	G	0.26	0/819	0.55	0/1106	
3	М	0.25	0/805	0.57	1/1088~(0.1%)	
3	Q	0.25	0/819	0.54	0/1106	
4	D	0.28	0/756	0.56	0/1015	
4	Н	0.26	0/737	0.55	0/993	
4	N	0.26	0/756	0.54	0/1015	
4	R	0.29	0/737	0.63	1/993~(0.1%)	
5	Ι	0.63	0/3308	1.05	2/5099~(0.0%)	
5	S	0.64	0/3308	1.05	2/5099~(0.0%)	
6	J	0.62	0/3354	1.05	5/5180~(0.1%)	
6	Т	0.62	0/3354	1.06	3/5180~(0.1%)	
7	U	0.26	0/3647	0.49	0/4916	
7	Ζ	0.25	0/3678	0.49	0/4955	
7	е	0.25	0/3695	0.50	0/4978	
7	j	0.25	0/3643	0.50	0/4908	
8	V	0.23	0/725	0.46	0/977	
8	a	0.24	0/735	0.48	0/990	
8	f	0.29	0/695	0.57	0/936	
8	k	0.25	0/701	0.44	0/944	
9	W	0.27	0/726	0.52	0/982	
9	b	0.26	0/718	0.52	0/971	
9	g	0.30	0/726	0.55	0/982	
9	1	0.27	0/747	0.54	0/1011	



Mol Chain		Bond lengths		В	ond angles
	Ullalli	RMSZ	# Z > 5	RMSZ	# Z > 5
10	Х	0.22	0/607	0.47	0/816
10	с	0.22	0/607	0.44	0/816
10	h	0.23	0/607	0.48	0/816
10	m	0.22	0/607	0.45	0/816
11	Y	0.25	0/678	0.61	0/915
11	d	0.25	0/725	0.56	0/978
11	i	0.26	0/725	0.58	0/978
11	n	0.25	0/707	0.55	0/953
All	All	0.39	0/51099	0.72	14/71422~(0.0%)

There are no bond length outliers.

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
3	М	81	ARG	NE-CZ-NH1	-7.82	116.39	120.30
4	R	83	ARG	NE-CZ-NH1	-6.16	117.22	120.30
5	Ι	-53	DG	OP2-P-O3'	6.14	118.71	105.20
5	Ι	-2	DC	O4'-C1'-N1	5.95	112.17	108.00
5	S	-53	DG	OP2-P-O3'	5.87	118.11	105.20
6	J	-61	DT	O4'-C1'-N1	5.34	111.73	108.00
6	Т	20	DG	C4'-C3'-C2'	-5.33	98.30	103.10
6	Т	20	DG	C3'-C2'-C1'	-5.27	96.17	102.50
5	S	-53	DG	P-O3'-C3'	5.25	126.00	119.70
6	Т	-63	DT	O4'-C1'-N1	5.23	111.66	108.00
6	J	20	DG	C3'-C2'-C1'	-5.17	96.29	102.50
6	J	-63	DT	C1'-O4'-C4'	-5.17	104.93	110.10
6	J	-63	DT	O4'-C1'-N1	5.13	111.59	108.00
6	J	20	DG	C4'-C3'-C2'	-5.00	98.60	103.10

All (14) bond angle outliers are listed below:

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	95/136~(70%)	84 (88%)	8 (8%)	3(3%)	4	32
1	Ε	95/136~(70%)	86 (90%)	9 (10%)	0	100	100
1	Κ	95/136~(70%)	84 (88%)	8 (8%)	3~(3%)	4	32
1	Ο	95/136~(70%)	84 (88%)	8 (8%)	3(3%)	4	32
2	В	81/103~(79%)	71 (88%)	8 (10%)	2(2%)	5	36
2	F	76/103~(74%)	69 (91%)	7 (9%)	0	100	100
2	L	81/103~(79%)	71 (88%)	7 (9%)	3 (4%)	3	29
2	Р	76/103~(74%)	65 (86%)	8 (10%)	3 (4%)	3	28
3	С	101/130~(78%)	89 (88%)	9 (9%)	3(3%)	4	33
3	G	103/130~(79%)	91 (88%)	12 (12%)	0	100	100
3	М	101/130~(78%)	90 (89%)	8 (8%)	3(3%)	4	33
3	Q	103/130~(79%)	91 (88%)	9 (9%)	3 (3%)	4	34
4	D	93/123~(76%)	84 (90%)	7 (8%)	2(2%)	6	38
4	Н	91/123~(74%)	81 (89%)	9 (10%)	1 (1%)	14	51
4	Ν	93/123~(76%)	83 (89%)	8 (9%)	2(2%)	6	38
4	R	91/123~(74%)	82 (90%)	7 (8%)	2(2%)	6	38
7	U	439/472~(93%)	430 (98%)	9 (2%)	0	100	100
7	Ζ	443/472~(94%)	434 (98%)	9 (2%)	0	100	100
7	е	445/472~(94%)	435 (98%)	9 (2%)	1 (0%)	47	78
7	j	437/472~(93%)	426 (98%)	11 (2%)	0	100	100
8	V	87/96~(91%)	87 (100%)	0	0	100	100
8	a	86/96~(90%)	86 (100%)	0	0	100	100
8	f	81/96~(84%)	81 (100%)	0	0	100	100
8	k	84/96~(88%)	84 (100%)	0	0	100	100
9	W	88/99~(89%)	86 (98%)	2 (2%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
9	b	87/99~(88%)	84 (97%)	2(2%)	1 (1%)	14 51
9	g	88/99~(89%)	86~(98%)	2(2%)	0	100 100
9	1	91/99~(92%)	89~(98%)	2(2%)	0	100 100
10	Х	74/76~(97%)	73~(99%)	1 (1%)	0	100 100
10	с	74/76~(97%)	73~(99%)	1 (1%)	0	100 100
10	h	74/76~(97%)	72~(97%)	2(3%)	0	100 100
10	m	74/76~(97%)	73~(99%)	1 (1%)	0	100 100
11	Y	78/104~(75%)	75~(96%)	3~(4%)	0	100 100
11	d	85/104~(82%)	80 (94%)	4(5%)	1 (1%)	13 49
11	i	85/104~(82%)	82~(96%)	3~(4%)	0	100 100
11	n	82/104~(79%)	79~(96%)	3 (4%)	0	100 100
All	All	4552/5356~(85%)	4320 (95%)	196 (4%)	36 (1%)	19 56

All (36) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	73	GLU
2	В	29	ILE
4	D	101	GLY
1	Κ	73	GLU
2	L	29	ILE
4	Ν	101	GLY
4	R	100	PRO
9	b	76	CYS
11	d	47	ASP
4	Н	101	GLY
1	Κ	77	ASP
4	Ν	109	SER
1	0	73	GLU
1	0	77	ASP
2	Р	29	ILE
1	А	77	ASP
4	D	109	SER
4	R	109	SER
7	е	292	CYS
2	В	26	ILE
3	С	64	GLU
3	М	64	GLU



Mol	Chain	Res	Type
3	Q	64	GLU
1	А	74	ILE
1	Κ	74	ILE
2	L	26	ILE
1	0	74	ILE
3	С	113	SER
3	М	113	SER
3	Q	113	SER
2	Р	26	ILE
3	Q	114	VAL
3	С	114	VAL
2	P	34	ILE
2	L	34	ILE
3	М	114	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	А	85/111~(77%)	78~(92%)	7 (8%)	11	40
1	Е	84/111~(76%)	82~(98%)	2(2%)	49	71
1	Κ	85/111~(77%)	79~(93%)	6~(7%)	14	45
1	Ο	84/111 (76%)	77 (92%)	7 (8%)	11	40
2	В	68/79~(86%)	64 (94%)	4 (6%)	19	51
2	F	63/79~(80%)	60~(95%)	3~(5%)	25	55
2	L	68/79~(86%)	64 (94%)	4 (6%)	19	51
2	Р	63/79~(80%)	57 (90%)	6 (10%)	8	34
3	С	82/102~(80%)	77 (94%)	5~(6%)	18	50
3	G	83/102 (81%)	79~(95%)	4 (5%)	25	55
3	М	82/102~(80%)	78~(95%)	4(5%)	25	55
3	Q	83/102 (81%)	80 (96%)	3 (4%)	35	63
4	D	81/103~(79%)	70~(86%)	11 (14%)	3	22



4ZUX	
-	

Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
4	Н	79/103~(77%)	75~(95%)	4 (5%)	24	54
4	Ν	81/103~(79%)	70~(86%)	11 (14%)	3	22
4	R	79/103~(77%)	69~(87%)	10 (13%)	4	23
7	U	406/428~(95%)	391~(96%)	15~(4%)	34	61
7	Z	409/428~(96%)	398~(97%)	11 (3%)	44	68
7	е	411/428~(96%)	401 (98%)	10 (2%)	49	71
7	j	405/428~(95%)	395~(98%)	10 (2%)	47	70
8	V	84/91~(92%)	82~(98%)	2(2%)	49	71
8	a	86/91~(94%)	83~(96%)	3(4%)	36	63
8	f	81/91~(89%)	79~(98%)	2(2%)	47	70
8	k	81/91~(89%)	79~(98%)	2(2%)	47	70
9	W	82/89~(92%)	80~(98%)	2(2%)	49	71
9	b	81/89~(91%)	78~(96%)	3~(4%)	34	61
9	g	82/89~(92%)	81 (99%)	1 (1%)	71	84
9	1	84/89~(94%)	81~(96%)	3~(4%)	35	63
10	Х	68/68~(100%)	67~(98%)	1 (2%)	65	80
10	с	68/68~(100%)	67~(98%)	1 (2%)	65	80
10	h	68/68~(100%)	67~(98%)	1 (2%)	65	80
10	m	68/68~(100%)	68~(100%)	0	100	100
11	Y	77/90~(86%)	73~(95%)	4(5%)	23	54
11	d	81/90~(90%)	76~(94%)	5(6%)	18	49
11	i	81/90 (90%)	75~(93%)	6(7%)	13	44
11	n	79/90 (88%)	$76 \ (96\%)$	3(4%)	33	60
All	All	4132/4644 (89%)	3956 (96%)	176 (4%)	29	58

All (176) residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	А	48	LEU
1	А	49	ARG
1	А	59	GLU
1	А	63	ARG
1	А	65	LEU
1	А	105	GLU



Mol	Chain	Res	Type
1	А	129	ARG
2	В	47	SER
2	В	73	THR
2	В	91	LYS
2	В	92	ARG
3	С	29	ARG
3	С	59	THR
3	С	81	ARG
3	С	114	VAL
3	С	118	LYS
4	D	31	LYS
4	D	39	TYR
4	D	48	ASP
4	D	49	THR
4	D	77	LEU
4	D	84	SER
4	D	85	THR
4	D	98	LEU
4	D	109	SER
4	D	116	THR
4	D	119	THR
1	Е	59	GLU
1	Ε	129	ARG
2	F	26	ILE
2	F	73	THR
2	F	92	ARG
3	G	72	ASP
3	G	74	LYS
3	G	88	ARG
3	G	114	VAL
4	Н	49	THR
4	Н	83	ARG
4	Н	88	SER
4	Н	93	THR
1	K	49	ARG
1	K	59	GLU
1	K	63	ARG
1	K	65	LEU
1	K	105	GLU
1	K	129	ARG
2	L	47	SER
2	L	73	THR



Mol	Chain	Res	Type
2	L	91	LYS
2	L	92	ARG
3	М	29	ARG
3	М	81	ARG
3	М	91	GLU
3	М	114	VAL
4	Ν	39	TYR
4	Ν	48	ASP
4	Ν	49	THR
4	Ν	60	ASN
4	N	77	LEU
4	N	84	SER
4	Ν	85	THR
4	N	93	THR
4	Ν	98	LEU
4	N	116	THR
4	Ν	119	THR
1	0	59	GLU
1	0	63	ARG
1	0	65	LEU
1	0	90	MET
1	0	105	GLU
1	0	123	ASP
1	0	134	ARG
2	Р	26	ILE
2	Р	47	SER
2	Р	73	THR
2	Р	85	ASP
2	P	92	ARG
2	Р	95	ARG
3	Q	81	ARG
3	Q	114	VAL
3	Q	118	LYS
4	R	39	TYR
4	R	48	ASP
4	R	49	THR
4	R	77	LEU
4	R	84	SER
4	R	93	THR
4	R	97	LEU
4	R	109	SER
4	R	116	THR



Mol	Chain	Res	Type
4	R	119	THR
7	U	1	MET
7	U	83	HIS
7	U	98	LYS
7	U	116	LYS
7	U	124	LYS
7	U	142	MET
7	U	156	ILE
7	U	176	VAL
7	U	192	HIS
7	U	219	LEU
7	U	228	GLN
7	U	296	SER
7	U	365	PHE
7	U	378	ASP
7	U	425	GLU
8	V	28	GLU
8	V	68	LYS
9	W	67	SER
9	W	85	LEU
10	Х	39	ASP
11	Y	18	GLU
11	Y	29	ASN
11	Y	87	LEU
11	Y	96	ASN
7	Ζ	1	MET
7	Ζ	98	LYS
7	Ζ	116	LYS
7	Ζ	142	MET
7	Ζ	156	ILE
7	Ζ	176	VAL
7	Ζ	219	LEU
7	Ζ	228	GLN
7	Ζ	365	PHE
7	Ζ	378	ASP
7	Ζ	397	LYS
8	a	28	GLU
8	a	68	LYS
8	a	78	ARG
9	b	67	SER
9	b	75	ASN
0	h	85	LEII



Mol	Chain	Res	Type
10	с	39	ASP
11	d	5	ASP
11	d	18	GLU
11	d	29	ASN
11	d	87	LEU
11	d	96	ASN
7	е	98	LYS
7	е	116	LYS
7	е	142	MET
7	е	156	ILE
7	е	176	VAL
7	е	219	LEU
7	е	228	GLN
7	е	292	CYS
7	е	365	PHE
7	е	397	LYS
8	f	28	GLU
8	f	68	LYS
9	g	85	LEU
10	h	39	ASP
11	i	5	ASP
11	i	18	GLU
11	i	29	ASN
11	i	87	LEU
11	i	92	ASP
11	i	96	ASN
7	j	98	LYS
7	j	142	MET
7	j	156	ILE
7	j	176	VAL
7	j	219	LEU
7	j	228	GLN
7	j	296	SER
7	j	365	PHE
7	j	378	ASP
7	j	425	GLU
8	k	28	GLU
8	k	68	LYS
9	1	67	SER
9	1	85	LEU
9	1	92	CYS
11	n	5	ASP



Continued from previous page...

Mol	Chain	Res	Type
11	n	87	LEU
11	n	96	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
2	В	75	HIS
2	L	75	HIS
4	N	60	ASN
4	N	64	ASN
1	0	39	HIS
3	Q	112	GLN
7	U	53	ASN
7	U	83	HIS
7	U	90	ASN
7	Ζ	88	ASN
7	Ζ	90	ASN
7	Ζ	328	GLN
9	b	75	ASN
7	е	348	GLN
11	i	61	ASN
9	1	88	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 32 ligands modelled in this entry, 32 are monoatomic - leaving 0 for Mogul analysis.



There are no bond length outliers. There are no bond angle outliers. There are no chirality outliers. There are no torsion outliers. There are no ring outliers. No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	# RSRZ > 2	$\mathrm{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	А	97/136~(71%)	-0.09	1 (1%) 82 76	134,166,198,212	0
1	Е	97/136~(71%)	-0.14	1 (1%) 82 76	139, 172, 213, 249	0
1	K	97/136~(71%)	-0.15	0 100 100	146, 175, 209, 216	0
1	Ο	97/136~(71%)	-0.14	2 (2%) 63 55	141, 177, 219, 249	0
2	В	83/103~(80%)	-0.18	1 (1%) 79 72	124, 155, 185, 198	0
2	F	78/103~(75%)	-0.04	2 (2%) 56 46	124, 166, 189, 201	0
2	L	83/103~(80%)	-0.10	1 (1%) 79 72	131, 152, 192, 204	0
2	Р	78/103~(75%)	0.13	4 (5%) 28 24	153, 183, 219, 230	0
3	С	103/130~(79%)	-0.12	0 100 100	128, 162, 187, 197	0
3	G	105/130~(80%)	-0.01	0 100 100	148, 177, 198, 207	0
3	М	103/130~(79%)	-0.09	1 (0%) 82 76	143, 168, 195, 216	0
3	Q	105/130~(80%)	-0.05	2 (1%) 66 59	141, 179, 205, 222	0
4	D	95/123~(77%)	-0.18	1 (1%) 80 74	125, 155, 201, 243	0
4	Н	93/123~(75%)	-0.21	1 (1%) 80 74	145, 172, 197, 209	0
4	Ν	95/123~(77%)	-0.17	0 100 100	137, 165, 190, 239	0
4	R	93/123~(75%)	-0.22	0 100 100	149, 172, 196, 207	0
5	Ι	145/145~(100%)	-0.85	0 100 100	176, 210, 241, 255	0
5	S	145/145~(100%)	-0.88	0 100 100	177, 208, 241, 257	0
6	J	145/145~(100%)	-0.87	0 100 100	169, 211, 243, 269	0
6	Т	145/145~(100%)	-0.87	0 100 100	177, 211, 243, 264	0
7	U	447/472~(94%)	-0.12	6 (1%) 77 70	133, 183, 235, 285	0
7	Z	$45\overline{1/472} \ (95\%)$	0.00	10 (2%) 62 53	161, 205, 261, 302	0
7	е	453/472~(95%)	-0.14	7 (1%) 73 66	148, 192, 244, 276	0
7	j	447/472 (94%)	-0.16	10 (2%) 62 53	146, 189, 232, 286	0



Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q<0.9
8	V	89/96~(92%)	-0.09	6 (6%) 17 13	160, 196, 260, 275	0
8	a	90/96~(93%)	0.10	6 (6%) 17 13	181, 238, 267, 283	0
8	f	85/96~(88%)	0.26	5 (5%) 22 17	196, 227, 267, 298	0
8	k	86/96~(89%)	-0.04	6 (6%) 16 12	188, 217, 249, 277	1 (1%)
9	W	90/99~(90%)	-0.31	1 (1%) 80 74	163, 196, 217, 231	0
9	b	89/99~(89%)	-0.23	0 100 100	173, 219, 252, 260	0
9	g	90/99~(90%)	-0.26	1 (1%) 80 74	170, 212, 266, 282	1 (1%)
9	1	93/99~(93%)	-0.19	2 (2%) 62 53	168, 201, 263, 275	0
10	Х	76/76~(100%)	-0.06	0 100 100	157, 193, 221, 237	0
10	с	76/76~(100%)	-0.04	2 (2%) 56 46	159, 192, 218, 252	0
10	h	76/76~(100%)	0.07	5 (6%) 18 14	170, 195, 214, 239	0
10	m	76/76~(100%)	-0.15	0 100 100	159, 188, 218, 233	0
11	Y	82/104 (78%)	-0.00	1 (1%) 79 72	159, 193, 257, 301	0
11	d	89/104~(85%)	0.12	4 (4%) 33 28	174, 212, 279, 315	0
11	i	89/104 (85%)	0.02	3 (3%) 45 37	162, 200, 266, 297	0
11	n	86/104 (82%)	0.49	6 (6%) 16 12	167, 197, 281, 349	0
All	All	5242/5936 (88%)	-0.17	98 (1%) 66 59	124, 192, 248, 349	2(0%)

Continued from previous page...

All (98) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
11	n	6	ALA	22.8
11	n	5	ASP	10.5
11	d	26	GLY	10.3
11	n	7	GLU	9.9
7	Ζ	397	LYS	6.7
11	d	27	PRO	6.3
7	j	267	ASN	6.2
11	i	11	ILE	6.1
7	Ζ	267	ASN	5.7
8	f	93	VAL	4.9
2	Р	98	TYR	4.7
9	1	3	GLU	4.4
2	Р	102	GLY	4.4
11	n	9	LYS	4.3
2	Р	100	PHE	4.2



Mol	Chain	Res	Type	RSRZ	
8	f	15	TYR	4.0	
11	Y	28	SER	4.0	
2	Р	101	GLY	3.8	
7	Ζ	209	SER	3.8	
8	k	56	THR	3.7	
11	i	10	GLY	3.7	
1	А	134	ARG	3.7	
7	U	3	ILE	3.6	
7	j	392	CYS	3.6	
8	f	89	LEU	3.5	
11	n	8	ILE	3.3	
7	j	349	LEU	3.2	
8	V	93	VAL	3.2	
8	a	60	GLN	3.2	
7	е	392	CYS	3.1	
10	h	61	ILE	3.0	
7	е	340	ASN	3.0	
10	h	13	ILE	3.0	
1	Е	135	ALA	3.0	
3	Q	78	ILE	3.0	
8	V	17	VAL	3.0	
7	Ζ	80	GLN	3.0	
2	F	44	LYS	3.0	
8	k	17	VAL	2.9	
2	L	43	VAL	2.9	
3	Q	117	PRO	2.9	
7	i	304	LEU	2.8	
9	1	9	ASP	2.8	
7	i	268	ASN	2.8	
7	e	38	GLU	2.8	
7	j	309	ASP	2.8	
8	k	14	GLN	2.8	
8	V	90	GLU	2.7	
10	с	43	LEU	2.7	
7	Z	310	ILE	2.7	
7	U	21	LEU	2.7	
7	j	38	GLU	2.7	
8	f	90	GLU	2.7	
10	h	62	GLN	2.6	
7	е	388	MET	2.6	
7	i	259	ASN	2.6	
8	V	14	GLN	2.6	



Mol	Chain	Res	Type	RSRZ	
11	n	10	GLY	2.6	
8	a	55	SER	2.6	
8	a	41	LYS	2.5	
8	V	15	TYR	2.5	
7	Z	450	ILE	2.5	
7	Z	345	ALA	2.5	
11	i	64	LEU	2.4	
7	U	177	ARG	2.4	
7	U	38	GLU	2.4	
2	F	68	ASP	2.4	
7	j	345	ALA	2.4	
8	k	13	GLN	2.4	
11	d	9	LYS	2.4	
8	a	44	ASP	2.4	
1	0	39	HIS	2.4	
7	Z	396	GLU	2.4	
7	е	150	SER	2.4	
7	U	175	LYS	2.3	
8	k	29	LEU	2.3	
10	h	56	LEU	2.3	
10	h	63	LYS	2.3	
11	d	8	ILE	2.3	
7	е	43	THR	2.2	
9	W	9	ASP	2.2	
8	V	18	GLU	2.2	
7	е	84	ILE	2.2	
4	D	31	LYS	2.2	
9	g	80	VAL	2.2	
1	0	40	ARG	2.2	
2	В	43	VAL	2.2	
8	k	12	ILE	2.2	
10	с	69	LEU	2.1	
7	Z	116	LYS	2.1	
8	a	56	THR	2.1	
3	М	23	LEU	2.1	
7	U	0	ALA	2.1	
7	Ζ	43	THR	2.1	
4	Н	86	ILE	2.0	
7	j	111	ASP	2.0	
8	f	92	ILE	2.0	
8	a	18	GLU	2.0	

Continued from previous page...



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
12	ZN	Z	502	1/1	0.58	0.20	233,233,233,233	0
12	ZN	j	501	1/1	0.62	0.11	286,286,286,286	0
12	ZN	е	506	1/1	0.68	0.04	251,251,251,251	0
12	ZN	е	503	1/1	0.70	0.13	229,229,229,229	0
12	ZN	U	501	1/1	0.71	0.16	228,228,228,228	0
12	ZN	U	506	1/1	0.74	0.13	252,252,252,252	0
12	ZN	W	101	1/1	0.76	0.14	199,199,199,199	0
12	ZN	e	502	1/1	0.83	0.10	279,279,279,279	0
12	ZN	Z	505	1/1	0.83	0.17	272,272,272,272	0
12	ZN	е	501	1/1	0.84	0.14	251,251,251,251	0
12	ZN	n	201	1/1	0.84	0.20	216,216,216,216	0
12	ZN	Z	501	1/1	0.85	0.23	234,234,234,234	0
12	ZN	j	504	1/1	0.85	0.16	198,198,198,198	0
12	ZN	d	201	1/1	0.85	0.20	226,226,226,226	0
12	ZN	b	101	1/1	0.86	0.13	218,218,218,218	0
12	ZN	1	101	1/1	0.88	0.19	173,173,173,173	0
12	ZN	j	503	1/1	0.89	0.18	179,179,179,179	0
12	ZN	Y	201	1/1	0.89	0.18	184,184,184,184	0
12	ZN	Z	506	1/1	0.91	0.14	290,290,290,290	0
12	ZN	Z	504	1/1	0.92	0.12	215,215,215,215	0
12	ZN	Z	503	1/1	0.94	0.10	237,237,237,237	0
12	ZN	е	505	1/1	0.94	0.21	152,152,152,152	0
12	ZN	U	503	1/1	0.95	0.18	209,209,209,209	0
12	ZN	j	506	1/1	0.95	0.20	192,192,192,192	0
12	ZN	g	101	1/1	0.97	0.17	206,206,206,206	0
12	ZN	U	504	1/1	0.97	0.12	201,201,201,201	0
12	ZN	j	505	1/1	0.97	0.12	212,212,212,212	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
12	ZN	i	201	1/1	0.98	0.17	$195,\!195,\!195,\!195$	0
12	ZN	U	502	1/1	0.98	0.17	195,195,195,195	0
12	ZN	е	504	1/1	0.99	0.14	188,188,188,188	0
12	ZN	U	505	1/1	0.99	0.18	160,160,160,160	0
12	ZN	j	502	1/1	0.99	0.15	160,160,160,160	0

Continued from previous page...

6.5 Other polymers (i)

There are no such residues in this entry.

