



# Full wwPDB X-ray Structure Validation Report i

Oct 28, 2024 – 07:59 AM JST

PDB ID : 5DUM  
Title : Crystal structure of influenza A virus H5 hemagglutinin globular head in complex with the Fab of antibody 65C6  
Authors : Sun, J.; Zuo, T.; Wang, G.; Zhou, P.; Zhang, L.; Wang, X.  
Deposited on : 2015-09-19  
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2511 (3.00-3.00)
Clashscore	180529	2866 (3.00-3.00)
Ramachandran outliers	177936	2778 (3.00-3.00)
Sidechain outliers	177891	2781 (3.00-3.00)
RSRZ outliers	164620	2523 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4922 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 65C6 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	H	219	Total	C 1694	N 1085	O 281	S 321	7	0	1	0

- Molecule 2 is a protein called 65C6 Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	L	211	Total	C 1621	N 1016	O 275	S 325	5	0	0	0


- Molecule 3 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
3	A	197	Total	C 1579	N 1006	O 272	S 295	6	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	ALA	-	expression tag	UNP Q1WDM0
A	43	ASP	-	expression tag	UNP Q1WDM0
A	44	PRO	-	expression tag	UNP Q1WDM0
A	269	HIS	-	expression tag	UNP Q1WDM0
A	270	HIS	-	expression tag	UNP Q1WDM0
A	271	HIS	-	expression tag	UNP Q1WDM0
A	272	HIS	-	expression tag	UNP Q1WDM0
A	273	HIS	-	expression tag	UNP Q1WDM0
A	274	HIS	-	expression tag	UNP Q1WDM0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total    C    N    O 14    8    1    5	0	0
4	A	1	Total    C    N    O 14    8    1    5	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 65C6 Heavy Chain



- Molecule 2: 65C6 Light Chain



- Molecule 3: Hemagglutinin



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 2 1 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.54Å 92.98Å 134.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.13 – 3.00 36.13 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.1 (36.13-3.00) 99.0 (36.13-3.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	3.04 (at 3.00Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
$R$ , $R_{free}$	0.215 , 0.269 0.217 , 0.267	Depositor DCC
$R_{free}$ test set	798 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	72.7	Xtriage
Anisotropy	0.448	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 17.7	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49$ , $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4922	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	H	0.56	0/1749	0.69	0/2385
2	L	0.53	0/1656	0.65	0/2253
3	A	0.62	1/1621 (0.1%)	0.67	0/2204
All	All	0.57	1/5026 (0.0%)	0.67	0/6842

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	186	GLU	CG-CD	5.04	1.59	1.51

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1694	0	1632	37	0
2	L	1621	0	1582	33	0
3	A	1579	0	1533	38	0
4	A	28	0	26	0	0
All	All	4922	0	4773	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 10.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:126:SER:HB3	1:H:127:SER:HA	1.47	0.95
3:A:82:LYS:HE3	3:A:85:PRO:HA	1.58	0.84
3:A:142:SER:OG	3:A:143:PHE:N	2.18	0.77
2:L:37:GLN:HB2	2:L:47:LEU:HD11	1.66	0.75
1:H:103[A]:HIS:HD2	3:A:163:SER:HB3	1.52	0.75
1:H:140:PRO:HG3	1:H:152:LEU:HB3	1.71	0.73
1:H:73:ASP:HB3	1:H:76:ILE:HD11	1.71	0.72
2:L:61:ARG:NH1	2:L:82:ASP:OD2	2.16	0.72
1:H:60:TYR:HB2	1:H:65:GLN:HG3	1.71	0.71
1:H:139:ALA:HB2	1:H:226:GLU:HG3	1.74	0.70
1:H:100:ARG:HB3	1:H:103[A]:HIS:ND1	2.07	0.70
1:H:52:ASP:OD2	1:H:101:ARG:NH1	2.24	0.69
1:H:126:SER:CB	1:H:127:SER:HA	2.18	0.68
2:L:6:GLN:NE2	2:L:86:TYR:O	2.24	0.67
1:H:103[A]:HIS:CD2	3:A:163:SER:HB3	2.29	0.67
2:L:185:ASP:HA	2:L:188:LYS:HD2	1.77	0.66
2:L:149:LYS:HG2	2:L:154:LEU:HD13	1.76	0.66
3:A:175:LEU:HD13	3:A:230:TRP:HB3	1.81	0.63
3:A:152:LYS:HE2	3:A:189:LYS:O	1.99	0.63
1:H:101:ARG:NH1	3:A:161:LYS:HD2	2.15	0.61
1:H:103[A]:HIS:HD2	3:A:163:SER:CB	2.13	0.60
1:H:101:ARG:CZ	3:A:161:LYS:HD2	2.31	0.59
1:H:214:HIS:CE1	1:H:216:PRO:HG2	2.37	0.59
3:A:53:ARG:HG3	3:A:54:ASP:H	1.68	0.58
2:L:134:CYS:HB2	2:L:148:TRP:CZ2	2.39	0.56
2:L:184:ALA:O	2:L:185:ASP:HB2	2.04	0.56
1:H:87:LYS:O	1:H:90:ASP:N	2.34	0.56
2:L:182:SER:OG	2:L:184:ALA:O	2.24	0.56
3:A:50:LEU:HD22	3:A:108:ILE:CD1	2.36	0.55
3:A:57:VAL:O	3:A:60:TRP:HB3	2.07	0.55
3:A:125:HIS:CE1	3:A:158:PRO:HG2	2.42	0.54
3:A:129:SER:O	3:A:131:VAL:N	2.40	0.54
1:H:18:LEU:HB2	1:H:86:LEU:HD11	1.90	0.54
1:H:57:TYR:OH	3:A:124:ASP:OD2	2.24	0.54
3:A:50:LEU:HD22	3:A:108:ILE:HD12	1.89	0.54
3:A:79:ILE:HG22	3:A:80:VAL:HG23	1.90	0.53
3:A:54:ASP:HB2	3:A:86:ALA:HB3	1.89	0.53
2:L:184:ALA:HA	2:L:187:GLU:OE2	2.08	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:56:SER:HB3	3:A:85:PRO:HG3	1.90	0.52
2:L:67:SER:O	2:L:69:THR:N	2.39	0.52
3:A:182:ASN:ND2	3:A:186:GLU:OE1	2.42	0.52
2:L:202:SER:OG	2:L:203:SER:N	2.42	0.51
2:L:141:PRO:HB2	2:L:143:GLU:OE1	2.11	0.51
1:H:23:LYS:HB3	1:H:78:THR:HG23	1.92	0.50
2:L:73:LEU:HD12	2:L:74:THR:H	1.77	0.50
3:A:53:ARG:CG	3:A:54:ASP:H	2.23	0.50
3:A:160:ILE:O	3:A:242:GLU:HA	2.11	0.50
1:H:13:LYS:HA	1:H:127:SER:OG	2.11	0.50
3:A:53:ARG:HG3	3:A:54:ASP:N	2.27	0.50
1:H:12:LYS:O	1:H:125:VAL:HA	2.12	0.49
1:H:60:TYR:HE2	1:H:70:ILE:HG13	1.76	0.49
1:H:174:THR:O	1:H:177:VAL:HG12	2.13	0.48
2:L:143:GLU:O	2:L:198:HIS:HD2	1.97	0.48
2:L:21:LEU:HD12	2:L:73:LEU:HD23	1.96	0.48
2:L:140:TYR:CD2	2:L:141:PRO:HA	2.49	0.48
3:A:61:LEU:O	3:A:146:ASN:ND2	2.39	0.48
3:A:154:ASN:O	3:A:156:THR:N	2.47	0.48
3:A:178:ILE:HB	3:A:198:ILE:HD12	1.96	0.47
1:H:126:SER:H	1:H:127:SER:HG	1.63	0.46
3:A:151:ILE:HD12	3:A:190:LEU:HD22	1.97	0.46
1:H:91:THR:HG23	1:H:124:THR:HA	1.97	0.46
1:H:87:LYS:HG2	1:H:89:SER:OG	2.16	0.46
1:H:106:GLY:O	3:A:164:TYR:OH	2.23	0.46
1:H:87:LYS:HG3	1:H:88:ALA:N	2.30	0.46
3:A:182:ASN:HA	3:A:214:ALA:O	2.16	0.46
2:L:127:SER:OG	2:L:128:GLY:N	2.47	0.45
3:A:183:ASP:OD1	3:A:185:ALA:HB3	2.16	0.45
1:H:13:LYS:HG2	1:H:127:SER:HB3	1.98	0.45
3:A:182:ASN:HB2	3:A:215:THR:HA	1.98	0.45
2:L:30:SER:O	2:L:30:SER:OG	2.22	0.45
2:L:39:MET:O	2:L:42:GLN:HB2	2.17	0.45
2:L:187:GLU:H	2:L:187:GLU:CD	2.18	0.45
3:A:130:GLY:HA3	3:A:149:TRP:HB3	1.98	0.44
3:A:64:ASN:HB3	3:A:67:CYS:SG	2.57	0.44
1:H:73:ASP:OD1	1:H:75:SER:HB3	2.17	0.44
3:A:64:ASN:ND2	3:A:90:CYS:SG	2.91	0.43
3:A:198:ILE:HD11	3:A:247:PHE:HA	2.00	0.43
1:H:87:LYS:O	1:H:89:SER:N	2.51	0.43
2:L:78:LEU:HD13	2:L:78:LEU:HA	1.88	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:25:PHE:HZ	1:H:79:VAL:HG23	1.84	0.43
2:L:128:GLY:HA2	2:L:183:LYS:HD3	2.00	0.43
2:L:167:ASP:OD2	2:L:168:SER:N	2.51	0.43
1:H:126:SER:N	1:H:127:SER:OG	2.40	0.42
3:A:175:LEU:CD1	3:A:230:TRP:HB3	2.48	0.42
2:L:124:GLN:O	2:L:127:SER:HB3	2.19	0.42
1:H:76:ILE:H	1:H:76:ILE:HG13	1.59	0.42
1:H:18:LEU:CB	1:H:86:LEU:HD11	2.50	0.42
2:L:198:HIS:CG	2:L:199:GLN:N	2.88	0.42
3:A:256:ILE:HD13	3:A:256:ILE:HA	1.83	0.42
1:H:223:LYS:HE2	2:L:123:GLU:OE1	2.20	0.41
2:L:7:SER:HA	2:L:8:PRO:HA	1.86	0.41
2:L:25:ALA:HB3	2:L:69:THR:HG23	2.02	0.41
2:L:187:GLU:O	2:L:211:ARG:NH2	2.52	0.41
2:L:124:GLN:HG2	2:L:129:THR:O	2.20	0.41
2:L:118:PHE:HA	2:L:119:PRO:HD3	1.69	0.41
2:L:128:GLY:C	2:L:183:LYS:HB2	2.42	0.41
1:H:184:LEU:HG	1:H:190:TYR:CE1	2.57	0.40
1:H:38:ARG:NH2	1:H:46:GLU:OE2	2.41	0.40
2:L:185:ASP:N	2:L:187:GLU:OE2	2.55	0.40
3:A:183:ASP:CG	3:A:185:ALA:HB3	2.42	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	H	216/237 (91%)	188 (87%)	21 (10%)	7 (3%)	3 19
2	L	209/214 (98%)	189 (90%)	14 (7%)	6 (3%)	3 20
3	A	193/233 (83%)	166 (86%)	21 (11%)	6 (3%)	3 19
All	All	618/684 (90%)	543 (88%)	56 (9%)	19 (3%)	3 19

All (19) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	226	GLU
2	L	127	SER
3	A	86	ALA
1	H	88	ALA
1	H	201	SER
2	L	68	GLY
3	A	129	SER
3	A	130	GLY
1	H	31	TYR
1	H	227	SER
2	L	128	GLY
2	L	170	ASP
3	A	90	CYS
3	A	134	ALA
3	A	155	ASN
1	H	75	SER
1	H	205	THR
2	L	17	GLU
2	L	40	PRO

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	H	189/204 (93%)	181 (96%)	8 (4%)	25 59
2	L	183/186 (98%)	178 (97%)	5 (3%)	40 71
3	A	176/208 (85%)	171 (97%)	5 (3%)	38 70
All	All	548/598 (92%)	530 (97%)	18 (3%)	33 67

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	5	VAL
1	H	21	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	H	28	SER
1	H	56	SER
1	H	84	SER
1	H	87	LYS
1	H	93	MET
1	H	130	THR
2	L	11	LEU
2	L	45	ARG
2	L	100	GLN
2	L	126	LYS
2	L	191	VAL
3	A	98	TYR
3	A	109	ASN
3	A	135	CYS
3	A	172	LEU
3	A	220	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	115	GLN
3	A	236	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	NAG	A	302	3	14,14,15	1.04	2 (14%)	17,19,21	0.86	1 (5%)
4	NAG	A	301	3	14,14,15	0.45	0	17,19,21	0.84	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	302	3	-	0/6/23/26	0/1/1/1
4	NAG	A	301	3	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	302	NAG	O5-C1	2.78	1.48	1.43
4	A	302	NAG	C1-C2	2.36	1.55	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	302	NAG	C1-O5-C5	2.79	115.97	112.19
4	A	301	NAG	C1-C2-N2	2.01	113.92	110.49

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	301	NAG	C4-C5-C6-O6
4	A	301	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	H	219/237 (92%)	-0.33	0	100	100	31, 61, 83, 88 1 (0%)
2	L	211/214 (98%)	-0.33	0	100	100	52, 68, 79, 88 0
3	A	197/233 (84%)	-0.31	2 (1%)	79	60	50, 66, 92, 96 0
All	All	627/684 (91%)	-0.32	2 (0%)	90	81	31, 65, 86, 96 1 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	49	PRO	2.6
3	A	50	LEU	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	A	301	14/15	0.45	0.13	64,64,64,64	0
4	NAG	A	302	14/15	0.78	0.12	64,64,64,64	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.