



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 02:05 AM EDT

PDB ID : 5HOB  
Title : p73 homo-tetramerization domain mutant I  
Authors : Coutandin, D.; Krojer, T.; Salah, E.; Mathea, S.; Knapp, S.; Dotsch, V.  
Deposited on : 2016-01-19  
Resolution : 1.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1232 (1.24-1.20)
Clashscore	141614	1294 (1.24-1.20)
Ramachandran outliers	138981	1251 (1.24-1.20)
Sidechain outliers	138945	1250 (1.24-1.20)
RSRZ outliers	127900	1209 (1.24-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	50	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 84%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div>
1	B	50	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 86%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div>
1	C	50	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 86%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: grey;"></div> </div>
1	D	50	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 80%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: grey;"></div> </div>
1	E	50	<div style="display: flex; align-items: center;"> <div style="width: 8%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 88%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	50	 <p>8% 86% 12%</p>
1	G	50	 <p>2% 90% 6%</p>
1	H	50	 <p>18% 80% 12% 8%</p>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6145 atoms, of which 2849 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor protein p73.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	45	703	227	343	61	70	2	0	0	0
1	B	45	742	241	372	62	65	2	4	1	0
1	C	46	762	245	382	64	69	2	0	1	0
1	D	44	705	228	351	58	66	2	0	0	1
1	E	48	714	238	343	56	75	2	0	0	0
1	F	44	711	228	354	61	66	2	0	2	0
1	G	47	756	244	377	63	70	2	0	0	0
1	H	46	675	224	327	57	65	2	0	0	1

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	349	GLY	-	expression tag	UNP O15350
A	350	SER	-	expression tag	UNP O15350
A	363	LYS	GLU	engineered mutation	UNP O15350
A	370	GLU	LYS	engineered mutation	UNP O15350
A	373	ARG	GLU	engineered mutation	UNP O15350
A	390	GLU	ARG	engineered mutation	UNP O15350
B	349	GLY	-	expression tag	UNP O15350
B	350	SER	-	expression tag	UNP O15350
B	363	LYS	GLU	engineered mutation	UNP O15350
B	370	GLU	LYS	engineered mutation	UNP O15350
B	373	ARG	GLU	engineered mutation	UNP O15350
B	390	GLU	ARG	engineered mutation	UNP O15350
C	349	GLY	-	expression tag	UNP O15350

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	350	SER	-	expression tag	UNP O15350
C	363	LYS	GLU	engineered mutation	UNP O15350
C	370	GLU	LYS	engineered mutation	UNP O15350
C	373	ARG	GLU	engineered mutation	UNP O15350
C	390	GLU	ARG	engineered mutation	UNP O15350
D	349	GLY	-	expression tag	UNP O15350
D	350	SER	-	expression tag	UNP O15350
D	363	LYS	GLU	engineered mutation	UNP O15350
D	370	GLU	LYS	engineered mutation	UNP O15350
D	373	ARG	GLU	engineered mutation	UNP O15350
D	390	GLU	ARG	engineered mutation	UNP O15350
E	349	GLY	-	expression tag	UNP O15350
E	350	SER	-	expression tag	UNP O15350
E	363	LYS	GLU	engineered mutation	UNP O15350
E	370	GLU	LYS	engineered mutation	UNP O15350
E	373	ARG	GLU	engineered mutation	UNP O15350
E	390	GLU	ARG	engineered mutation	UNP O15350
F	349	GLY	-	expression tag	UNP O15350
F	350	SER	-	expression tag	UNP O15350
F	363	LYS	GLU	engineered mutation	UNP O15350
F	370	GLU	LYS	engineered mutation	UNP O15350
F	373	ARG	GLU	engineered mutation	UNP O15350
F	390	GLU	ARG	engineered mutation	UNP O15350
G	349	GLY	-	expression tag	UNP O15350
G	350	SER	-	expression tag	UNP O15350
G	363	LYS	GLU	engineered mutation	UNP O15350
G	370	GLU	LYS	engineered mutation	UNP O15350
G	373	ARG	GLU	engineered mutation	UNP O15350
G	390	GLU	ARG	engineered mutation	UNP O15350
H	349	GLY	-	expression tag	UNP O15350
H	350	SER	-	expression tag	UNP O15350
H	363	LYS	GLU	engineered mutation	UNP O15350
H	370	GLU	LYS	engineered mutation	UNP O15350
H	373	ARG	GLU	engineered mutation	UNP O15350
H	390	GLU	ARG	engineered mutation	UNP O15350

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	F	1	Total Mg 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	44	Total O 44 44	0	0
3	B	48	Total O 48 48	0	0
3	C	66	Total O 66 66	0	0
3	D	53	Total O 53 53	0	0
3	E	32	Total O 32 32	0	0
3	F	38	Total O 38 38	0	0
3	G	52	Total O 52 52	0	0
3	H	43	Total O 43 43	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tumor protein p73




- Molecule 1: Tumor protein p73




- Molecule 1: Tumor protein p73





- Molecule 1: Tumor protein p73




- Molecule 1: Tumor protein p73





- Molecule 1: Tumor protein p73



- Molecule 1: Tumor protein p73



- Molecule 1: Tumor protein p73





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	29.05Å 48.40Å 75.73Å 89.60° 83.20° 74.27°	Depositor
Resolution (Å)	46.57 – 1.22 46.57 – 1.22	Depositor EDS
% Data completeness (in resolution range)	68.1 (46.57-1.22) 68.1 (46.57-1.22)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.47 (at 1.22Å)	Xtrriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, $R_{free}$	0.188 , 0.214 0.189 , 0.214	Depositor DCC
$R_{free}$ test set	4012 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.1	Xtrriage
Anisotropy	0.377	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 49.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.008 for -h,-h+k,-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6145	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/364	0.62	0/491
1	B	0.46	0/378	0.63	0/510
1	C	0.51	0/388	0.67	1/522 (0.2%)
1	D	0.52	0/359	0.65	0/483
1	E	0.37	0/376	0.54	0/509
1	F	0.39	0/372	0.59	0/502
1	G	0.46	0/384	0.62	0/517
1	H	0.44	0/353	0.57	0/476
All	All	0.45	0/2974	0.61	1/4010 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	C	357	LEU	CA-CB-CG	5.65	128.30	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	360	343	343	1	0
1	B	370	372	372	2	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	380	382	382	2	0
1	D	354	351	351	2	0
1	E	371	343	343	4	0
1	F	357	354	341	1	0
1	G	379	377	377	2	0
1	H	348	327	327	5	0
2	F	1	0	0	0	0
3	A	44	0	0	0	0
3	B	48	0	0	0	1
3	C	66	0	0	0	0
3	D	53	0	0	0	1
3	E	32	0	0	1	0
3	F	38	0	0	0	1
3	G	52	0	0	0	0
3	H	43	0	0	0	0
All	All	3296	2849	2836	12	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:373:ARG:NH2	1:G:390:GLU:OE2	2.25	0.68
1:E:371:LEU:HD11	1:H:371:LEU:HD11	1.78	0.64
1:B:371:LEU:HD11	1:C:371:LEU:HD11	1.90	0.54
1:A:386:VAL:O	1:A:390:GLU:HG2	2.09	0.52
1:G:362:ARG:HD3	1:H:351:ASP:OD1	2.10	0.51
1:F:378:MET:SD	1:H:373:ARG:NH1	2.85	0.50
1:H:375:LEU:O	1:H:378:MET:HG2	2.13	0.48
1:E:364:ASN:HA	1:E:367:ILE:HD12	1.96	0.47
1:E:373:ARG:NH1	3:E:404:HOH:O	2.49	0.45
1:B:389:TYR:HB2	1:D:377:LEU:HD21	1.98	0.45
1:C:356:TYR:CE2	1:D:356:TYR:HB2	2.53	0.43
1:H:357:LEU:HD12	1:H:357:LEU:C	2.42	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:430:HOH:O	3:D:439:HOH:O[1_455]	1.97	0.23
3:B:438:HOH:O	3:F:535:HOH:O[1_546]	2.12	0.08

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	43/50 (86%)	43 (100%)	0	0	100	100
1	B	44/50 (88%)	44 (100%)	0	0	100	100
1	C	45/50 (90%)	45 (100%)	0	0	100	100
1	D	42/50 (84%)	42 (100%)	0	0	100	100
1	E	46/50 (92%)	46 (100%)	0	0	100	100
1	F	44/50 (88%)	44 (100%)	0	0	100	100
1	G	45/50 (90%)	45 (100%)	0	0	100	100
1	H	44/50 (88%)	43 (98%)	1 (2%)	0	100	100
All	All	353/400 (88%)	352 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	38/48 (79%)	37 (97%)	1 (3%)	46	9

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	39/48 (81%)	39 (100%)	0	100	100
1	C	41/48 (85%)	41 (100%)	0	100	100
1	D	38/48 (79%)	36 (95%)	2 (5%)	22	2
1	E	38/48 (79%)	38 (100%)	0	100	100
1	F	39/48 (81%)	39 (100%)	0	100	100
1	G	41/48 (85%)	41 (100%)	0	100	100
1	H	34/48 (71%)	34 (100%)	0	100	100
All	All	308/384 (80%)	305 (99%)	3 (1%)	76	47

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	394	GLN
1	D	353	ASP
1	D	358	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	45/50 (90%)	0.21	2 (4%) 34 33	15, 24, 43, 56	0
1	B	45/50 (90%)	0.18	1 (2%) 62 59	14, 20, 32, 38	1 (2%)
1	C	46/50 (92%)	0.15	1 (2%) 62 59	13, 18, 28, 44	0
1	D	44/50 (88%)	0.14	1 (2%) 60 57	12, 21, 35, 51	0
1	E	48/50 (96%)	0.44	4 (8%) 11 10	16, 29, 45, 51	0
1	F	44/50 (88%)	0.24	4 (9%) 9 9	16, 25, 43, 64	0
1	G	47/50 (94%)	-0.14	1 (2%) 63 60	14, 20, 36, 42	0
1	H	46/50 (92%)	0.70	9 (19%) 1 1	14, 24, 56, 67	0
All	All	365/400 (91%)	0.24	23 (6%) 20 18	12, 22, 46, 67	1 (0%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	396	LEU	5.2
1	F	354	THR	4.8
1	H	384	PRO	4.7
1	E	395	LEU	4.5
1	B	356	TYR	4.0
1	H	382	PRO	4.0
1	H	393	GLN	3.8
1	H	388	SER	3.8
1	D	395	LEU	3.8
1	H	386	VAL	3.7
1	E	361	GLY	3.5
1	F	353	ASP	3.1
1	E	367	ILE	3.0
1	F	355	TYR	2.9
1	H	349	GLY	2.8
1	A	395	LEU	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	H	392	GLN	2.7
1	F	396	LEU	2.4
1	H	390	GLU	2.3
1	H	381	VAL	2.2
1	A	394	GLN	2.2
1	G	397	GLN	2.1
1	C	356	TYR	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	F	401	1/1	0.75	0.10	51,51,51,51	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.