



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 1, 2023 – 01:18 PM JST

PDB ID : 5HXP
Title : Crystal Structure of Z,Z-Farnesyl Diphosphate Synthase (D71M, E75A and H103Y Mutants) Complexed with IPP
Authors : Lee, C.C.; Chan, Y.T.; Wang, A.H.J.
Deposited on : 2016-01-31
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	
1	C	233	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3870 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called (2Z,6Z)-farnesyl diphosphate synthase, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	224	Total 1798	C 1151	N 300	O 337	S 10	0	0	0
1	C	223	Total 1789	C 1148	N 295	O 336	S 10	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	71	MET	-	initiating methionine	UNP B8XA40
A	75	ALA	GLU	engineered mutation	UNP B8XA40
A	103	TYR	HIS	engineered mutation	UNP B8XA40
C	71	MET	-	initiating methionine	UNP B8XA40
C	75	ALA	GLU	engineered mutation	UNP B8XA40
C	103	TYR	HIS	engineered mutation	UNP B8XA40

- Molecule 2 is ISOPENTYL PYROPHOSPHATE (three-letter code: IPR) (formula: $C_5H_{14}O_7P_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			14	5	7	2		
2	C	1	Total	C	O	P	0	0
			14	5	7	2		
2	C	1	Total	C	O	P	0	0
			14	5	7	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is 1,4,7,10,13,16-HEXAOXACYCLOOCTADECANE (three-letter code: O4B) (formula: C₁₂H₂₄O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			18	12	6		
4	C	1	Total	C	O	0	0
			18	12	6		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	95	Total	O	0	0
			95	95		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	109	Total 109	O 109	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: (2Z,6Z)-farnesyl diphosphate synthase, chloroplastic



- Molecule 1: (2Z,6Z)-farnesyl diphosphate synthase, chloroplastic



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.31Å 64.55Å 127.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 1.95 24.52 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.4 (25.00-1.95) 98.5 (24.52-1.95)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.29 (at 1.95Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.199 , 0.243 0.208 , 0.248	Depositor DCC
R_{free} test set	1761 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtrriage
Anisotropy	0.001	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 62.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3870	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: O4B, MG, IPR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/1832	0.68	0/2466
1	C	0.46	0/1823	0.62	0/2456
All	All	0.45	0/3655	0.65	0/4922

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	141	GLU	Peptide
1	C	293	GLN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1798	0	1787	20	0
1	C	1789	0	1790	11	0
2	A	14	0	10	0	0
2	C	28	0	20	0	0
3	C	1	0	0	0	0
4	C	36	0	48	0	0
5	A	95	0	0	6	0
5	C	109	0	0	3	0
All	All	3870	0	3655	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:GLU:N	1:A:142:VAL:HB	1.67	1.08
1:A:141:GLU:HB3	1:A:142:VAL:HG23	1.63	0.78
1:C:182:LEU:O	1:C:186:THR:HG23	1.84	0.78
1:A:260:LEU:C	5:A:504:HOH:O	2.27	0.72
1:A:131:SER:OG	1:A:134:ASN:ND2	2.31	0.63
1:A:171:ASP:OD1	1:A:171:ASP:N	2.31	0.62
1:A:141:GLU:N	1:A:142:VAL:CB	2.55	0.61
1:A:301:HIS:C	5:A:563:HOH:O	2.42	0.58
1:A:89:ARG:NH1	5:A:502:HOH:O	2.37	0.56
1:A:152:LEU:HD21	1:A:197:ILE:CD1	2.37	0.54
1:A:103:TYR:CD1	1:A:145:LEU:HD21	2.43	0.53
1:C:76:LEU:HA	5:C:564:HOH:O	2.09	0.53
1:A:264:LEU:HD12	5:A:504:HOH:O	2.09	0.52
1:C:174:MET:HG3	5:C:599:HOH:O	2.10	0.52
1:A:215:VAL:O	1:A:219:MET:HG2	2.10	0.51
1:C:152:LEU:HD22	1:C:183:THR:HG21	1.93	0.51
1:A:260:LEU:N	5:A:504:HOH:O	2.44	0.50
1:A:261:LEU:N	5:A:504:HOH:O	2.43	0.49
1:C:103:TYR:OH	5:C:502:HOH:O	2.16	0.48
1:A:268:GLU:OE2	1:C:255:ARG:NH1	2.45	0.48
1:C:81:ILE:HD12	1:C:122:ILE:HD13	1.95	0.48
1:C:152:LEU:HD11	1:C:165:ILE:HD11	1.97	0.46
1:A:182:LEU:O	1:A:186:THR:HG23	2.15	0.46
1:A:164:SER:C	1:A:165:ILE:HD12	2.39	0.43
1:C:292:GLN:O	1:C:294:ARG:HB2	2.18	0.43
1:A:260:LEU:HB3	1:A:263:GLN:HB2	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:ILE:HD12	1:A:122:ILE:HD13	2.00	0.42
1:A:168:CYS:HB2	1:A:200:ASN:ND2	2.35	0.42
1:C:153:TYR:CE1	1:C:157:SER:OG	2.72	0.41
1:C:245:ASP:HB3	1:C:291:PHE:CE1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	220/233 (94%)	216 (98%)	3 (1%)	1 (0%)	29	17
1	C	221/233 (95%)	218 (99%)	3 (1%)	0	100	100
All	All	441/466 (95%)	434 (98%)	6 (1%)	1 (0%)	47	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	142	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/204 (96%)	190 (97%)	6 (3%)	40	28

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	195/204 (96%)	189 (97%)	6 (3%)	40	28
All	All	391/408 (96%)	379 (97%)	12 (3%)	40	28

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	128	PHE
1	A	151	GLU
1	A	171	ASP
1	A	175	THR
1	A	182	LEU
1	A	233	PHE
1	C	79	LYS
1	C	104	LYS
1	C	128	PHE
1	C	152	LEU
1	C	158	ARG
1	C	294	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	134	ASN
1	A	190	ASN
1	C	190	ASN
1	C	243	ASN
1	C	293	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	IPR	A	401	-	11,13,13	0.97	0	16,19,19	1.67	2 (12%)
2	IPR	C	405	3	11,13,13	0.88	0	16,19,19	1.42	4 (25%)
2	IPR	C	404	3	11,13,13	0.91	0	16,19,19	1.52	2 (12%)
4	O4B	C	403	-	18,18,18	0.74	0	18,18,18	0.89	0
4	O4B	C	402	-	18,18,18	0.84	0	18,18,18	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IPR	A	401	-	-	2/13/13/13	-
2	IPR	C	405	3	-	2/13/13/13	-
2	IPR	C	404	3	-	7/13/13/13	-
4	O4B	C	403	-	-	7/18/18/18	0/1/1/1
4	O4B	C	402	-	-	5/18/18/18	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	IPR	P7-O10-P11	-5.26	114.78	132.83
2	C	404	IPR	P7-O10-P11	-4.25	118.26	132.83
2	C	405	IPR	O12-P11-O10	2.66	113.54	104.64
2	C	405	IPR	P7-O10-P11	-2.66	123.71	132.83
2	C	404	IPR	O12-P11-O10	2.31	112.39	104.64
2	C	405	IPR	O14-P11-O10	2.25	112.19	104.64
2	A	401	IPR	C3-C2-C1	2.18	120.56	110.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	405	IPR	C3-C2-C4	2.08	124.36	111.54

There are no chirality outliers.

All (23) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	404	IPR	C5-O6-P7-O10
2	C	404	IPR	C5-O6-P7-O8
2	C	404	IPR	C5-O6-P7-O9
2	C	405	IPR	C3-C2-C4-C5
4	C	403	O4B	OAM-CAA-CAB-OAN
2	C	404	IPR	C4-C5-O6-P7
4	C	402	O4B	OAN-CAE-CAF-OAP
4	C	402	O4B	CAK-CAL-OAR-CAJ
2	A	401	IPR	C3-C2-C4-C5
4	C	402	O4B	CAB-CAA-OAM-CAC
4	C	402	O4B	CAC-CAD-OAO-CAG
4	C	403	O4B	CAH-CAG-OAO-CAD
2	A	401	IPR	C2-C4-C5-O6
4	C	403	O4B	CAF-CAE-OAN-CAB
4	C	402	O4B	CAJ-CAI-OAP-CAF
4	C	403	O4B	CAB-CAA-OAM-CAC
4	C	403	O4B	OAP-CAI-CAJ-OAR
2	C	404	IPR	C1-C2-C4-C5
4	C	403	O4B	CAI-CAJ-OAR-CAL
4	C	403	O4B	OAQ-CAK-CAL-OAR
2	C	404	IPR	C3-C2-C4-C5
2	C	404	IPR	C2-C4-C5-O6
2	C	405	IPR	C2-C4-C5-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/233 (96%)	0.68	25 (11%) 5 8	9, 22, 42, 54	0
1	C	223/233 (95%)	0.68	21 (9%) 8 13	7, 22, 40, 53	0
All	All	447/466 (95%)	0.68	46 (10%) 6 10	7, 22, 41, 54	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	96	GLY	4.7
1	A	144	PHE	4.6
1	C	76	LEU	4.1
1	A	143	ASP	4.1
1	C	83	LEU	3.8
1	C	222	LEU	3.7
1	A	192	GLY	3.6
1	A	295	HIS	3.4
1	A	158	ARG	3.3
1	A	142	VAL	3.2
1	C	84	ILE	3.1
1	C	106	LEU	3.1
1	C	72	LEU	3.0
1	A	110	LEU	3.0
1	A	191	LYS	3.0
1	C	224	ASP	3.0
1	A	301	HIS	3.0
1	A	83	LEU	2.8
1	C	110	LEU	2.7
1	A	147	GLN	2.6
1	C	95	LYS	2.6
1	C	73	ASP	2.6
1	C	158	ARG	2.6
1	A	159	SER	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	115	ASP	2.5
1	C	91	TRP	2.5
1	C	75	ALA	2.5
1	A	145	LEU	2.5
1	C	93	LYS	2.5
1	A	222	LEU	2.5
1	A	152	LEU	2.4
1	C	129	ALA	2.3
1	A	149	PHE	2.3
1	A	84	ILE	2.2
1	C	154	ASP	2.2
1	A	219	MET	2.2
1	A	134	ASN	2.2
1	C	155	GLU	2.1
1	A	171	ASP	2.1
1	A	207	ILE	2.1
1	A	82	ALA	2.1
1	A	141	GLU	2.1
1	A	284	LEU	2.1
1	A	150	GLU	2.0
1	C	282	GLU	2.0
1	C	276	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	O4B	C	403	18/18	0.84	0.18	41,43,47,48	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	IPR	A	401	14/14	0.86	0.25	48,51,55,56	0
4	O4B	C	402	18/18	0.90	0.16	26,28,33,34	0
2	IPR	C	404	14/14	0.94	0.15	23,29,33,33	0
2	IPR	C	405	14/14	0.96	0.12	25,26,28,28	0
3	MG	C	401	1/1	0.99	0.08	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.