



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 10:35 AM EDT

PDB ID : 5KKD  
Title : Truncated hemolysin A Y134A from *P. mirabilis* at 2.1 Angstroms resolution crystallized in a high salt condition  
Authors : Novak, W.R.P.; Bhattacharyya, B.; Weaver, T.M.  
Deposited on : 2016-06-21  
Resolution : 2.13 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	242	
1	B	242	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3135 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemolysin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	204	1494	923	266	302	3	0	1	0
1	B	205	1474	915	260	296	3	0	2	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	134	ALA	TYR	engineered mutation	UNP P16466
A	266	HIS	-	expression tag	UNP P16466
A	267	HIS	-	expression tag	UNP P16466
A	268	HIS	-	expression tag	UNP P16466
A	269	HIS	-	expression tag	UNP P16466
A	270	HIS	-	expression tag	UNP P16466
A	271	HIS	-	expression tag	UNP P16466
B	134	ALA	TYR	engineered mutation	UNP P16466
B	266	HIS	-	expression tag	UNP P16466
B	267	HIS	-	expression tag	UNP P16466
B	268	HIS	-	expression tag	UNP P16466
B	269	HIS	-	expression tag	UNP P16466
B	270	HIS	-	expression tag	UNP P16466
B	271	HIS	-	expression tag	UNP P16466

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	O S	0	0
			5	4 1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	87	Total	O	0	0
			87	87		
3	B	75	Total	O	0	0
			75	75		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.08Å 69.08Å 234.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.82 – 2.13 47.82 – 2.13	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.82-2.13) 95.9 (47.82-2.13)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.09 (at 2.12Å)	Xtrriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, $R_{free}$	0.170 , 0.195 0.172 , 0.198	Depositor DCC
$R_{free}$ test set	1650 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.9	Xtrriage
Anisotropy	0.573	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3135	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/1517	0.58	0/2066
1	B	0.41	0/1496	0.56	0/2039
All	All	0.41	0/3013	0.57	0/4105

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1494	0	1443	9	0
1	B	1474	0	1417	22	0
2	B	5	0	0	1	0
3	A	87	0	0	0	0
3	B	75	0	0	0	0
All	All	3135	0	2860	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:ASN:HD22	1:A:120:PHE:H	1.18	0.89
1:A:176:ASN:H	1:A:231:GLN:HE22	1.22	0.86
1:B:72:ASN:HD22	1:B:120:PHE:H	1.27	0.83
1:B:176:ASN:H	1:B:231:GLN:HE22	1.26	0.82
1:B:56:VAL:H	1:B:66:ASN:HD21	1.39	0.68
1:B:176:ASN:H	1:B:231:GLN:NE2	1.97	0.59
1:A:72:ASN:ND2	1:A:120:PHE:H	1.97	0.58
1:B:223:GLN:CD	1:B:223:GLN:H	2.11	0.54
1:B:223:GLN:NE2	2:B:301:SO4:O1	2.33	0.54
1:B:33:VAL:HG22	1:B:70:ASP:HB2	1.90	0.53
1:A:72:ASN:HD22	1:A:120:PHE:N	1.97	0.53
1:B:41:PRO:HB3	1:B:66:ASN:HD22	1.74	0.51
1:A:176:ASN:H	1:A:231:GLN:NE2	2.01	0.49
1:B:176:ASN:N	1:B:231:GLN:HE22	2.02	0.48
1:A:211:GLU:OE2	1:B:206:LYS:HD2	2.13	0.48
1:B:81:ASN:HD22	1:B:81:ASN:C	2.15	0.47
1:A:144[A]:CYS:SG	1:A:157:LEU:HD13	2.54	0.47
1:B:71:PHE:HZ	1:B:80:PHE:HZ	1.64	0.46
1:A:63:ILE:HG12	1:A:108:LEU:HD23	1.98	0.45
1:B:127:VAL:HG23	1:B:152:THR:HG22	1.98	0.45
1:B:41:PRO:HB3	1:B:66:ASN:ND2	2.33	0.44
1:B:56:VAL:H	1:B:66:ASN:ND2	2.12	0.44
1:A:32:ILE:HD13	1:A:53:ILE:HD11	1.99	0.43
1:B:108:LEU:HD11	1:B:135:VAL:HG23	2.00	0.43
1:B:171:GLY:HA3	1:B:221:PHE:O	2.19	0.42
1:B:170:LYS:HD2	1:B:170:LYS:HA	1.79	0.42
1:B:81:ASN:HD21	1:B:83:ALA:HB3	1.85	0.42
1:B:33:VAL:HG23	1:B:69:GLN:HB2	2.02	0.42
1:B:198:ALA:O	1:B:216:THR:HA	2.19	0.41
1:B:81:ASN:HD22	1:B:83:ALA:H	1.70	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	203/242 (84%)	199 (98%)	4 (2%)	0	100	100
1	B	205/242 (85%)	200 (98%)	5 (2%)	0	100	100
All	All	408/484 (84%)	399 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/194 (84%)	161 (99%)	2 (1%)	71	74
1	B	157/194 (81%)	154 (98%)	3 (2%)	57	59
All	All	320/388 (82%)	315 (98%)	5 (2%)	62	65

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	147	CYS
1	A	155	SER
1	B	81	ASN
1	B	147	CYS
1	B	223	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	ASN
1	A	90	GLN
1	A	111	ASN
1	A	140	ASN
1	A	179	ASN
1	A	231	GLN
1	B	66	ASN
1	B	72	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	81	ASN
1	B	111	ASN
1	B	231	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	B	301	-	4,4,4	0.17	0	6,6,6	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	204/242 (84%)	0.04	4 (1%) 65 71	25, 39, 66, 101	0
1	B	205/242 (84%)	0.14	12 (5%) 22 28	26, 41, 74, 116	0
All	All	409/484 (84%)	0.09	16 (3%) 39 47	25, 40, 70, 116	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	47	ASN	5.1
1	B	102	ASN	4.1
1	B	234	VAL	3.8
1	B	36	ALA	3.7
1	B	48	GLY	3.5
1	A	38	HIS	3.4
1	A	166	ASN	3.3
1	B	103	GLY	2.8
1	B	30	ASN	2.8
1	A	102	ASN	2.3
1	B	99	SER	2.3
1	A	152	THR	2.2
1	B	166	ASN	2.2
1	B	37	GLY	2.1
1	B	96	ASN	2.1
1	B	193	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	B	301	5/5	0.93	0.16	52,68,74,78	5

### 6.5 Other polymers [i](#)

There are no such residues in this entry.