



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 08:57 PM EDT

PDB ID : 5L3B  
Title : Human LSD1/CoREST: LSD1 D556G mutation  
Authors : Pilotto, S.; Speranzini, V.; Marabelli, C.  
Deposited on : 2016-04-06  
Resolution : 3.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	852	
2	B	482	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6342 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine-specific histone demethylase 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	666	5213	3322	906	965	20	0	0	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	556	GLY	ASP	engineered mutation	UNP O60341

- Molecule 2 is a protein called REST corepressor 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	133	1076	676	194	203	3	0	0	0

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	53	27	9	15	2	0	0





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	119.09Å 179.12Å 234.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.30 45.79 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (50.00-3.30) 99.5 (45.79-3.30)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.96 (at 3.32Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.195 , 0.208 0.202 , 0.216	Depositor DCC
$R_{free}$ test set	740 reflections (1.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	81.5	Xtrriage
Anisotropy	0.034	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 53.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6342	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	85.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/5327	0.66	2/7226 (0.0%)
2	B	0.45	0/1091	0.69	0/1471
All	All	0.44	0/6418	0.67	2/8697 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	820	ARG	NE-CZ-NH1	5.44	123.02	120.30
1	A	815	LEU	CA-CB-CG	5.41	127.74	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5213	0	5251	39	0
2	B	1076	0	1091	9	0
3	A	53	0	31	3	0
All	All	6342	0	6373	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:ILE:HD11	1:A:248:LEU:HD11	1.71	0.71
1:A:670:PHE:CZ	1:A:740:VAL:HG12	2.38	0.59
1:A:801:GLU:HG3	1:A:809:ALA:HA	1.83	0.59
1:A:566:THR:HG21	1:A:697:LEU:HD13	1.86	0.57
1:A:631:LYS:HE2	1:A:651:VAL:O	2.06	0.56
1:A:353:LEU:HB3	1:A:565:LEU:HD23	1.89	0.55
1:A:392:LEU:HD23	1:A:398:PHE:CD2	2.41	0.55
1:A:761:TYR:CD1	1:A:809:ALA:HB1	2.40	0.55
1:A:801:GLU:O	1:A:802:HIS:ND1	2.40	0.54
1:A:670:PHE:CE1	1:A:740:VAL:HG12	2.44	0.52
1:A:331:ALA:HA	3:A:901:FAD:N5	2.25	0.52
1:A:564:HIS:C	1:A:565:LEU:HD12	2.32	0.50
1:A:442:LYS:HE3	2:B:355:THR:HG21	1.94	0.49
1:A:538:PHE:CE1	1:A:706:LEU:HD23	2.48	0.49
1:A:694:PHE:HA	1:A:704:LEU:O	2.12	0.48
1:A:435:VAL:HG13	2:B:349:ILE:HD12	1.95	0.48
1:A:760:SER:HB2	3:A:901:FAD:HM83	1.96	0.47
1:A:366:ASN:OD1	1:A:367:GLY:N	2.47	0.47
1:A:756:TRP:O	1:A:757:ALA:HB2	2.15	0.46
1:A:340:ASN:OD1	1:A:342:MET:N	2.43	0.45
1:A:366:ASN:OD1	1:A:368:GLN:N	2.49	0.45
2:B:368:GLU:OE2	2:B:371:ARG:NH1	2.48	0.45
2:B:327:ASN:ND2	2:B:330:ALA:HB2	2.32	0.45
1:A:801:GLU:O	1:A:802:HIS:CG	2.71	0.44
2:B:395:ILE:HG22	2:B:433:VAL:HG12	1.99	0.43
1:A:659:LEU:HD11	3:A:901:FAD:HM73	2.00	0.43
1:A:418:LEU:CD1	2:B:324:VAL:HG21	2.48	0.43
1:A:214:ARG:O	1:A:218:LEU:HD12	2.19	0.42
1:A:671:TRP:O	1:A:673:PRO:HD3	2.20	0.42
1:A:700:ALA:HB1	1:A:701:PRO:HD2	2.02	0.42
1:A:384:ARG:NH1	2:B:312:LYS:O	2.48	0.41
1:A:804:ILE:O	1:A:804:ILE:HG23	2.20	0.41
1:A:534:ALA:HB2	1:A:688:ARG:HG3	2.02	0.41
1:A:632:GLN:CD	1:A:758:ARG:HE	2.24	0.41
1:A:374:LYS:NZ	1:A:525:ASP:OD1	2.54	0.41
1:A:666:PHE:O	1:A:701:PRO:HG2	2.21	0.41
1:A:668:ARG:HD3	1:A:741:PRO:HG3	2.03	0.41
1:A:695:TRP:CE3	1:A:697:LEU:HD11	2.56	0.41
1:A:473:ASP:OD1	1:A:475:THR:N	2.53	0.40
1:A:418:LEU:HD12	2:B:324:VAL:HG21	2.03	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:456:LYS:HA	2:B:370:TYR:CD1	2.57	0.40
1:A:594:ARG:HD2	1:A:603:ILE:HD11	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	664/852 (78%)	634 (96%)	28 (4%)	2 (0%)	41	71
2	B	131/482 (27%)	119 (91%)	10 (8%)	2 (2%)	10	38
All	All	795/1334 (60%)	753 (95%)	38 (5%)	4 (0%)	29	61

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	757	ALA
2	B	326	ALA
2	B	439	ALA
1	A	468	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	565/698 (81%)	533 (94%)	32 (6%)	20	51
2	B	117/395 (30%)	105 (90%)	12 (10%)	7	26
All	All	682/1093 (62%)	638 (94%)	44 (6%)	17	46

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	191	GLN
1	A	243	ASN
1	A	246	THR
1	A	269	ARG
1	A	273	LEU
1	A	328	ASP
1	A	368	GLN
1	A	437	THR
1	A	448	MET
1	A	458	LEU
1	A	469	LYS
1	A	473	ASP
1	A	479	LEU
1	A	485	ARG
1	A	492	LYS
1	A	508	LEU
1	A	511	LEU
1	A	523	SER
1	A	525	ASP
1	A	526	ARG
1	A	571	TYR
1	A	591	ARG
1	A	594	ARG
1	A	610	THR
1	A	611	SER
1	A	668	ARG
1	A	677	LEU
1	A	684	THR
1	A	696	ASN
1	A	719	SER
1	A	750	ARG
1	A	776	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	314	MET
2	B	316	LEU
2	B	317	SER
2	B	332	THR
2	B	337	GLN
2	B	344	SER
2	B	349	ILE
2	B	371	ARG
2	B	384	THR
2	B	385	THR
2	B	425	ARG
2	B	426	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	243	ASN
1	A	828	GLN
2	B	327	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FAD	A	901	-	53,58,58	1.36	7 (13%)	68,89,89	1.50	13 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	901	-	-	2/30/50/50	0/6/6/6

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	901	FAD	C9A-C5X	5.52	1.50	1.41
3	A	901	FAD	C8-C7	3.29	1.49	1.40
3	A	901	FAD	C5X-N5	-2.83	1.34	1.39
3	A	901	FAD	C4-N3	-2.37	1.34	1.38
3	A	901	FAD	C5A-C4A	2.35	1.47	1.40
3	A	901	FAD	C2A-N3A	2.25	1.35	1.32
3	A	901	FAD	C10-N10	2.24	1.42	1.37

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	FAD	O4-C4-C4X	-3.73	116.70	126.60
3	A	901	FAD	C4X-C10-N1	-3.21	117.28	124.73
3	A	901	FAD	N3A-C2A-N1A	-3.13	123.79	128.68
3	A	901	FAD	C10-N1-C2	2.82	122.55	116.90
3	A	901	FAD	C4X-C4-N3	2.57	119.71	113.19
3	A	901	FAD	C6-C5X-C9A	2.54	122.53	118.94
3	A	901	FAD	C5'-C4'-C3'	-2.54	107.30	112.20
3	A	901	FAD	O4'-C4'-C3'	2.47	115.12	109.10
3	A	901	FAD	O4B-C1B-C2B	-2.39	103.44	106.93
3	A	901	FAD	N6A-C6A-N1A	2.32	123.39	118.57
3	A	901	FAD	P-O3P-PA	-2.24	125.13	132.83
3	A	901	FAD	N10-C10-N1	2.19	124.65	118.35
3	A	901	FAD	C1'-N10-C9A	2.11	124.04	120.51

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	901	FAD	PA-O3P-P-O5'
3	A	901	FAD	O4B-C4B-C5B-O5B

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	901	FAD	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	666/852 (78%)	0.05	4 (0%) 89 90	47, 79, 113, 143	0
2	B	133/482 (27%)	0.35	3 (2%) 60 59	71, 107, 130, 148	0
All	All	799/1334 (59%)	0.10	7 (0%) 84 84	47, 83, 120, 148	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	242	TYR	2.5
1	A	171	PRO	2.5
2	B	374	GLU	2.4
1	A	509	GLN	2.2
1	A	270	ILE	2.2
2	B	312	LYS	2.2
2	B	365	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	FAD	A	901	53/53	0.97	0.22	49,58,68,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.