



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 28, 2024 – 08:24 pm GMT

PDB ID : 5LHR
Title : The catalytic domain of murine urokinase-type plasminogen activator in complex with the active site binding inhibitory nanobody Nb22
Authors : Kromann-Hansen, T.; Lange, E.L.; Sorensen, H.P.; Ghassabeh, G.H.; Huang, M.; Jensen, J.K.; Muyldermans, S.; Declerck, P.J.; Andreasen, P.A.
Deposited on : 2016-07-12
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	 2% 80% 19% ..
2	B	140	 % 65% 18% 17%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2975 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Urokinase-type plasminogen activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	245	1926	1225	322	365	14	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	122	ALA	CYS	engineered mutation	UNP P06869

- Molecule 2 is a protein called Camelid-Derived Antibody Fragment Nb22.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	116	876	544	153	175	4	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	132	Total	O	0	0
			132	132		
3	B	41	Total	O	0	0
			41	41		

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.09Å 66.51Å 57.03Å 90.00° 91.38° 90.00°	Depositor
Resolution (Å)	32.47 – 2.30 32.47 – 2.30	Depositor EDS
% Data completeness (in resolution range)	89.4 (32.47-2.30) 89.4 (32.47-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.29Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.188 , 0.242 0.189 , 0.242	Depositor DCC
R_{free} test set	720 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	17.3	Xtrriage
Anisotropy	0.218	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.046 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2975	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.46	3/1979 (0.2%)	0.60	4/2680 (0.1%)
2	B	0.49	2/893 (0.2%)	0.49	0/1208
All	All	0.47	5/2872 (0.2%)	0.57	4/3888 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	57	SER	CB-OG	-6.70	1.33	1.42
1	A	218	GLY	C-N	-6.55	1.19	1.34
1	A	217	ARG	C-N	-5.99	1.22	1.33
2	B	55	GLY	C-O	-5.84	1.14	1.23
1	A	193	GLY	C-O	-5.75	1.14	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	GLY	O-C-N	-10.23	106.33	122.70
1	A	218	GLY	CA-C-N	7.00	132.60	117.20
1	A	172	TYR	N-CA-C	6.43	128.36	111.00
1	A	218	GLY	C-N-CA	6.15	137.09	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1926	0	1848	38	0
2	B	876	0	830	16	0
3	A	132	0	0	2	0
3	B	41	0	0	2	0
All	All	2975	0	2678	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:ILE:O	1:A:242:ILE:HD13	1.36	1.25
1:A:238:ILE:O	1:A:242:ILE:CD1	2.05	1.05
1:A:27:GLN:HG3	1:A:139:THR:HG21	1.53	0.88
1:A:238:ILE:HG22	1:A:242:ILE:HD11	1.59	0.83
1:A:177:ASN:ND2	3:A:301:HOH:O	2.14	0.81
1:A:51:TRP:CD2	1:A:242:ILE:HG22	2.18	0.78
1:A:48:SER:HB3	1:A:51:TRP:HB2	1.69	0.74
1:A:238:ILE:HG22	1:A:242:ILE:CD1	2.19	0.72
1:A:77:ASN:HB2	1:A:80:GLU:HG3	1.73	0.70
1:A:51:TRP:CE2	1:A:242:ILE:HG22	2.28	0.68
2:B:32:TYR:O	2:B:72:ARG:NH2	2.25	0.68
1:A:116:ARG:NH2	3:A:302:HOH:O	2.28	0.63
2:B:19:ARG:NH2	3:B:201:HOH:O	2.33	0.59
1:A:135:ASP:OD1	1:A:161:LYS:NZ	2.37	0.58
2:B:83:MET:HB3	2:B:86:LEU:HD21	1.87	0.56
1:A:238:ILE:C	1:A:242:ILE:CD1	2.74	0.55
1:A:238:ILE:CG2	1:A:242:ILE:CD1	2.85	0.53
1:A:180:MET:HE3	1:A:227:VAL:HG11	1.91	0.53
1:A:238:ILE:CG2	1:A:242:ILE:HD12	2.40	0.52
1:A:242:ILE:HG12	1:A:243:GLY:N	2.24	0.52
1:A:124:PRO:HD3	1:A:209:LEU:O	2.11	0.50
2:B:39:GLN:NE2	3:B:206:HOH:O	2.42	0.50
1:A:42:CYS:HB2	1:A:195:SER:O	2.12	0.49
1:A:238:ILE:HG23	1:A:242:ILE:HD12	1.95	0.49
2:B:40:ALA:HB3	2:B:43:LYS:HD2	1.94	0.48
2:B:38:ARG:NE	2:B:46:GLU:OE2	2.35	0.48
1:A:180:MET:HE3	1:A:215:TRP:HE1	1.79	0.48
2:B:53:TRP:HA	2:B:72:ARG:CZ	2.43	0.48
1:A:213:VAL:HG22	1:A:228:TYR:HE2	1.78	0.47
1:A:216:GLY:O	2:B:103:SER:OG	2.25	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:71:SER:O	1:A:154:ASN:ND2	2.48	0.47
2:B:34:MET:HG2	2:B:72:ARG:NH2	2.30	0.47
1:A:165:HIS:CD2	1:A:178:TYR:HA	2.50	0.46
1:A:45:SER:OG	1:A:198:PRO:HB3	2.16	0.46
1:A:57:HIS:ND1	1:A:102:ASP:OD2	2.45	0.46
1:A:181:LEU:HD23	1:A:230:ARG:HG3	1.98	0.46
1:A:242:ILE:HD13	1:A:242:ILE:H	1.81	0.45
1:A:51:TRP:CG	1:A:242:ILE:HG22	2.53	0.44
2:B:5:GLN:NE2	2:B:6:GLU:O	2.43	0.44
2:B:72:ARG:HD3	2:B:74:ASN:OD1	2.17	0.44
2:B:87:LYS:HE2	2:B:87:LYS:HB2	1.90	0.43
1:A:125:PRO:HG2	1:A:128:THR:HB	2.00	0.43
1:A:97(B):LEU:HD23	1:A:217:ARG:HG3	2.00	0.43
1:A:49:PRO:O	1:A:111:CYS:HB2	2.19	0.43
2:B:22:CYS:HB3	2:B:79:VAL:HG13	2.01	0.42
1:A:97(B):LEU:HD12	1:A:97(B):LEU:HA	1.86	0.42
1:A:27:GLN:CG	1:A:139:THR:HG21	2.35	0.42
1:A:165:HIS:NE2	1:A:180:MET:O	2.53	0.42
2:B:87:LYS:O	2:B:119:VAL:HG11	2.20	0.41
2:B:2:VAL:HB	2:B:110:TYR:CE2	2.55	0.41
1:A:71:SER:C	1:A:72:LYS:HD2	2.42	0.40
1:A:126:ARG:HA	1:A:235:LEU:HD12	2.03	0.40
2:B:2:VAL:HG23	2:B:3:GLN:H	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/247 (98%)	232 (96%)	9 (4%)	0	100	100
2	B	112/140 (80%)	106 (95%)	6 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	353/387 (91%)	338 (96%)	15 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/212 (100%)	208 (98%)	4 (2%)	52	69
2	B	92/111 (83%)	92 (100%)	0	100	100
All	All	304/323 (94%)	300 (99%)	4 (1%)	65	79

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	ARG
1	A	167	GLN
1	A	191	CYS
1	A	242	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	154	ASN
1	A	167	GLN
1	A	233	HIS
2	B	39	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	218:GLY	C	220:CYS	N	1.19

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/247 (99%)	-0.05	4 (1%) 70 71	7, 15, 30, 47	0
2	B	116/140 (82%)	0.20	2 (1%) 69 69	11, 22, 45, 57	0
All	All	361/387 (93%)	0.03	6 (1%) 69 69	7, 17, 36, 57	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	154	ASN	3.7
1	A	243	GLY	3.7
2	B	110	TYR	2.9
2	B	121	SER	2.1
1	A	242	ILE	2.1
1	A	167	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.