



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 04:09 PM EDT

PDB ID : 5LQB
Title : Complex structure of human IL2 mutant, Proleukin, with Fab fragment of NARA1 antibody
Authors : Zou, C.; Wirth, E.
Deposited on : 2016-08-16
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	132	
2	H	224	
3	L	218	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4828 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	128	1035	667	169	193	6	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	145	SER	CYS	engineered mutation	UNP P60568

- Molecule 2 is a protein called anti-hIL2 FAB fragment heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	219	1627	1028	265	328	6	0	0	0

- Molecule 3 is a protein called anti-hIL2 FAB fragment light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	217	1672	1037	279	350	6	0	1	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	108	Total 108	O 108	0	0
4	H	224	Total 224	O 224	0	0
4	L	162	Total 162	O 162	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Interleukin-2



- Molecule 2: anti-hIL2 FAB fragment heavy chain



- Molecule 3: anti-hIL2 FAB fragment light chain



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	201.76Å 36.23Å 88.71Å 90.00° 102.93° 90.00°	Depositor
Resolution (Å)	58.74 – 1.95 58.74 – 1.95	Depositor EDS
% Data completeness (in resolution range)	84.9 (58.74-1.95) 84.9 (58.74-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 1.95Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.189 , 0.291 0.196 , 0.293	Depositor DCC
R_{free} test set	1972 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	23.4	Xtrriage
Anisotropy	0.865	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4828	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.10	2/1052 (0.2%)	1.10	3/1423 (0.2%)
2	H	1.28	3/1668 (0.2%)	1.07	2/2283 (0.1%)
3	L	1.02	1/1710 (0.1%)	0.99	3/2327 (0.1%)
All	All	1.14	6/4430 (0.1%)	1.05	8/6033 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	H	0	1
All	All	0	3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	80	GLU	CD-OE1	8.10	1.34	1.25
2	H	35	GLU	CD-OE2	7.07	1.33	1.25
2	H	94	TYR	CG-CD1	6.51	1.47	1.39
2	H	36	TRP	CG-CD1	-5.57	1.28	1.36
1	A	133	THR	CB-CG2	5.54	1.70	1.52
3	L	100	TYR	CE1-CZ	-5.06	1.31	1.38

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	109	ASP	CB-CG-OD1	8.22	125.69	118.30
3	L	98	ASP	CB-CG-OD1	7.23	124.81	118.30
1	A	66	MET	CG-SD-CE	-7.22	88.64	100.20
1	A	58	ARG	NE-CZ-NH1	5.72	123.16	120.30
3	L	215	ARG	NE-CZ-NH2	-5.37	117.61	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	112	ARG	NE-CZ-NH2	-5.35	117.62	120.30
2	H	76	SER	N-CA-CB	-5.29	102.57	110.50
1	A	66	MET	CA-CB-CG	5.02	121.83	113.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	118	GLY	Peptide
1	A	93	ALA	Peptide
2	H	210	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1035	0	1060	22	0
2	H	1627	0	1563	19	0
3	L	1672	0	1570	23	0
4	A	108	0	0	3	0
4	H	224	0	0	5	2
4	L	162	0	0	3	0
All	All	4828	0	4193	60	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:20:VAL:HG22	4:H:335:HOH:O	1.74	0.88
1:A:66:MET:HE1	1:A:137:PHE:HD1	1.50	0.75
2:H:20:VAL:CG2	4:H:335:HOH:O	2.31	0.74
3:L:24:LYS:HE2	3:L:74:ASP:OD1	1.88	0.73
3:L:191:GLU:HG2	3:L:215:ARG:NH2	2.05	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:121:SER:OG	4:H:301:HOH:O	2.15	0.65
3:L:8:PRO:HG3	3:L:11:LEU:HD13	1.78	0.65
3:L:78:ASN:ND2	4:L:301:HOH:O	2.31	0.61
1:A:92:LEU:O	1:A:93:ALA:HB3	2.02	0.59
3:L:170:GLN:HG2	3:L:175:SER:HA	1.83	0.59
3:L:125:SER:OG	3:L:127:GLU:O	2.20	0.58
3:L:191:GLU:HA	3:L:215:ARG:NH2	2.18	0.58
3:L:191:GLU:HG2	3:L:215:ARG:HH22	1.67	0.58
3:L:116:ALA:HB2	3:L:204:THR:OG1	2.04	0.58
3:L:193:HIS:O	3:L:215:ARG:HD2	2.04	0.57
1:A:66:MET:HE1	1:A:137:PHE:CD1	2.35	0.57
2:H:52:ASN:C	2:H:52:ASN:HD22	2.09	0.56
3:L:128:GLN:HG2	3:L:133:GLY:O	2.06	0.56
1:A:66:MET:HE3	1:A:137:PHE:HB2	1.87	0.55
1:A:73:LEU:HD21	1:A:148:ILE:HG21	1.87	0.55
1:A:55:LYS:HB3	1:A:93:ALA:HB2	1.87	0.54
3:L:199:GLU:HG2	3:L:210:VAL:HG22	1.89	0.54
2:H:6:GLN:HE22	2:H:95:PHE:HA	1.74	0.53
2:H:202:THR:HG21	2:H:214:VAL:HG13	1.92	0.52
2:H:196:TRP:CG	2:H:197:PRO:HA	2.45	0.51
1:A:94:GLN:HG3	2:H:103:GLY:C	2.32	0.51
1:A:55:LYS:HG3	1:A:93:ALA:HA	1.93	0.51
1:A:27:THR:HB	4:A:281:HOH:O	2.12	0.49
2:H:160:LEU:C	2:H:160:LEU:HD23	2.33	0.49
2:H:219:GLU:N	4:H:305:HOH:O	2.38	0.49
1:A:28:LYS:HE3	4:A:274:HOH:O	2.12	0.49
1:A:55:LYS:CB	1:A:93:ALA:HB2	2.43	0.48
2:H:135:VAL:HG11	3:L:213:PHE:HB3	1.98	0.46
3:L:128:GLN:NE2	4:L:305:HOH:O	2.49	0.46
1:A:50:ASN:HD22	1:A:51:TYR:N	2.15	0.45
3:L:187:LYS:O	3:L:191:GLU:HG3	2.18	0.44
1:A:99:HIS:CE1	1:A:100:LEU:HG	2.54	0.43
3:L:172:SER:O	3:L:173:LYS:HB3	2.18	0.43
3:L:191:GLU:HA	3:L:215:ARG:CZ	2.48	0.43
1:A:75:HIS:HD2	1:A:123:PHE:CZ	2.36	0.43
2:H:107:TYR:CD1	3:L:50:LEU:HD22	2.54	0.43
2:H:209:ALA:HB1	4:H:422:HOH:O	2.18	0.43
2:H:6:GLN:NE2	2:H:114:GLY:H	2.16	0.42
1:A:124:MET:SD	1:A:124:MET:N	2.91	0.42
1:A:59:MET:SD	1:A:93:ALA:CB	3.07	0.42
1:A:80:GLU:OE1	1:A:117:LYS:NZ	2.46	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:147:GLY:O	2:H:217:LYS:HE2	2.20	0.42
2:H:40:ARG:HA	2:H:41:PRO:HD3	1.92	0.42
1:A:50:ASN:HB2	4:A:214:HOH:O	2.20	0.41
2:H:29:PHE:CD2	2:H:77:SER:HA	2.54	0.41
1:A:92:LEU:HG	2:H:99:TRP:HZ3	1.85	0.41
3:L:125:SER:HB2	4:L:402:HOH:O	2.20	0.41
3:L:16:GLY:HA2	3:L:81:PRO:HB2	2.02	0.41
1:A:118:GLY:C	1:A:120:GLU:H	2.23	0.41
2:H:12:VAL:HG22	2:H:16:THR:OG1	2.21	0.41
3:L:127:GLU:O	3:L:128:GLN:CB	2.69	0.41
3:L:188:ASP:O	3:L:192:ARG:HG3	2.21	0.41
3:L:112:ARG:HD3	3:L:113:ALA:O	2.21	0.40
1:A:53:ASN:C	1:A:53:ASN:HD22	2.25	0.40
1:A:73:LEU:CD2	1:A:148:ILE:HG21	2.50	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:508:HOH:O	4:H:508:HOH:O[2_554]	1.49	0.71
4:H:518:HOH:O	4:H:518:HOH:O[2_554]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	126/132 (96%)	119 (94%)	3 (2%)	4 (3%)	4 0
2	H	217/224 (97%)	205 (94%)	11 (5%)	1 (0%)	29 17
3	L	216/218 (99%)	205 (95%)	8 (4%)	3 (1%)	11 3
All	All	559/574 (97%)	529 (95%)	22 (4%)	8 (1%)	11 3

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	120	GLU
3	L	128	GLN
3	L	173	LYS
1	A	94	GLN
1	A	95	SER
2	H	142	SER
1	A	119	SER
3	L	155	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	119/126 (94%)	110 (92%)	9 (8%)	13	4
2	H	180/187 (96%)	162 (90%)	18 (10%)	7	2
3	L	188/190 (99%)	170 (90%)	18 (10%)	8	2
All	All	487/503 (97%)	442 (91%)	45 (9%)	9	2

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	SER
1	A	31	GLN
1	A	49	ASN
1	A	50	ASN
1	A	53	ASN
1	A	92	LEU
1	A	96	LYS
1	A	124	MET
1	A	150	SER
2	H	1	GLN
2	H	5	GLN
2	H	17	SER
2	H	52	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	85	SER
2	H	121	SER
2	H	135	VAL
2	H	140	THR
2	H	144	VAL
2	H	169	SER
2	H	194	SER
2	H	204	ASN
2	H	210	SER
2	H	212	THR
2	H	215	ASP
2	H	217	LYS
2	H	218	ILE
2	H	219	GLU
3	L	13	VAL
3	L	15	LEU
3	L	22	SER
3	L	112	ARG
3	L	120	SER
3	L	129	LEU
3	L	135	SER
3	L	146	LYS
3	L	149	ASN
3	L	151	LYS
3	L	161	ASN
3	L	173	LYS
3	L	174	ASP
3	L	179	MET
3	L	182	THR
3	L	187	LYS
3	L	195	SER
3	L	207	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	49	ASN
1	A	50	ASN
1	A	53	ASN
1	A	94	GLN
1	A	99	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	1	GLN
2	H	3	GLN
2	H	6	GLN
2	H	31	ASN
2	H	52	ASN
2	H	172	HIS
2	H	207	HIS
3	L	17	GLN
3	L	96	ASN
3	L	128	GLN
3	L	161	ASN
3	L	202	HIS
3	L	214	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	128/132 (96%)	0.95	17 (13%) 3 5	21, 38, 77, 122	0
2	H	219/224 (97%)	0.82	15 (6%) 17 25	20, 34, 56, 121	0
3	L	217/218 (99%)	0.75	23 (10%) 6 10	20, 46, 84, 103	0
All	All	564/574 (98%)	0.82	55 (9%) 7 12	20, 37, 80, 122	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	138	ASP	15.3
1	A	121	THR	14.6
2	H	140	THR	14.2
2	H	137	GLY	11.6
2	H	139	THR	9.3
1	A	119	SER	8.9
3	L	206	THR	7.5
1	A	153	THR	6.5
2	H	141	GLY	6.4
2	H	136	CYS	6.2
3	L	205	SER	5.7
1	A	120	GLU	5.4
1	A	118	GLY	4.9
1	A	122	THR	4.8
3	L	195	SER	4.6
3	L	173	LYS	4.5
3	L	216	ASN	4.1
3	L	155	ASP	3.9
3	L	214	ASN	3.9
1	A	152	LEU	3.8
3	L	172	SER	3.7
3	L	154	ILE	3.7
1	A	123	PHE	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	L	217	GLU	3.6
2	H	142	SER	3.5
2	H	194	SER	3.5
1	A	26	SER	3.5
2	H	192	THR	3.5
3	L	157	SER	3.4
3	L	194	ASN	3.4
1	A	93	ALA	3.3
3	L	212	SER	3.1
2	H	135	VAL	3.1
3	L	207	SER	3.0
3	L	113	ALA	2.9
3	L	161	ASN	2.9
2	H	214	VAL	2.9
1	A	95	SER	2.8
3	L	185	LEU	2.8
1	A	27	THR	2.7
2	H	195	THR	2.7
3	L	12	ALA	2.6
1	A	124	MET	2.6
3	L	193	HIS	2.5
3	L	210	VAL	2.5
1	A	151	THR	2.4
2	H	208	PRO	2.3
1	A	51	TYR	2.3
2	H	143	SER	2.3
3	L	188	ASP	2.3
3	L	115	ALA	2.2
1	A	73	LEU	2.2
1	A	78	CYS	2.2
2	H	50	VAL	2.1
3	L	192	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.