



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 3, 2024 – 02:21 am GMT

PDB ID : 5LYC
Title : Cytochrome c in complex with phosphonato-calix[6]arene
Authors : Rennie, M.L.; Crowley, P.B.
Deposited on : 2016-09-27
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	108	 96%
1	B	108	 98%

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is phosphonato-calix[6]arene (three-letter code: 7AZ) (formula: $C_{42}H_{42}O_{24}P_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			72	42	24	6		
3	B	1	Total	C	O	P	0	0
			72	42	24	6		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	93	Total O 94 94	0	1
5	B	87	Total O 89 89	0	2

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cytochrome c iso-1



- Molecule 1: Cytochrome c iso-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	54.57Å 54.57Å 208.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.78 – 1.80 52.78 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (52.78-1.80) 100.0 (52.78-1.80)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.169 , 0.199 0.179 , 0.206	Depositor DCC
R_{free} test set	1595 reflections (5.27%)	wwPDB-VP
Wilson B-factor (Å ²)	23.8	Xtrriage
Anisotropy	0.129	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2183	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, HEC, 7AZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/893	0.74	0/1191
1	B	0.63	0/918	0.75	1/1225 (0.1%)
All	All	0.65	0/1811	0.75	1/2416 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	91	ARG	NE-CZ-NH2	5.22	122.91	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	863	0	890	3	0
1	B	882	0	909	1	0
2	A	43	0	31	2	0
2	B	43	0	30	3	0
3	A	72	0	0	0	0
3	B	72	0	0	0	0
4	A	10	0	0	0	0
4	B	15	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	94	0	0	1	0
5	B	89	0	0	0	0
All	All	2183	0	1860	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (8) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103:GLU:OE2	5:A:301:HOH:O	2.15	0.64
1:B:94[B]:LEU:HD21	2:B:201:HEC:HBB3	1.86	0.58
2:B:201:HEC:HMC1	2:B:201:HEC:HBC3	1.89	0.54
2:A:201:HEC:HBC3	2:A:201:HEC:HMC1	1.93	0.51
2:B:201:HEC:HBB3	2:B:201:HEC:HMB1	1.95	0.48
1:A:99[B]:LYS:HB3	1:A:99[B]:LYS:HE2	1.83	0.43
2:A:201:HEC:HMB1	2:A:201:HEC:HBB3	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	109/108 (101%)	106 (97%)	3 (3%)	0	100	100
1	B	112/108 (104%)	110 (98%)	2 (2%)	0	100	100
All	All	221/216 (102%)	216 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	92/88 (104%)	92 (100%)	0	100	100
1	B	94/88 (107%)	94 (100%)	0	100	100
All	All	186/176 (106%)	186 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	203	-	4,4,4	0.33	0	6,6,6	0.31	0
4	SO4	B	204	-	4,4,4	0.35	0	6,6,6	0.16	0
3	7AZ	A	202	-	78,78,78	1.39	13 (16%)	126,126,126	1.41	17 (13%)
3	7AZ	B	202	-	78,78,78	1.25	12 (15%)	126,126,126	1.34	15 (11%)
4	SO4	A	204	-	4,4,4	0.30	0	6,6,6	0.15	0
4	SO4	B	203	-	4,4,4	0.29	0	6,6,6	0.21	0
4	SO4	B	205	-	4,4,4	0.32	0	6,6,6	0.05	0
2	HEC	B	201	1	32,50,50	1.72	5 (15%)	24,82,82	1.83	5 (20%)
2	HEC	A	201	1	32,50,50	1.31	4 (12%)	24,82,82	2.03	7 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	7AZ	A	202	-	-	1/60/60/60	-
2	HEC	B	201	1	-	2/10/54/54	-
3	7AZ	B	202	-	-	3/60/60/60	-
2	HEC	A	201	1	-	0/10/54/54	-

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	201	HEC	C2B-C3B	-5.37	1.35	1.40
3	A	202	7AZ	P53-O64	-4.41	1.43	1.54
3	B	202	7AZ	P53-O64	-3.64	1.45	1.54
2	B	201	HEC	CBB-CAB	-3.56	1.36	1.49
2	A	201	HEC	CBC-CAC	-3.56	1.36	1.49
2	B	201	HEC	CBC-CAC	-3.55	1.36	1.49
3	A	202	7AZ	P49-O50	-3.48	1.46	1.54
3	A	202	7AZ	P51-O72	-3.44	1.46	1.54
3	B	202	7AZ	P51-O72	-3.30	1.46	1.54
2	A	201	HEC	C4B-C3B	3.25	1.49	1.43
3	A	202	7AZ	P53-O54	-3.21	1.46	1.54
2	B	201	HEC	C4B-C3B	3.15	1.48	1.43
3	B	202	7AZ	P49-O50	-3.06	1.47	1.54
3	A	202	7AZ	P51-O52	-2.91	1.47	1.54
2	A	201	HEC	CBB-CAB	-2.89	1.38	1.49
2	B	201	HEC	C3C-C2C	-2.89	1.37	1.40
3	B	202	7AZ	P53-O54	-2.74	1.47	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	202	7AZ	P49-C1	-2.68	1.74	1.79
2	A	201	HEC	C2B-C3B	-2.68	1.37	1.40
3	B	202	7AZ	P57-O67	-2.53	1.48	1.54
3	A	202	7AZ	P49-O62	-2.52	1.48	1.54
3	A	202	7AZ	P59-O60	-2.48	1.48	1.54
3	A	202	7AZ	P57-O67	-2.43	1.48	1.54
3	A	202	7AZ	P59-C13	-2.38	1.75	1.79
3	B	202	7AZ	P59-O70	-2.32	1.48	1.54
3	B	202	7AZ	P49-C1	-2.30	1.75	1.79
3	A	202	7AZ	P55-O56	-2.29	1.49	1.54
3	B	202	7AZ	P55-O56	-2.27	1.49	1.54
3	B	202	7AZ	P59-O60	-2.26	1.49	1.54
3	B	202	7AZ	P51-O52	-2.24	1.49	1.54
3	A	202	7AZ	P49-O61	-2.24	1.45	1.49
3	B	202	7AZ	C30-C33	-2.22	1.37	1.39
3	A	202	7AZ	P57-O58	-2.15	1.49	1.54
3	B	202	7AZ	P55-C26	-2.14	1.75	1.79

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	202	7AZ	O54-P53-O65	-4.61	99.31	112.45
2	B	201	HEC	CBA-CAA-C2A	-4.51	105.00	112.60
3	A	202	7AZ	O54-P53-O64	4.47	122.91	107.90
3	B	202	7AZ	O62-P49-C1	4.17	116.28	106.77
2	B	201	HEC	CBD-CAD-C3D	-4.08	105.66	112.62
3	B	202	7AZ	O65-P53-C33	-4.06	101.11	109.69
2	A	201	HEC	CBD-CAD-C3D	-4.02	105.75	112.62
3	A	202	7AZ	O65-P53-C33	-3.97	101.29	109.69
3	B	202	7AZ	O60-P59-C13	3.86	115.58	106.77
2	A	201	HEC	CBA-CAA-C2A	-3.39	106.89	112.60
3	B	202	7AZ	O69-P55-C26	3.36	114.42	106.77
2	A	201	HEC	CMB-C2B-C1B	-3.25	123.46	128.46
2	B	201	HEC	CMB-C2B-C1B	-3.21	123.53	128.46
2	A	201	HEC	CMD-C2D-C1D	-3.19	123.56	128.46
3	A	202	7AZ	O64-P53-O65	-3.15	103.47	112.45
3	A	202	7AZ	O52-P51-C44	3.12	113.88	106.77
2	A	201	HEC	CMD-C2D-C3D	3.03	130.66	124.94
3	B	202	7AZ	O71-P59-C13	-3.01	103.32	109.69
3	A	202	7AZ	O60-P59-C13	3.01	113.63	106.77
3	B	202	7AZ	O52-P51-C44	2.96	113.51	106.77
3	A	202	7AZ	O64-P53-C33	2.83	113.22	106.77

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	202	7AZ	P53-C33-C30	-2.80	116.84	120.49
2	A	201	HEC	CMB-C2B-C3B	2.67	128.96	125.82
2	A	201	HEC	CMC-C2C-C1C	-2.66	124.37	128.46
3	A	202	7AZ	C13-C10-C8	-2.61	118.97	121.31
3	A	202	7AZ	O69-P55-C26	2.59	112.67	106.77
3	B	202	7AZ	P53-C33-C32	2.59	123.87	120.49
3	A	202	7AZ	O50-P49-C1	2.58	112.66	106.77
3	A	202	7AZ	O54-P53-C33	2.49	112.44	106.77
3	A	202	7AZ	O67-P57-C20	2.46	112.38	106.77
2	B	201	HEC	CMD-C2D-C1D	-2.38	124.81	128.46
3	B	202	7AZ	O62-P49-O61	-2.30	105.89	112.45
3	A	202	7AZ	C5-C39-C40	2.29	118.99	112.67
3	B	202	7AZ	O64-P53-C33	2.24	111.88	106.77
3	A	202	7AZ	C14-C15-C17	-2.23	116.80	120.23
3	B	202	7AZ	O58-P57-C20	2.19	111.75	106.77
3	A	202	7AZ	O52-P51-O63	-2.16	106.30	112.45
3	B	202	7AZ	O56-P55-O68	2.13	118.51	112.45
3	B	202	7AZ	C33-C32-C29	-2.12	119.41	121.31
3	A	202	7AZ	C15-C14-C12	2.10	118.47	112.67
2	B	201	HEC	CMA-C3A-C2A	2.09	128.88	124.94
3	B	202	7AZ	C32-C29-C31	2.06	120.95	118.41
3	A	202	7AZ	O60-P59-O70	-2.02	101.14	107.90
3	B	202	7AZ	O68-P55-C26	-2.01	105.44	109.69

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	202	7AZ	C4-C1-P49-O62
2	B	201	HEC	CAA-CBA-CGA-O2A
3	B	202	7AZ	C3-C1-P49-O62
2	B	201	HEC	CAA-CBA-CGA-O1A
3	B	202	7AZ	C4-C1-P49-O61
3	A	202	7AZ	C30-C33-P53-O64

There are no ring outliers.


2 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	201	HEC	3	0
2	A	201	HEC	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand 7AZ B 202







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	107/108 (99%)	-0.17	1 (0%) 81 80	13, 25, 41, 50	4 (3%)
1	B	108/108 (100%)	-0.17	1 (0%) 81 80	11, 24, 39, 51	6 (5%)
All	All	215/216 (99%)	-0.17	2 (0%) 81 80	11, 24, 40, 51	10 (4%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	-5	ALA	4.5
1	A	-4	GLU	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	B	204	5/5	0.74	0.17	77,79,82,82	5
4	SO4	B	203	5/5	0.81	0.15	58,61,65,66	5
4	SO4	A	204	5/5	0.81	0.16	49,50,51,51	5

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	B	205	5/5	0.84	0.15	50,51,52,54	5
4	SO4	A	203	5/5	0.87	0.13	50,56,56,59	5
3	7AZ	B	202	72/72	0.94	0.09	24,30,38,40	0
3	7AZ	A	202	72/72	0.95	0.08	24,30,38,40	0
2	HEC	B	201	43/43	0.99	0.05	15,17,19,21	0
2	HEC	A	201	43/43	0.99	0.05	16,17,21,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 7AZ A 202:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEC B 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.