



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 8, 2024 – 06:15 am GMT

PDB ID : 5M6D
Title : Streptococcus pneumoniae Glyceraldehyde-3-Phosphate Dehydrogenase (SpGAPDH) crystal structure
Authors : Gaboriaud, C.; Moreau, C.P.; Di Guilmi, A.M.
Deposited on : 2016-10-25
Resolution : 2.00 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	355	
1	B	355	

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 10677 atoms, of which 5177 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glyceraldehyde-3-phosphate dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	346	5286	1672	2614	465	524	11	18	10	0
1	B	334	5094	1601	2533	443	507	10	0	9	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP I6L8L9
A	2	ALA	-	expression tag	UNP I6L8L9
A	3	HIS	-	expression tag	UNP I6L8L9
A	4	HIS	-	expression tag	UNP I6L8L9
A	5	HIS	-	expression tag	UNP I6L8L9
A	6	HIS	-	expression tag	UNP I6L8L9
A	7	HIS	-	expression tag	UNP I6L8L9
A	8	HIS	-	expression tag	UNP I6L8L9
A	9	GLY	-	expression tag	UNP I6L8L9
A	10	HIS	-	expression tag	UNP I6L8L9
A	11	HIS	-	expression tag	UNP I6L8L9
A	12	HIS	-	expression tag	UNP I6L8L9
A	13	GLN	-	expression tag	UNP I6L8L9
A	14	LEU	-	expression tag	UNP I6L8L9
A	15	GLU	-	expression tag	UNP I6L8L9
A	16	ASN	-	expression tag	UNP I6L8L9
A	17	LEU	-	expression tag	UNP I6L8L9
A	18	TYR	-	expression tag	UNP I6L8L9
A	19	PHE	-	expression tag	UNP I6L8L9
A	20	GLN	-	expression tag	UNP I6L8L9
A	21	GLY	-	expression tag	UNP I6L8L9
B	1	MET	-	initiating methionine	UNP I6L8L9
B	2	ALA	-	expression tag	UNP I6L8L9
B	3	HIS	-	expression tag	UNP I6L8L9
B	4	HIS	-	expression tag	UNP I6L8L9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	5	HIS	-	expression tag	UNP I6L8L9
B	6	HIS	-	expression tag	UNP I6L8L9
B	7	HIS	-	expression tag	UNP I6L8L9
B	8	HIS	-	expression tag	UNP I6L8L9
B	9	GLY	-	expression tag	UNP I6L8L9
B	10	HIS	-	expression tag	UNP I6L8L9
B	11	HIS	-	expression tag	UNP I6L8L9
B	12	HIS	-	expression tag	UNP I6L8L9
B	13	GLN	-	expression tag	UNP I6L8L9
B	14	LEU	-	expression tag	UNP I6L8L9
B	15	GLU	-	expression tag	UNP I6L8L9
B	16	ASN	-	expression tag	UNP I6L8L9
B	17	LEU	-	expression tag	UNP I6L8L9
B	18	TYR	-	expression tag	UNP I6L8L9
B	19	PHE	-	expression tag	UNP I6L8L9
B	20	GLN	-	expression tag	UNP I6L8L9
B	21	GLY	-	expression tag	UNP I6L8L9

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	B	1	Total Cl 1 1	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0

- Molecule 4 is ACETIC ACID (three-letter code: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			7	2	3	2		
4	B	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	H	O	0	0
			14	3	8	3		
5	B	1	Total	C	H	O	0	0
			14	3	8	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	141	Total	O	0	0
			141	141		
6	B	97	Total	O	0	0
			97	97		

4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	81.04Å 130.24Å 79.84Å 90.00° 119.64° 90.00°	Depositor
Resolution (Å)	40.52 – 2.00 47.48 – 1.91	Depositor EDS
% Data completeness (in resolution range)	99.7 (40.52-2.00) 90.2 (47.48-1.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 1.91Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.195 , 0.215 0.196 , 0.217	Depositor DCC
R_{free} test set	2800 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	28.0	Xtrriage
Anisotropy	0.304	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.019 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10677	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSU, CA, CL, GOL, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	1/2737 (0.0%)	0.57	3/3709 (0.1%)
1	B	0.29	0/2614	0.54	0/3542
All	All	0.32	1/5351 (0.0%)	0.56	3/7251 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	295	ASP	C-N	5.74	1.45	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	10	HIS	CB-CA-C	8.30	127.00	110.40
1	A	10	HIS	C-N-CA	7.60	140.69	121.70
1	A	10	HIS	CA-CB-CG	5.16	122.37	113.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	10	HIS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2672	2614	2630	17	0
1	B	2561	2533	2543	22	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
4	A	4	3	3	0	0
4	B	4	3	3	0	0
5	B	18	24	24	2	0
6	A	141	0	0	3	0
6	B	97	0	0	1	0
All	All	5500	5177	5203	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:171:CSU:HB2	1:B:339:TYR:HB2	1.40	0.98
1:B:51:ARG:NH2	1:B:108:ASP:OD1	2.12	0.82
1:A:171:CSU:HB3	1:A:339:TYR:HB2	1.65	0.75
1:A:40[B]:ARG:NH2	6:A:502:HOH:O	2.23	0.71
1:A:11:HIS:HA	1:A:13:GLN:OE1	1.91	0.70
1:A:40[B]:ARG:NH2	6:A:501:HOH:O	2.20	0.70
1:A:11:HIS:HD2	1:A:13:GLN:OE1	1.75	0.69
1:B:51:ARG:NE	1:B:95:SER:OG	2.26	0.68
1:B:53:ASN:HD21	1:B:103:ILE:HD13	1.58	0.68
1:A:11:HIS:HA	1:A:13:GLN:HG3	1.79	0.64
1:B:218[A]:ARG:NH1	1:B:229[A]:ASN:OD1	2.35	0.60
1:B:335:ASN:ND2	5:B:401:GOL:O1	2.33	0.56
1:A:11:HIS:CD2	1:A:13:GLN:OE1	2.60	0.54
1:A:171:CSU:SG	1:A:333:TYR:HE2	2.33	0.51
1:B:259:THR:CG2	1:B:303:VAL:HG13	2.41	0.51
1:B:136:LYS:HZ3	1:B:355:LYS:HE3	1.76	0.50
1:B:257[B]:THR:HG21	6:B:513:HOH:O	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:100:PRO:HB3	1:B:129:HIS:ND1	2.28	0.48
1:B:51:ARG:NH1	1:B:95:SER:OG	2.46	0.47
1:A:354:ALA:O	1:A:355:LYS:HG2	2.15	0.47
1:B:33:ILE:HD11	5:B:401:GOL:O2	2.15	0.47
1:A:11:HIS:HA	1:A:13:GLN:CG	2.45	0.46
1:B:114:LEU:HG	1:B:347:LEU:HD11	1.97	0.46
1:B:99:ASP:HB3	1:B:102:GLN:HG3	1.97	0.46
1:A:11:HIS:CA	1:A:13:GLN:HG3	2.46	0.45
1:B:105:TRP:HB3	1:B:110:VAL:HB	1.99	0.44
1:A:196[A]:THR:HG21	1:A:333:TYR:OH	2.18	0.44
1:A:105:TRP:HB3	1:A:110:VAL:HB	2.00	0.44
1:B:26:GLY:HA3	1:B:105:TRP:CZ3	2.54	0.42
1:B:55:LEU:HG	1:B:55:LEU:O	2.20	0.42
1:A:257[A]:THR:HG21	6:A:555:HOH:O	2.20	0.42
1:B:53:ASN:O	1:B:53:ASN:OD1	2.38	0.42
1:A:115:GLU:HB3	1:A:139:ILE:HD13	2.01	0.42
1:B:259:THR:HG23	1:B:303:VAL:HG13	2.02	0.42
1:A:142:PRO:HA	1:A:143:GLY:HA3	1.77	0.41
1:B:144:GLY:HA2	1:B:147:VAL:HB	2.02	0.41
1:A:319:LEU:O	1:A:325:GLN:HA	2.21	0.41
1:B:100:PRO:HB2	1:B:128:LYS:CB	2.51	0.40
1:B:104:ASP:OD2	1:B:107:THR:OG1	2.40	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	353/355 (99%)	337 (96%)	15 (4%)	1 (0%)	41 37
1	B	340/355 (96%)	319 (94%)	20 (6%)	1 (0%)	41 37
All	All	693/710 (98%)	656 (95%)	35 (5%)	2 (0%)	41 37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	11	HIS
1	B	143	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/285 (100%)	279 (98%)	7 (2%)	49	51
1	B	275/285 (96%)	270 (98%)	5 (2%)	59	63
All	All	561/570 (98%)	549 (98%)	12 (2%)	59	57

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	98	ARG
1	A	145	ASN
1	A	229	ASN
1	A	257[A]	THR
1	A	257[B]	THR
1	A	333	TYR
1	A	355	LYS
1	B	95	SER
1	B	111	GLU
1	B	114	LEU
1	B	229[A]	ASN
1	B	229[B]	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	HIS
1	B	53	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSU	A	171	1	6,9,10	2.99	2 (33%)	3,12,14	2.32	1 (33%)
1	CSU	B	171	1	6,9,10	3.09	3 (50%)	3,12,14	2.29	2 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSU	A	171	1	-	1/4/8/10	-
1	CSU	B	171	1	-	2/4/8/10	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	171	CSU	OD1-S	6.20	1.64	1.45
1	B	171	CSU	OD3-S	5.86	1.63	1.45
1	B	171	CSU	CB-SG	-3.50	1.75	1.82
1	A	171	CSU	CB-SG	-2.95	1.77	1.82
1	B	171	CSU	OD1-S	-2.86	1.36	1.45

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	171	CSU	OD2-S-OD3	3.43	125.23	112.78

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	171	CSU	OD2-S-OD1	3.01	123.68	112.78
1	B	171	CSU	OD2-S-OD3	-2.33	104.34	112.78

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	171	CSU	N-CA-CB-SG
1	B	171	CSU	N-CA-CB-SG
1	B	171	CSU	OD1-S-SG-CB

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	171	CSU	2	0
1	B	171	CSU	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACY	B	405	-	3,3,3	0.78	0	3,3,3	0.64	0
5	GOL	B	401	-	5,5,5	0.40	0	5,5,5	0.28	0
4	ACY	A	403	-	3,3,3	0.77	0	3,3,3	0.75	0
5	GOL	B	403	-	5,5,5	0.39	0	5,5,5	0.44	0
5	GOL	B	402	-	5,5,5	0.37	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	401	-	-	2/4/4/4	-
5	GOL	B	403	-	-	4/4/4/4	-
5	GOL	B	402	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	402	GOL	O1-C1-C2-O2
5	B	402	GOL	O1-C1-C2-C3
5	B	403	GOL	O1-C1-C2-O2
5	B	403	GOL	O1-C1-C2-C3
5	B	401	GOL	O1-C1-C2-C3
5	B	401	GOL	O1-C1-C2-O2
5	B	403	GOL	O2-C2-C3-O3
5	B	403	GOL	C1-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	401	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	345/355 (97%)	0.21	23 (6%) 17 17	26, 50, 77, 117	0
1	B	333/355 (93%)	1.00	70 (21%) 1 0	24, 47, 100, 126	0
All	All	678/710 (95%)	0.60	93 (13%) 3 2	24, 49, 92, 126	0

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	10	HIS	10.2
1	A	11	HIS	9.1
1	B	131	LYS	7.2
1	B	55	LEU	6.4
1	B	124	GLU	6.4
1	B	102	GLN	6.2
1	B	96	ALA	6.2
1	B	106	ALA	5.9
1	B	130	LEU	5.3
1	B	119	PHE	5.2
1	B	125	ALA	4.9
1	B	107	THR	4.9
1	B	110	VAL	4.8
1	B	101	GLU	4.8
1	B	103	ILE	4.7
1	B	126	ALA	4.7
1	B	127	GLU	4.6
1	B	122	LYS	4.5
1	B	212	ARG	4.5
1	B	109	GLY	4.5
1	A	19	PHE	4.5
1	B	105	TRP	4.4
1	B	135	LYS	4.3
1	A	12	HIS	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	100	PRO	4.1
1	B	98	ARG	4.1
1	B	82	GLU	4.0
1	B	120	PHE	3.9
1	B	97	GLU	3.9
1	A	119	PHE	3.9
1	A	13	GLN	3.8
1	B	132	GLY	3.8
1	B	111	GLU	3.7
1	B	129	HIS	3.6
1	A	122	LYS	3.5
1	B	128	LYS	3.5
1	B	67	TYR	3.5
1	B	134	ALA	3.5
1	B	133	GLY	3.5
1	B	144	GLY	3.4
1	B	121	ALA	3.4
1	B	99	ASP	3.2
1	B	22	VAL	3.1
1	B	354	ALA	3.0
1	B	118	GLY	3.0
1	B	257[A]	THR	3.0
1	A	17	LEU	3.0
1	A	212	ARG	2.9
1	A	224	ALA	2.9
1	A	223	ALA	2.9
1	A	256	PRO	2.8
1	B	123	LYS	2.8
1	A	55	LEU	2.8
1	B	145	ASN	2.8
1	B	146	ASP	2.8
1	B	298	VAL	2.8
1	B	136	LYS	2.8
1	A	226	ILE	2.8
1	A	14	LEU	2.7
1	A	255	VAL	2.7
1	A	227	VAL	2.7
1	A	18	TYR	2.6
1	B	93	LYS	2.6
1	B	300[A]	SER	2.6
1	A	222	GLY	2.5
1	B	53	ASN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	28	ASN	2.5
1	B	164	THR	2.5
1	B	165	VAL	2.5
1	B	303	VAL	2.5
1	B	258	PRO	2.4
1	A	124	GLU	2.4
1	B	355	LYS	2.4
1	B	71	GLN	2.4
1	B	91	PHE	2.3
1	B	36	LEU	2.2
1	B	69	THR	2.2
1	B	255	VAL	2.2
1	B	54	ASP	2.2
1	A	257[A]	THR	2.2
1	B	199	ALA	2.2
1	A	221	ALA	2.2
1	A	89	GLY	2.1
1	B	108	ASP	2.1
1	B	137	VAL	2.1
1	B	72	GLY	2.1
1	B	92	ILE	2.1
1	B	139	ILE	2.0
1	B	45	GLU	2.0
1	B	299	SER	2.0
1	B	112	ILE	2.0
1	B	68	ASP	2.0
1	B	304	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
1	CSU	A	171	10/11	0.85	0.20	43,49,52,52	0
1	CSU	B	171	10/11	0.85	0.20	37,39,41,42	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	B	402	6/6	0.67	0.21	74,89,91,92	0
5	GOL	B	403	6/6	0.69	0.18	56,67,71,72	0
4	ACY	A	403	4/4	0.80	0.14	70,71,84,84	0
5	GOL	B	401	6/6	0.92	0.23	62,74,75,75	0
3	CA	A	402	1/1	0.92	0.07	87,87,87,87	0
4	ACY	B	405	4/4	0.92	0.32	54,54,65,65	0
2	CL	B	404	1/1	0.93	0.14	59,59,59,59	0
2	CL	A	401	1/1	0.98	0.06	41,41,41,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.