

# Full wwPDB X-ray Structure Validation Report (i)

Oct 13, 2024 - 09:36 am BST

PDB ID	:	5OUG
Title	:	Humanized alpha-AChBP (acetylcholine binding protein) in complex with lo-
		beline and allosteric binder fragment 4.
Authors	:	Delbart, F.; Gruss, F.; Ulens, C.
Deposited on	:	2017-08-23
Resolution	:	2.57  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\;DIFFRACTION$ 

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution				
Metric	$(\# { m Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$				
$R_{free}$	164625	4456 (2.60-2.56)				
Clashscore	180529	4905 (2.60-2.56)				
Ramachandran outliers	177936	4847 (2.60-2.56)				
Sidechain outliers	177891	4847 (2.60-2.56)				
RSRZ outliers	164620	4456 (2.60-2.56)				

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	205	2% 	9%
-	D	2005	3%	570
1	В	205	90% <u>3%</u>	9% •
1	С	205	89%	11%
1	D	205	91%	9%
1	Б	205	3%	
	E	205	91%	9%



Mol	Chain	Length	Quality of chain							
2	F	2		100%						
2	J	2	50%	50%						
3	G	5	20%	80%						
4	Н	3		100%						
4	Ι	3		100%						



#### 50UG

# 2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 8950 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf	Trace	
1	Δ	205	Total	С	Ν	0	$\mathbf{S}$	0	0	0	
	А	205	1680	1077	282	314	7	0	0	0	
1	В	205	Total	С	Ν	0	S	0	0	0	
	D	205	1680	1077	282	314	7	0	0	0	
1	С	205	Total	С	Ν	Ο	S	0	0	0	
	U	205	1680	1077	282	314	7	0		0	
1	Л	205	Total	С	Ν	0	S	0	0	0	
	D	205	1680	1077	282	314	7	0	0	0	
1	F	205	Total	С	Ν	0	S	0	0	0	
1	Ľ	205	1676	1075	282	312	7		0		

• Molecule 1 is a protein called Humanized alpha-AChBP.

• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	F	2	Total 28	C 16	N 2	O 10	0	0	0
2	J	2	Total 28	C 16	N 2	O 10	0	0	0

• Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyran ose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.





Mol	Chain	Residues	Atoms		ms		ms		ZeroOcc	AltConf	Trace
3	G	5	Total 61	С 34	N 2	O 25	0	0	0		

• Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-b eta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	Н	3	TotalC3922	N 2	0 15	0	0	0
4	Ι	3	TotalC3922	N 2	0 15	0	0	0

• Molecule 5 is Alpha-Lobeline (three-letter code: L0B) (formula:  $C_{22}H_{27}NO_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total         C         N         O           25         22         1         2	0	0
5	А	1	Total         C         N         O           25         22         1         2	0	0
5	В	1	Total         C         N         O           25         22         1         2	0	0
5	С	1	$\begin{array}{cccc} \text{Total} & \overline{\text{C}} & \text{N} & \text{O} \\ 25 & 22 & 1 & 2 \end{array}$	0	0



Mol	Chain	Residues	A	Aton	ns		ZeroOcc	AltConf
5	D	1	Total 25	C 22	N 1	O 2	0	0

• Molecule 6 is 4,5-dibromo-N-(3-hydroxypropyl)-1H-pyrrole-2-carboxamide (three-letter code: 9Z0) (formula:  $C_8H_{10}Br_2N_2O_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	Λ	1	Total	Br	С	Ν	Ο	0	0
0	Л	T	14	2	8	2	2	0	0
6	Δ	1	Total	Br	С	Ν	0	0	0
0	Λ	I	14	2	8	2	2	0	0
6	В	1	Total	Br	С	Ν	Ο	0	0
0	D	I	14	2	8	2	2	0	0
6	В	1	Total	$\operatorname{Br}$	С	Ν	Ο	0	0
0	D	I	14	2	8	2	2	0	0
6	С	1	Total	$\operatorname{Br}$	С	Ν	Ο	0	0
0	U	I	14	2	8	2	2		0
6	С	1	Total	$\operatorname{Br}$	С	Ν	Ο	0	0
0	0	Ĩ	14	2	8	2	2	0	0
6	Л	1	Total	$\operatorname{Br}$	С	Ν	Ο	0	0
0		1	14	2	8	2	2	0	0
6	Л	1	Total	$\operatorname{Br}$	С	Ν	Ο	0	0
		1	14	2	8	2	2		0
6	E	1	Total	$\operatorname{Br}$	С	Ν	Ο	0	0
		1	14	2	8	2	2		U
6	E	1	Total	$\operatorname{Br}$	С	Ν	Ο	0	0
			14	2	8	2	2		U



• Molecule 7 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula:  $C_2H_6OS$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	Е	1	Total 4	$\begin{array}{c} \mathrm{C} \\ \mathrm{2} \end{array}$	0 1	S 1	0	0

• Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	А	17	Total O 17 17	0	0
8	В	26	TotalO2626	0	0
8	С	19	Total O 19 19	0	0
8	D	18	Total O 18 18	0	0
8	Е	10	Total         O           10         10	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Humanized alpha-AChBP



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc opyranose

Chain F:

100%

#### NAG1 NAG2

• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:	50%	50%
NGC NGC		

 $\bullet$  Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)] beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose nose

Chain G:	20%	80%
NAG1 NAG2 BMA3 MAN4 MAN5		

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:

100%

#### NAG1 NAG2 BMA3

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:

100%

NAG1 NAG2 BMA3



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	68.67Å 119.81Å 84.54Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $107.29^{\circ}$ $90.00^{\circ}$	Depositor
Bosolution(Å)	80.72 - 2.57	Depositor
Resolution (A)	80.72 - 2.57	EDS
% Data completeness	99.7 (80.72-2.57)	Depositor
(in resolution range)	99.9 (80.72-2.57)	EDS
R <sub>merge</sub>	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.54 (at 2.58 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
P. P.	0.214 , $0.240$	Depositor
$n, n_{free}$	0.222 , $0.245$	DCC
$R_{free}$ test set	2107 reflections $(5.07%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	63.6	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34 , $44.3$	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8950	wwPDB-VP
Average B, all atoms $(Å^2)$	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 9Z0, BMA, MAN, DMS, L0B, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.64	0/1724	0.82	4/2344~(0.2%)
1	В	0.64	0/1724	0.86	5/2344~(0.2%)
1	С	0.61	0/1724	0.82	2/2344~(0.1%)
1	D	0.64	0/1724	0.83	3/2344~(0.1%)
1	Е	0.64	0/1720	0.84	4/2339~(0.2%)
All	All	0.63	0/8616	0.83	18/11715~(0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1
1	В	0	1
1	Е	0	1
All	All	0	3

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	В	131	ASP	N-CA-CB	8.34	125.61	110.60
1	D	131	ASP	N-CA-CB	8.33	125.59	110.60
1	Е	122	ARG	NE-CZ-NH2	7.01	123.81	120.30
1	В	122	ARG	NE-CZ-NH2	6.94	123.77	120.30
1	D	120	ARG	NE-CZ-NH1	6.32	123.46	120.30
1	А	120	ARG	NE-CZ-NH1	5.63	123.12	120.30
1	Е	120	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	Е	150	ARG	NE-CZ-NH1	5.40	123.00	120.30



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	4	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	Е	182	ARG	NE-CZ-NH1	5.18	122.89	120.30
1	В	120	ARG	NE-CZ-NH2	-5.16	117.72	120.30
1	А	120	ARG	NE-CZ-NH2	-5.16	117.72	120.30
1	В	120	ARG	NE-CZ-NH1	5.15	122.87	120.30
1	D	150	ARG	NE-CZ-NH1	5.13	122.87	120.30
1	С	150	ARG	NE-CZ-NH2	-5.13	117.73	120.30
1	А	4	ARG	NE-CZ-NH1	5.06	122.83	120.30
1	C	4	ARG	NE-CZ-NH1	5.03	122.81	120.30
1	А	179	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	186	CYS	Peptide
1	В	186	CYS	Peptide
1	Е	186	CYS	Peptide

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1680	0	1651	10	0
1	В	1680	0	1651	11	0
1	С	1680	0	1651	13	0
1	D	1680	0	1651	12	0
1	Е	1676	0	1647	10	0
2	F	28	0	25	0	0
2	J	28	0	25	0	0
3	G	61	0	52	1	0
4	Н	39	0	34	0	0
4	Ι	39	0	34	0	0
5	А	50	0	54	6	0
5	В	25	0	27	0	0
5	С	25	0	27	1	0
5	D	25	0	27	2	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	А	28	0	20	4	0
6	В	28	0	20	4	0
6	С	28	0	20	4	0
6	D	28	0	20	4	0
6	Е	28	0	20	3	0
7	Е	4	0	6	0	0
8	А	17	0	0	0	0
8	В	26	0	0	0	0
8	С	19	0	0	1	0
8	D	18	0	0	0	0
8	Е	10	0	0	0	0
All	All	8950	0	8662	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:D:182:ARG:NH1	1:D:193:ASP:OD2	2.16	0.77
1:C:78:VAL:CG1	1:C:83:LEU:HD11	2.25	0.66
1:C:78:VAL:HG12	1:C:83:LEU:HD11	1.80	0.64
1:D:7:TYR:HA	6:D:305:9Z0:BR2	2.52	0.63
5:C:306:L0B:C5	1:D:116:LEU:HD13	2.30	0.62
1:E:103:GLN:NE2	1:E:115:TYR:OH	2.35	0.59
1:D:103:GLN:NE2	1:D:115:TYR:OH	2.34	0.59
1:C:83:LEU:N	1:C:83:LEU:HD12	2.18	0.59
1:B:7:TYR:HA	6:B:307:9Z0:BR2	2.59	0.58
1:C:159:ALA:HB2	8:C:1104:HOH:O	2.03	0.58
1:C:119:ILE:HD12	6:C:304:9Z0:BR2	2.60	0.56
1:B:119:ILE:HD12	6:B:306:9Z0:BR2	2.61	0.56
1:C:119:ILE:CD1	6:C:304:9Z0:BR2	3.10	0.55
1:C:54:LEU:CD1	6:C:304:9Z0:BR1	3.10	0.54
1:E:7:TYR:HA	6:E:302:9Z0:BR2	2.64	0.52
1:E:25:ARG:HE	1:E:26:PRO:HD2	1.74	0.52
1:D:119:ILE:HD12	6:D:304:9Z0:BR2	2.65	0.52
1:B:186:CYS:SG	1:B:187:CYS:N	2.84	0.51
1:A:119:ILE:CD1	6:A:1104:9Z0:BR2	3.14	0.51
1:A:22:GLN:HG2	1:A:23:ARG:HG3	1.92	0.50
1:A:186:CYS:SG	1:A:187:CYS:N	2.85	0.50
1:C:186:CYS:SG	1:C:187:CYS:N	2.84	0.50



		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:E:186:CYS:SG	1:E:187:CYS:N	2.85	0.50	
5:A:1106:L0B:C7	1:B:116:LEU:HD13	2.42	0.50	
1:D:186:CYS:SG	1:D:187:CYS:N	2.84	0.50	
1:C:7:TYR:HA	6:C:305:9Z0:BR2	2.66	0.50	
1:B:119:ILE:CD1	6:B:306:9Z0:BR2	3.15	0.50	
5:D:306:L0B:C4	1:E:106:LEU:HD12	2.42	0.49	
1:A:106:LEU:HD12	5:A:1101:L0B:C2	2.42	0.49	
1:E:119:ILE:CD1	6:E:301:9Z0:BR2	3.16	0.49	
1:D:121:GLN:OE1	6:D:304:9Z0:BR2	2.86	0.48	
1:D:46:GLN:OE1	1:D:126:ASP:HA	2.13	0.48	
1:A:86:PRO:HB2	1:A:88:ILE:HG12	1.96	0.47	
1:A:7:TYR:HA	6:A:1105:9Z0:BR2	2.69	0.47	
1:D:119:ILE:CD1	6:D:304:9Z0:BR2	3.19	0.45	
1:B:86:PRO:HB2	1:B:88:ILE:HG12	1.98	0.45	
1:C:13:ASN:O	1:D:4:ARG:NH2	2.50	0.45	
1:E:119:ILE:HD12	6:E:301:9Z0:BR2	2.72	0.45	
1:D:115:TYR:C	1:D:116:LEU:HD12	2.38	0.44	
1:C:22:GLN:HE21	1:C:23:ARG:HE	1.66	0.44	
1:D:13:ASN:O	1:E:4:ARG:NH2	2.51	0.44	
1:B:115:TYR:C	1:B:116:LEU:HD12	2.38	0.44	
1:E:86:PRO:HB2	1:E:88:ILE:HG12	1.99	0.44	
1:B:109:SER:OG	3:G:1:NAG:H82	2.18	0.43	
1:C:86:PRO:HB2	1:C:88:ILE:HG13	1.99	0.43	
5:D:306:L0B:H82C	5:D:306:L0B:H221	1.85	0.43	
1:B:54:LEU:CD1	6:B:306:9Z0:BR1	3.21	0.43	
1:A:106:LEU:HD12	5:A:1101:L0B:H2	2.01	0.42	
5:A:1106:L0B:H82C	5:A:1106:L0B:H221	1.85	0.42	
1:A:107:VAL:HG11	6:A:1105:9Z0:BR1	2.74	0.42	
5:A:1106:L0B:C4	1:B:106:LEU:HD12	2.50	0.41	
5:A:1101:L0B:H81C	1:E:145:TRP:O	2.20	0.41	
1:C:123:PHE:CZ	1:C:140:LEU:HD11	2.56	0.41	
1:A:103:GLN:OE1	1:A:115:TYR:OH	2.39	0.40	
1:B:60:ASP:HB3	1:B:63:LEU:HD12	2.03	0.40	
1:A:119:ILE:HD12	6:A:1104:9Z0:BR2	2.77	0.40	

There are no symmetry-related clashes.



### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	203/205~(99%)	199 (98%)	3~(2%)	1 (0%)	25	45
1	В	203/205~(99%)	198 (98%)	4 (2%)	1 (0%)	25	45
1	С	203/205~(99%)	199 (98%)	3(2%)	1 (0%)	25	45
1	D	203/205~(99%)	198 (98%)	4 (2%)	1 (0%)	25	45
1	Е	203/205~(99%)	199 (98%)	3(2%)	1 (0%)	25	45
All	All	1015/1025~(99%)	993 (98%)	17 (2%)	5 (0%)	25	45

All (5) Ramachandran outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type
1	А	24	ASP
1	В	24	ASP
1	С	24	ASP
1	Е	24	ASP
1	D	24	ASP

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	in Analysed Rotameric Outliers		Percentiles		
1	А	192/192~(100%)	189~(98%)	3(2%)	58	78
1	В	192/192~(100%)	189~(98%)	3(2%)	58	78
1	С	192/192~(100%)	188~(98%)	4 (2%)	48	71
1	D	192/192~(100%)	190~(99%)	2(1%)	73	87



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	Ε	191/192 (100%)	190 (100%)	1 (0%)	86 95
All	All	959/960~(100%)	946 (99%)	13 (1%)	62 81

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	37	GLN
1	А	131	ASP
1	А	133	GLU
1	В	25	ARG
1	В	131	ASP
1	В	133	GLU
1	С	25	ARG
1	С	131	ASP
1	С	178	LYS
1	С	182	ARG
1	D	23	ARG
1	D	131	ASP
1	Е	131	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	177	GLN
1	В	37	GLN
1	В	61	HIS
1	С	3	GLN
1	С	22	GLN
1	С	37	GLN
1	С	61	HIS
1	С	177	GLN
1	D	37	GLN
1	D	103	GLN
1	D	121	GLN
1	D	177	GLN
1	Е	37	GLN
1	Е	61	HIS
1	Е	103	GLN
1	Е	157	GLN
1	Е	177	GLN



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

15 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tuno	Chain	Dog	Link	Bo	ond leng	ths	В	ond ang	les
	Type	Ullaili	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
2	NAG	F	1	1,2	14,14,15	0.42	0	17,19,21	1.09	2 (11%)
2	NAG	F	2	2	14,14,15	0.54	0	17,19,21	1.36	2 (11%)
3	NAG	G	1	1,3	14,14,15	0.47	0	17,19,21	0.96	0
3	NAG	G	2	3	14,14,15	0.43	0	17,19,21	0.79	0
3	BMA	G	3	3	11,11,12	0.46	0	$15,\!15,\!17$	1.22	2 (13%)
3	MAN	G	4	3	11,11,12	0.53	0	$15,\!15,\!17$	1.03	1 (6%)
3	MAN	G	5	3	11,11,12	0.32	0	$15,\!15,\!17$	1.06	1 (6%)
4	NAG	Н	1	1,4	14,14,15	0.40	0	17,19,21	1.34	3 (17%)
4	NAG	Н	2	4	14,14,15	0.49	0	17,19,21	1.04	1 (5%)
4	BMA	Н	3	4	11,11,12	0.62	0	$15,\!15,\!17$	1.00	1 (6%)
4	NAG	Ι	1	1,4	14,14,15	0.50	0	17,19,21	1.27	1 (5%)
4	NAG	Ι	2	4	14,14,15	0.66	0	17,19,21	1.38	3 (17%)
4	BMA	Ι	3	4	11,11,12	0.67	0	$15,\!15,\!17$	1.48	2 (13%)
2	NAG	J	1	1,2	14,14,15	0.48	0	17,19,21	1.31	2 (11%)
2	NAG	J	2	2	14,14,15	0.36	0	17,19,21	1.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	F	1	1,2	_	0/6/23/26	0/1/1/1
2	NAG	F	2	2	-	0/6/23/26	0/1/1/1
3	NAG	G	1	1,3	_	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	0/6/23/26	0/1/1/1
3	BMA	G	3	3	-	0/2/19/22	0/1/1/1
3	MAN	G	4	3	-	0/2/19/22	0/1/1/1
3	MAN	G	5	3	-	0/2/19/22	0/1/1/1
4	NAG	Н	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	Н	2	4	-	2/6/23/26	0/1/1/1
4	BMA	Н	3	4	-	0/2/19/22	0/1/1/1
4	NAG	Ι	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	Ι	2	4	-	0/6/23/26	0/1/1/1
4	BMA	Ι	3	4	-	0/2/19/22	0/1/1/1
2	NAG	J	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	J	2	2	-	2/6/23/26	0/1/1/1

'-' means no outliers of that kind were identified.

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	J	1	NAG	O5-C1-C2	-3.48	105.79	111.29
4	Н	1	NAG	O5-C1-C2	-3.38	105.94	111.29
4	Ι	3	BMA	C3-C4-C5	3.24	116.02	110.24
4	Ι	1	NAG	O5-C1-C2	-3.23	106.19	111.29
3	G	3	BMA	C1-O5-C5	3.19	116.52	112.19
4	Ι	3	BMA	C1-O5-C5	3.13	116.43	112.19
2	F	2	NAG	C4-C3-C2	3.11	115.58	111.02
3	G	5	MAN	C1-O5-C5	2.78	115.96	112.19
2	F	2	NAG	C3-C4-C5	2.61	114.89	110.24
3	G	3	BMA	C1-C2-C3	2.53	112.78	109.67
2	J	1	NAG	C1-O5-C5	2.44	115.50	112.19
4	Н	3	BMA	C1-C2-C3	2.38	112.59	109.67
3	G	4	MAN	C1-O5-C5	2.35	115.38	112.19
2	F	1	NAG	O5-C1-C2	-2.33	107.61	111.29
4	Н	1	NAG	C1-O5-C5	2.27	115.27	112.19
4	Н	1	NAG	C1-C2-N2	2.22	114.27	110.49
4	Ι	2	NAG	C3-C4-C5	-2.19	106.33	110.24
4	Н	2	NAG	C2-N2-C7	2.17	126.00	122.90
2	F	1	NAG	C1-C2-N2	2.07	114.03	110.49
4	Ι	2	NAG	O4-C4-C5	2.04	114.35	109.30
4	Ι	2	NAG	C4-C3-C2	2.02	113.98	111.02



There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	J	2	NAG	O5-C5-C6-O6
2	J	2	NAG	C4-C5-C6-O6
4	Н	2	NAG	C4-C5-C6-O6
4	Н	2	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





















## 5.6 Ligand geometry (i)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Chain		Link	Bo	ond leng	$_{\rm ths}$	Bond angles				
MOI	туре	Unain	nes	nes	res Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	9Z0	В	306	-	12,14,14	0.98	1 (8%)	10,18,18	0.90	0	
6	9Z0	С	305	-	12,14,14	0.87	1 (8%)	10,18,18	1.31	2 (20%)	
5	L0B	А	1106	-	27,27,27	0.54	0	33,36,36	1.19	1 (3%)	
6	9Z0	А	1104	-	12,14,14	0.91	1 (8%)	10,18,18	0.80	0	
6	9Z0	E	302	-	12,14,14	0.92	1 (8%)	10,18,18	0.85	0	



Mal	Tuno	Chain	in Res Link		Bo	Bond lengths			Bond angles		
IVIOI	Type	Ullalli	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2	
5	L0B	С	306	-	27,27,27	0.49	0	33,36,36	1.20	2 (6%)	
6	9Z0	В	307	-	12,14,14	0.88	1 (8%)	10,18,18	0.96	0	
6	9Z0	А	1105	-	12,14,14	0.87	1 (8%)	10,18,18	0.95	0	
7	DMS	Е	303	-	3,3,3	0.54	0	3,3,3	0.74	0	
6	9Z0	Е	301	-	12,14,14	1.00	1 (8%)	10,18,18	0.98	0	
5	L0B	В	308	-	27,27,27	0.47	0	33,36,36	1.17	3 (9%)	
5	L0B	D	306	-	27,27,27	0.50	0	33,36,36	1.25	2 (6%)	
6	9Z0	D	304	-	12,14,14	1.07	1 (8%)	10,18,18	0.91	1 (10%)	
6	9Z0	D	305	-	12,14,14	0.80	1 (8%)	10,18,18	0.95	0	
5	L0B	А	1101	-	27,27,27	0.49	0	33,36,36	1.27	3 (9%)	
6	9Z0	С	304	-	12,14,14	1.01	1 (8%)	10,18,18	0.93	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	9Z0	В	306	-	-	1/6/9/9	0/1/1/1
6	9Z0	С	305	-	-	2/6/9/9	0/1/1/1
5	L0B	А	1106	-	-	4/16/30/30	0/3/3/3
6	9Z0	А	1104	-	-	2/6/9/9	0/1/1/1
6	9Z0	Е	302	-	-	1/6/9/9	0/1/1/1
5	L0B	С	306	-	-	6/16/30/30	0/3/3/3
6	9Z0	В	307	-	-	2/6/9/9	0/1/1/1
6	9Z0	А	1105	-	-	2/6/9/9	0/1/1/1
6	9Z0	Е	301	-	-	2/6/9/9	0/1/1/1
5	L0B	В	308	-	-	9/16/30/30	0/3/3/3
5	L0B	D	306	-	-	8/16/30/30	0/3/3/3
6	9Z0	D	304	-	-	3/6/9/9	0/1/1/1
6	9Z0	D	305	-	-	1/6/9/9	0/1/1/1
5	L0B	А	1101	-	-	4/16/30/30	0/3/3/3
6	9Z0	C	304	-	-	0/6/9/9	0/1/1/1

All (10) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
6	С	304	9Z0	BR2-C12	-2.81	1.85	1.90
6	Е	301	9Z0	BR2-C12	-2.77	1.85	1.90
6	В	306	9Z0	BR2-C12	-2.67	1.86	1.90
6	А	1104	9Z0	BR2-C12	-2.60	1.86	1.90
6	D	304	9Z0	BR2-C12	-2.54	1.86	1.90
6	С	305	9Z0	BR2-C12	-2.30	1.86	1.90
6	В	307	9Z0	BR2-C12	-2.18	1.86	1.90
6	Е	302	9Z0	BR2-C12	-2.18	1.86	1.90
6	D	305	9Z0	BR2-C12	-2.13	1.86	1.90
6	А	1105	9Z0	BR2-C12	-2.07	1.86	1.90

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	А	1106	L0B	C22-N1-C11	4.89	117.11	113.16
5	D	306	L0B	C22-N1-C11	4.74	116.99	113.16
5	В	308	L0B	C22-N1-C11	4.39	116.71	113.16
5	А	1101	L0B	C22-N1-C11	4.30	116.63	113.16
5	С	306	LOB	C22-N1-C11	3.96	116.36	113.16
5	С	306	L0B	C22-N1-C9	3.12	115.68	113.16
5	А	1101	LOB	C22-N1-C9	2.57	115.23	113.16
6	С	305	9Z0	C08-C02-N03	2.53	119.13	115.59
6	С	305	9Z0	BR1-C10-C12	-2.40	125.02	127.63
6	D	304	9Z0	BR1-C10-C12	-2.25	125.18	127.63
5	А	1101	LOB	C8-C3-C1	2.13	121.01	118.71
5	В	308	LOB	C22-N1-C9	2.09	114.84	113.16
5	В	308	LOB	C14-C11-C13	2.08	116.82	113.03
5	D	306	LOB	C14-C11-C13	2.08	116.81	113.03

There are no chirality outliers.

All (47) torsion outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	Atoms
5	А	1101	LOB	C11-C14-C16-C17
5	А	1101	L0B	C11-C14-C16-O2
5	А	1106	LOB	C11-C14-C16-C17
5	А	1106	LOB	C11-C14-C16-O2
5	В	308	LOB	C11-C14-C16-C17
5	В	308	LOB	C11-C14-C16-O2
5	С	306	LOB	C11-C14-C16-C17
5	С	306	L0B	C11-C14-C16-O2
5	С	306	L0B	C3-C8-C9-C12



Mol	Chain	Res	Type	Atoms
5	D	306	LOB	C11-C14-C16-C17
5	D	306	LOB	C11-C14-C16-O2
6	А	1104	9Z0	C08-C02-N03-C04
6	А	1105	9Z0	N03-C04-C05-C06
6	В	306	9Z0	N03-C04-C05-C06
6	В	307	9Z0	N03-C04-C05-C06
6	С	305	9Z0	C04-C05-C06-O07
6	D	305	9Z0	N03-C04-C05-C06
6	А	1104	9Z0	O01-C02-N03-C04
6	В	307	9Z0	C04-C05-C06-O07
6	Е	302	9Z0	C04-C05-C06-O07
6	Е	301	9Z0	C08-C02-N03-C04
5	А	1101	L0B	O1-C3-C8-C9
5	В	308	L0B	O1-C3-C8-C9
5	D	306	L0B	O1-C3-C8-C9
5	А	1106	L0B	C1-C3-C8-C9
5	В	308	L0B	C1-C3-C8-C9
5	С	306	L0B	C1-C3-C8-C9
5	D	306	L0B	C1-C3-C8-C9
6	D	304	9Z0	C08-C02-N03-C04
6	Е	301	9Z0	O01-C02-N03-C04
6	D	304	9Z0	C04-C05-C06-O07
5	А	1106	L0B	O1-C3-C8-C9
5	С	306	LOB	O1-C3-C8-C9
5	А	1101	L0B	C1-C3-C8-C9
6	D	304	9Z0	O01-C02-N03-C04
5	В	308	L0B	C3-C8-C9-C12
6	С	305	9Z0	C08-C02-N03-C04
5	В	308	LOB	C2-C1-C3-O1
5	D	306	LOB	C2-C1-C3-O1
5	В	308	L0B	C4-C1-C3-O1
5	D	306	LOB	C4-C1-C3-O1
5	В	308	LOB	C2-C1-C3-C8
5	С	306	LOB	C4-C1-C3-O1
5	D	306	LOB	C2-C1-C3-C8
5	В	308	LOB	C4-C1-C3-C8
5	D	306	LOB	C4-C1-C3-C8
6	А	1105	9Z0	C04-C05-C06-O07

Continued from previous page...

There are no ring outliers.

14 monomers are involved in 28 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	В	306	9Z0	3	0
6	С	305	9Z0	1	0
5	А	1106	L0B	3	0
6	А	1104	9Z0	2	0
6	Е	302	9Z0	1	0
5	С	306	L0B	1	0
6	В	307	9Z0	1	0
6	А	1105	9Z0	2	0
6	Е	301	9Z0	2	0
5	D	306	L0B	2	0
6	D	304	9Z0	3	0
6	D	305	9Z0	1	0
5	А	1101	L0B	3	0
6	С	304	9Z0	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.













![](_page_30_Picture_4.jpeg)

![](_page_31_Figure_3.jpeg)

![](_page_31_Picture_4.jpeg)

![](_page_32_Figure_3.jpeg)

![](_page_32_Picture_4.jpeg)

![](_page_33_Figure_3.jpeg)

![](_page_33_Picture_4.jpeg)

![](_page_34_Figure_3.jpeg)

![](_page_34_Picture_4.jpeg)

![](_page_35_Figure_3.jpeg)

# 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

![](_page_35_Picture_8.jpeg)

# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	205/205~(100%)	0.14	4 (1%) 64 61	42, 68, 111, 131	0
1	В	205/205~(100%)	0.24	7 (3%) 48 44	44, 65, 115, 133	0
1	C	205/205~(100%)	0.25	6 (2%) 54 49	43, 71, 121, 146	0
1	D	205/205~(100%)	0.27	7 (3%) 48 44	46, 72, 122, 134	0
1	E	205/205~(100%)	0.19	6 (2%) 54 49	46, 68, 114, 126	0
All	All	1025/1025~(100%)	0.22	30 (2%) 54 49	42, 69, 117, 146	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	36	LEU	3.5
1	В	36	LEU	3.4
1	В	161	ILE	3.4
1	D	161	ILE	3.4
1	D	159	ALA	3.3
1	А	11	VAL	3.3
1	D	67	VAL	3.2
1	А	204	GLY	3.1
1	Ε	0	GLY	3.0
1	В	67	VAL	2.9
1	D	3	GLN	2.9
1	Ε	1	GLU	2.7
1	В	11	VAL	2.7
1	D	11	VAL	2.6
1	Е	161	ILE	2.6
1	С	0	GLY	2.6
1	Е	11	VAL	2.6
1	А	159	ALA	2.6
1	В	45	ASN	2.5
1	Е	186	CYS	2.5

![](_page_36_Picture_10.jpeg)

	v	1	1 0		
Mol	Chain	Res	Type	RSRZ	
1	С	159	ALA	2.5	
1	В	0	GLY	2.4	
1	С	161	ILE	2.4	
1	С	173	VAL	2.3	
1	С	204	GLY	2.3	
1	D	0	GLY	2.3	
1	D	2	PHE	2.2	
1	В	159	ALA	2.2	
1	А	161	ILE	2.2	
1	Е	13	ASN	2.2	

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
4	BMA	Ι	3	11/12	0.48	0.15	104,122,125,126	0
4	BMA	Н	3	11/12	0.62	0.12	90,104,111,114	0
4	NAG	Н	2	14/15	0.74	0.13	96,111,115,116	0
2	NAG	F	2	14/15	0.75	0.13	70,110,116,118	0
2	NAG	J	2	14/15	0.76	0.12	94,120,129,130	0
4	NAG	Ι	2	14/15	0.81	0.11	87,108,118,120	0
3	MAN	G	4	11/12	0.82	0.10	77,88,94,96	0
3	BMA	G	3	11/12	0.83	0.08	81,89,90,91	0
4	NAG	Н	1	14/15	0.83	0.10	73,86,98,106	0
3	MAN	G	5	11/12	0.85	0.10	86,87,92,93	0
2	NAG	F	1	14/15	0.88	0.08	62,74,83,88	0
3	NAG	G	2	14/15	0.90	0.07	64,78,84,87	0
3	NAG	G	1	14/15	0.91	0.08	58,63,71,76	0
4	NAG	Ι	1	14/15	0.91	0.08	68,77,86,92	0
2	NAG	J	1	14/15	0.92	0.07	73,82,88,97	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

![](_page_37_Picture_11.jpeg)

![](_page_38_Figure_3.jpeg)

![](_page_39_Figure_3.jpeg)

![](_page_39_Picture_4.jpeg)

![](_page_40_Figure_3.jpeg)

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
5	L0B	В	308	25/25	0.73	0.27	82,108,123,126	0
6	9Z0	D	305	14/14	0.73	0.26	92,103,159,160	14
5	L0B	А	1106	25/25	0.74	0.24	84,104,114,119	0
5	L0B	С	306	25/25	0.76	0.21	82,102,111,120	0
6	9Z0	D	304	14/14	0.77	0.34	69,94,106,125	14
5	L0B	А	1101	25/25	0.79	0.24	77,98,116,118	0
5	L0B	D	306	25/25	0.79	0.26	92,110,118,120	0
6	9Z0	В	307	14/14	0.80	0.24	92,112,136,174	14
6	9Z0	Е	301	14/14	0.80	0.32	60,95,115,115	14
6	9Z0	С	304	14/14	0.81	0.29	70,87,101,117	14
6	9Z0	Е	302	14/14	0.81	0.30	115,161,214,246	14
6	9Z0	А	1104	14/14	0.82	0.33	72,93,125,130	14
6	9Z0	А	1105	14/14	0.82	0.22	81,94,117,125	14
6	9Z0	В	306	14/14	0.83	0.29	64,91,107,118	14

![](_page_40_Picture_8.jpeg)

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	9Z0	С	305	14/14	0.84	0.20	72,79,105,121	14
7	DMS	Ε	303	4/4	0.87	0.19	110,112,113,123	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

![](_page_41_Figure_6.jpeg)

![](_page_41_Picture_7.jpeg)

![](_page_42_Figure_3.jpeg)

![](_page_42_Picture_4.jpeg)

![](_page_43_Figure_3.jpeg)

![](_page_43_Picture_4.jpeg)

![](_page_44_Figure_3.jpeg)

![](_page_44_Picture_4.jpeg)

![](_page_45_Figure_3.jpeg)

![](_page_45_Picture_4.jpeg)

![](_page_46_Figure_3.jpeg)

![](_page_46_Picture_4.jpeg)

![](_page_47_Figure_3.jpeg)

![](_page_47_Picture_4.jpeg)

![](_page_48_Figure_3.jpeg)

![](_page_48_Picture_4.jpeg)

![](_page_49_Figure_3.jpeg)

![](_page_49_Picture_4.jpeg)

![](_page_50_Figure_3.jpeg)

# 6.5 Other polymers (i)

There are no such residues in this entry.

![](_page_50_Picture_6.jpeg)