

Full wwPDB X-ray Structure Validation Report (i)

Jun 18, 2024 – 06:16 AM EDT

PDB ID : 5RKZ

Title : PanDDA analysis group deposition of ground-state model of human NUDT22

screened against the DSPL fragment library by X-ray Crystallography

Authors: Diaz-Saez, L.; Talon, R.; Krojer, T.; Burgess-Brown, N.A.; Arrowsmith, C.H.;

Edwards, A.M.; Bountra, C.; von Delft, F.; Huber, K.V.M.

Deposited on : 2020-06-19

Resolution : 1.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : FAILED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	141614	3037 (1.40-1.36)
Ramachandran outliers	138981	2970 (1.40-1.36)
Sidechain outliers	138945	2969 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain	
1	A	304	86%	9% • •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	DMS	A	401	-	-	X	-
2	DMS	A	402	-	-	X	-



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2419 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called Uridine diphosphate glucose pyrophosphatase NUDT22.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	291	Total 2241	C 1421	N 387	O 428	S 5	0	7	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9BRQ3
A	260	ARG	GLN	conflict	UNP Q9BRQ3
A	263	PRO	LEU	conflict	UNP Q9BRQ3

• Molecule 2 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total C O 4 2 1	S 1	0	0
2	A	1	Total C O 4 2 1	S 1	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O S 4 2 1 1	0	0
2	A	1	Total C O S 4 2 1 1	0	0

• Molecule 3 is water.

Mo	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	162	Total O 162 162	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.

• Molecule 1: Uridine diphosphate glucose pyrophosphatase NUDT22





4 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	49.55Å 52.39Å 101.73Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.92 - 1.38	Depositor
% Data completeness	99.8 (50.92-1.38)	Depositor
(in resolution range)	33.0 (90.32-1.90)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.45 (at 1.38Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.183 , 0.211	Depositor
Wilson B-factor (\mathring{A}^2)	17.7	Xtriage
Anisotropy	0.088	Xtriage
L-test for twinning ²	$ < L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2419	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.82% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Boı	nd lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	1.17	$2/2307 \ (0.1\%)$	1.13	$15/3152 \ (0.5\%)$	

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
1	A	255	GLU	CD-OE1	8.81	1.35	1.25
1	A	224	TYR	CE1-CZ	-7.17	1.29	1.38

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}({}^o)$	$\operatorname{Ideal}({}^{o})$
1	A	235	ARG	NE-CZ-NH1	10.25	125.42	120.30
1	A	211	ARG	NE-CZ-NH1	-10.20	115.20	120.30
1	A	211	ARG	NE-CZ-NH2	10.13	125.36	120.30
1	A	88	ARG	NE-CZ-NH1	8.38	124.49	120.30
1	A	62	ARG	NE-CZ-NH1	7.48	124.04	120.30
1	A	141	ARG	NE-CZ-NH1	6.35	123.47	120.30
1	A	141	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	A	218	ARG	NE-CZ-NH2	-5.98	117.31	120.30
1	A	222	GLU	OE1-CD-OE2	-5.78	116.37	123.30
1	A	284	ARG	NE-CZ-NH2	-5.72	117.44	120.30
1	A	81	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	A	88	ARG	CG-CD-NE	5.48	123.31	111.80
1	A	261	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	109	ASP	CB-CG-OD1	5.28	123.05	118.30
1	A	255	GLU	CG-CD-OE1	5.17	128.65	118.30

There are no chirality outliers.

There are no planarity outliers.



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2241	0	2196	26	0
2	A	16	0	24	11	0
3	A	162	0	0	8	4
All	All	2419	0	2220	28	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:ARG:CD	3:A:501:HOH:O	1.67	1.18
1:A:88:ARG:HD3	3:A:501:HOH:O	1.42	1.00
1:A:211:ARG:HD3	1:A:213:GLU:OE2	1.69	0.91
1:A:141:ARG:HE	2:A:401:DMS:H13	1.53	0.71
1:A:198:LEU:CD1	2:A:402:DMS:H11	2.22	0.70
1:A:4:GLU:HB2	1:A:214:THR:HG23	1.73	0.70
1:A:211:ARG:CD	1:A:213:GLU:OE2	2.39	0.69
2:A:401:DMS:H12	3:A:521:HOH:O	1.98	0.63
1:A:42:THR:OG1	1:A:88:ARG:NH2	2.32	0.62
1:A:71:ILE:HG23	3:A:504:HOH:O	2.00	0.61
1:A:18:GLN:HE22	1:A:180:HIS:HD2	1.49	0.61
1:A:71:ILE:N	3:A:504:HOH:O	2.39	0.56
1:A:145:GLU:OE2	1:A:218:ARG:NH2	2.40	0.55
1:A:6:THR:OG1	1:A:211:ARG:HD2	2.09	0.53
1:A:198:LEU:O	2:A:402:DMS:H12	2.11	0.51
1:A:18:GLN:HE22	1:A:180:HIS:CD2	2.27	0.50
1:A:2[B]:ASP:OD1	1:A:4:GLU:OE1	2.30	0.50
1:A:88:ARG:CG	3:A:501:HOH:O	2.33	0.48
1:A:211:ARG:HD3	1:A:213:GLU:CD	2.34	0.47
1:A:274:SER:OG	2:A:404:DMS:O	2.24	0.47
1:A:4:GLU:HG3	1:A:273:PRO:HG3	1.97	0.46
1:A:141:ARG:HE	2:A:401:DMS:C1	2.25	0.46
1:A:198:LEU:HD12	2:A:402:DMS:H11	1.97	0.46
1:A:198:LEU:HD11	2:A:402:DMS:H11	1.97	0.45

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:124:GLY:HA3	2:A:404:DMS:H13	1.99	0.44
1:A:141:ARG:H	2:A:401:DMS:C1	2.30	0.44
1:A:243:PRO:O	3:A:502:HOH:O	2.21	0.42
2:A:401:DMS:C1	3:A:521:HOH:O	2.62	0.41

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
3:A:517:HOH:O	3:A:572:HOH:O[3_655]	0.78	1.42
3:A:582:HOH:O	3:A:649:HOH:O[4_565]	1.83	0.37
3:A:539:HOH:O	3:A:649:HOH:O[4_565]	2.03	0.17
3:A:516:HOH:O	3:A:584:HOH:O[2_564]	2.19	0.01

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed Outliers		Percentiles	
1	A	293/304 (96%)	287 (98%)	4 (1%)	2 (1%)	22 5	

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	132[A]	ASP
1	A	132[B]	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Rotameric Outliers		Percentiles	
1	A	236/244 (97%)	234 (99%)	2 (1%)	81 61

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	211	ARG
1	A	232	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	GLN
1	A	77	GLN
1	A	176	GLN
1	A	180	HIS
1	A	203	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	ol Type Chain Res L		Link	Bond lengths			Bond angles			
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DMS	A	404	-	3,3,3	0.73	0	3,3,3	1.10	0
2	DMS	A	402	-	3,3,3	0.72	0	3,3,3	0.95	0
2	DMS	A	403	-	3,3,3	0.68	0	3,3,3	0.83	0
2	DMS	A	401	-	3,3,3	0.82	0	3,3,3	0.79	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	404	DMS	2	0
2	A	402	DMS	4	0
2	A	401	DMS	5	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

6.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

6.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

6.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

