



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 8, 2024 – 08:57 PM EDT

PDB ID : 5SFF
Title : CRYSTAL STRUCTURE OF HUMAN PHOSPHODIESTERASE 10 IN COMPLEX WITH c1cc(nn2c1nc(c2CO)C)C#Cc3nc(en3C)c4cccc4, micromolar IC₅₀=0.00103
Authors : Joseph, C.; Groebke-Zbinden, K.; Benz, J.; Schlatter, D.; Rudolph, M.G.
Deposited on : 2022-01-21
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1881 (2.16-2.16)
Clashscore	180529	2047 (2.16-2.16)
Ramachandran outliers	177936	2027 (2.16-2.16)
Sidechain outliers	177891	2026 (2.16-2.16)
RSRZ outliers	164620	1882 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	 80% 11% 9%
1	B	343	 81% 11% 7%
1	C	343	 83% 8% 9%
1	D	343	 83% 9% 8%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10720 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	313	Total 2549	C 1629	N 435	O 461	S 24	0	1	0
1	B	320	Total 2600	C 1660	N 441	O 475	S 24	0	0	0
1	C	313	Total 2549	C 1629	N 435	O 461	S 24	0	1	0
1	D	317	Total 2577	C 1647	N 438	O 468	S 24	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	447	GLY	-	expression tag	UNP Q9Y233
A	448	SER	-	expression tag	UNP Q9Y233
B	447	GLY	-	expression tag	UNP Q9Y233
B	448	SER	-	expression tag	UNP Q9Y233
C	447	GLY	-	expression tag	UNP Q9Y233
C	448	SER	-	expression tag	UNP Q9Y233
D	447	GLY	-	expression tag	UNP Q9Y233
D	448	SER	-	expression tag	UNP Q9Y233

- Molecule 2 is {(4S)-2-methyl-6-[(1-methyl-4-phenyl-1H-imidazol-2-yl)ethynyl]imidazo[1,2-b]pyridazin-3-yl}methanol (three-letter code: IKH) (formula: C₂₀H₁₇N₅O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			26	20	5	1		
2	B	1	Total	C	N	O	0	0
			26	20	5	1		
2	C	1	Total	C	N	O	0	0
			26	20	5	1		
2	D	1	Total	C	N	O	0	0
			26	20	5	1		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Mg 1	0	0
4	C	1	Total 1	Mg 1	0	0
4	D	1	Total 1	Mg 1	0	0


- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	95	Total 95	O 95	0	0
5	B	105	Total 105	O 105	0	0
5	C	104	Total 104	O 104	0	0
5	D	29	Total 29	O 29	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A

Chain A: 




- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A

Chain B: 




- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A


Chain C: 



- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A

Chain D: 





4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	134.93Å 134.93Å 235.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.41 – 2.16 43.41 – 2.16	Depositor EDS
% Data completeness (in resolution range)	96.3 (43.41-2.16) 96.3 (43.41-2.16)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 2.16Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.173 , 0.227 0.181 , 0.231	Depositor DCC
R_{free} test set	4287 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	41.6	Xtrriage
Anisotropy	0.156	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.027 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10720	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IKH, ZN, MG, CME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/2603	0.85	0/3521
1	B	0.74	1/2653 (0.0%)	0.85	0/3591
1	C	0.75	0/2603	0.85	0/3521
1	D	0.79	1/2630 (0.0%)	0.80	0/3560
All	All	0.76	2/10489 (0.0%)	0.84	0/14193

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	455	GLU	CD-OE2	6.26	1.32	1.25
1	B	455	GLU	CD-OE2	5.33	1.31	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2549	0	2524	25	0
1	B	2600	0	2555	25	0
1	C	2549	0	2524	14	0
1	D	2577	0	2541	17	0
2	A	26	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	26	0	0	0	0
2	C	26	0	0	0	0
2	D	26	0	0	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	95	0	0	2	0
5	B	105	0	0	2	0
5	C	104	0	0	3	0
5	D	29	0	0	0	0
All	All	10720	0	10144	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:469:CYS:SG	5:A:989:HOH:O	2.19	0.99
1:C:469:CYS:SG	5:C:995:HOH:O	2.32	0.85
1:B:509:CME:HB2	5:B:992:HOH:O	1.88	0.73
1:C:716:ARG:O	1:C:719:LYS:HG3	1.94	0.68
1:A:461:GLN:HA	1:A:461:GLN:OE1	1.93	0.67
1:B:546:THR:HG21	1:D:767:ARG:NH2	2.14	0.63
1:D:740:THR:HA	1:D:743:GLN:HE21	1.66	0.61
1:C:714:MET:HE2	5:C:977:HOH:O	2.01	0.60
1:B:727:LEU:HD12	1:B:759:LEU:CD1	2.32	0.60
1:A:571:SER:OG	1:A:699:GLU:OE2	2.18	0.60
1:A:727:LEU:CD2	1:A:759:LEU:CD1	2.80	0.60
1:A:617:SER:O	1:A:621:GLN:HG3	2.01	0.59
1:B:727:LEU:CD1	1:B:759:LEU:HD11	2.33	0.58
1:B:545:HIS:CE1	1:D:760:SER:HB2	2.39	0.58
1:A:493:MET:O	1:A:497:SER:HB2	2.03	0.58
1:C:700:GLY:HA3	1:C:714:MET:O	2.05	0.56
1:B:470:LYS:HE2	1:D:746:PRO:HG3	1.88	0.56
1:B:749:GLU:N	1:B:750:PRO:CD	2.69	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:727:LEU:CD1	1:B:759:LEU:CD1	2.84	0.55
1:D:498:CYS:SG	1:D:554:LYS:HG3	2.47	0.55
1:A:735:ILE:HB	1:A:736:PRO:HD3	1.88	0.55
1:D:713:MET:HG3	2:D:801:IKH:C15	2.35	0.55
1:B:700:GLY:HA3	1:B:714:MET:O	2.07	0.54
1:A:542:GLN:NE2	1:A:542:GLN:HA	2.23	0.54
1:A:492:TYR:CZ	1:A:496:ARG:HD2	2.43	0.53
1:A:719:LYS:O	1:A:722:VAL:HG23	2.08	0.53
1:A:721:GLU:HB2	2:A:801:IKH:C24	2.40	0.52
1:A:498:CYS:SG	1:A:554:LYS:HG3	2.49	0.52
1:A:700:GLY:HA3	1:A:714:MET:O	2.10	0.52
1:D:624:GLU:OE2	1:D:627:ARG:NH2	2.41	0.51
1:D:700:GLY:HA3	1:D:714:MET:O	2.11	0.51
1:A:722:VAL:HB	1:A:723:PRO:HD3	1.94	0.49
1:C:510:ARG:HG2	1:C:608:HIS:CE1	2.48	0.49
1:D:493:MET:SD	1:D:535:HIS:HA	2.54	0.48
1:A:730:TYR:HA	1:A:734:ALA:HB3	1.96	0.48
1:C:677:SER:OG	1:C:688:THR:HG21	2.14	0.47
1:A:507:LYS:NZ	5:A:904:HOH:O	2.48	0.47
1:C:749:GLU:HB3	1:C:750:PRO:HD3	1.96	0.47
1:B:727:LEU:HD12	1:B:759:LEU:HD11	1.95	0.46
1:B:473:GLU:OE2	1:B:496:ARG:NH1	2.49	0.45
1:D:638:TYR:OH	1:D:642:ARG:HD3	2.15	0.45
1:A:749:GLU:N	1:A:750:PRO:CD	2.79	0.45
1:B:504:GLU:HB3	1:B:507:LYS:HD2	1.99	0.45
1:D:725:GLY:HA3	2:D:801:IKH:C18	2.47	0.45
1:B:650:GLN:HA	1:B:650:GLN:OE1	2.17	0.45
1:B:497:SER:O	1:B:553:ARG:HD2	2.17	0.44
1:A:551:LEU:HD23	1:A:551:LEU:HA	1.86	0.44
1:B:740:THR:O	1:B:744:ILE:HG13	2.17	0.44
1:C:651:THR:HG23	5:C:999:HOH:O	2.17	0.44
1:B:627:ARG:O	1:B:631:ILE:HG12	2.18	0.44
1:D:647:GLU:O	1:D:651:THR:HG23	2.18	0.44
1:B:544:ASN:HB3	1:B:547:LEU:HD22	1.99	0.43
1:A:542:GLN:HA	1:A:542:GLN:HE21	1.82	0.43
1:A:727:LEU:HD23	1:A:759:LEU:CD1	2.48	0.43
1:D:693:TYR:OH	2:D:801:IKH:N19	2.52	0.43
1:B:648:MET:HE1	5:B:945:HOH:O	2.18	0.43
1:A:727:LEU:CD2	1:A:759:LEU:HD12	2.48	0.42
1:C:551:LEU:HD23	1:C:551:LEU:HA	1.84	0.42
1:B:611:PHE:HB3	1:B:614:LEU:HD12	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:716:ARG:O	1:B:719:LYS:HB2	2.19	0.42
1:C:727:LEU:HD22	1:C:759:LEU:CD1	2.49	0.42
1:C:722:VAL:N	1:C:723:PRO:CD	2.83	0.42
1:A:658:ASN:O	1:A:661:HIS:HB2	2.20	0.42
1:C:486:TRP:N	1:C:487:PRO:CD	2.83	0.42
1:D:766:ILE:HG22	1:D:766:ILE:O	2.19	0.42
1:B:492:TYR:CZ	1:B:496:ARG:HD3	2.55	0.42
1:A:722:VAL:N	1:A:723:PRO:CD	2.82	0.42
1:D:564:ASP:O	1:D:567:HIS:HB2	2.20	0.42
1:B:646:GLU:O	1:B:650:GLN:HG2	2.20	0.41
1:B:585:LEU:HD12	1:B:585:LEU:HA	1.86	0.41
1:C:740:THR:HA	1:C:743:GLN:HE21	1.86	0.41
1:D:483:GLU:HA	1:D:486:TRP:CE3	2.55	0.41
1:A:727:LEU:HD21	1:A:763:GLU:HG3	2.02	0.41
1:D:635:LEU:O	1:D:638:TYR:HB3	2.21	0.41
1:B:545:HIS:CD2	1:B:545:HIS:H	2.39	0.41
1:C:627:ARG:O	1:C:631:ILE:HG12	2.21	0.40
1:B:557:LEU:HD23	1:B:557:LEU:HA	1.89	0.40
1:A:477:PHE:HB3	1:A:535:HIS:CE1	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	311/343 (91%)	305 (98%)	6 (2%)	0	100	100
1	B	317/343 (92%)	306 (96%)	11 (4%)	0	100	100
1	C	311/343 (91%)	304 (98%)	7 (2%)	0	100	100
1	D	314/343 (92%)	299 (95%)	15 (5%)	0	100	100
All	All	1253/1372 (91%)	1214 (97%)	39 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/305 (92%)	273 (97%)	9 (3%)	34	34
1	B	287/305 (94%)	277 (96%)	10 (4%)	31	30
1	C	282/305 (92%)	277 (98%)	5 (2%)	54	59
1	D	285/305 (93%)	280 (98%)	5 (2%)	54	59
All	All	1136/1220 (93%)	1107 (97%)	29 (3%)	41	43

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	497	SER
1	A	517	LYS
1	A	521	ARG
1	A	542	GLN
1	A	550	ASP
1	A	638	TYR
1	A	653	SER
1	A	677	SER
1	A	727	LEU
1	B	452	THR
1	B	453	SER
1	B	454	GLU
1	B	496	ARG
1	B	587	SER
1	B	617	SER
1	B	660	SER
1	B	677	SER
1	B	749	GLU
1	B	760	SER
1	C	510	ARG
1	C	616	SER
1	C	617	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	638	TYR
1	C	760	SER
1	D	460	MET
1	D	617	SER
1	D	656	LEU
1	D	662	ARG
1	D	760	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	542	GLN
1	A	545	HIS
1	A	604	GLN
1	A	621	GLN
1	A	743	GLN
1	B	542	GLN
1	B	604	GLN
1	B	743	GLN
1	B	761	GLN
1	C	542	GLN
1	C	604	GLN
1	C	743	GLN
1	D	484	ASN
1	D	542	GLN
1	D	604	GLN
1	D	650	GLN
1	D	726	GLN
1	D	743	GLN
1	D	761	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CME	D	509	1	8,9,10	0.62	0	6,9,11	0.69	0
1	CME	B	509	1	8,9,10	0.49	0	6,9,11	0.88	0
1	CME	C	509	1	8,9,10	0.44	0	6,9,11	0.60	0
1	CME	A	509	1	8,9,10	0.70	0	6,9,11	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	D	509	1	-	1/5/8/10	-
1	CME	B	509	1	-	3/5/8/10	-
1	CME	C	509	1	-	3/5/8/10	-
1	CME	A	509	1	-	1/5/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	509	CME	SD-CE-CZ-OH
1	C	509	CME	SD-CE-CZ-OH
1	C	509	CME	CE-SD-SG-CB
1	D	509	CME	SD-CE-CZ-OH
1	A	509	CME	CZ-CE-SD-SG
1	B	509	CME	CE-SD-SG-CB
1	B	509	CME	CZ-CE-SD-SG
1	C	509	CME	CZ-CE-SD-SG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	509	CME	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	IKH	B	801	-	19,29,29	2.27	5 (26%)	21,41,41	1.17	2 (9%)
2	IKH	D	801	-	19,29,29	2.62	9 (47%)	21,41,41	1.17	1 (4%)
2	IKH	A	801	-	19,29,29	1.90	7 (36%)	21,41,41	1.96	6 (28%)
2	IKH	C	801	-	19,29,29	1.77	6 (31%)	21,41,41	1.07	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IKH	B	801	-	-	0/6/11/11	0/4/4/4
2	IKH	D	801	-	-	0/6/11/11	0/4/4/4
2	IKH	A	801	-	-	0/6/11/11	0/4/4/4
2	IKH	C	801	-	-	0/6/11/11	0/4/4/4

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	801	IKH	C05-N09	6.57	1.39	1.33
2	B	801	IKH	C03-N04	5.19	1.44	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	801	IKH	C03-N04	4.47	1.42	1.34
2	B	801	IKH	C05-N09	4.35	1.37	1.33
2	D	801	IKH	C01-C05	3.69	1.46	1.40
2	C	801	IKH	C03-N04	3.64	1.41	1.34
2	B	801	IKH	C01-C05	3.63	1.46	1.40
2	B	801	IKH	C01-C02	3.62	1.44	1.36
2	D	801	IKH	C01-C02	3.52	1.44	1.36
2	A	801	IKH	C03-N04	3.46	1.41	1.34
2	A	801	IKH	C01-C05	2.98	1.45	1.40
2	B	801	IKH	C21-C18	-2.89	1.44	1.49
2	A	801	IKH	C23-C22	2.80	1.43	1.38
2	C	801	IKH	C05-N09	2.78	1.36	1.33
2	A	801	IKH	C01-C02	2.69	1.42	1.36
2	C	801	IKH	C01-C02	2.50	1.42	1.36
2	C	801	IKH	C25-C26	2.48	1.43	1.38
2	C	801	IKH	C01-C05	2.44	1.44	1.40
2	D	801	IKH	C22-C21	2.43	1.44	1.39
2	A	801	IKH	C02-C03	2.39	1.45	1.39
2	A	801	IKH	C25-C26	2.39	1.43	1.38
2	D	801	IKH	C02-C03	2.36	1.45	1.39
2	C	801	IKH	C23-C22	2.33	1.42	1.38
2	D	801	IKH	C25-C26	2.22	1.42	1.38
2	A	801	IKH	C20-N16	-2.17	1.42	1.47
2	D	801	IKH	C03-C13	2.05	1.48	1.44
2	D	801	IKH	C21-C18	-2.01	1.45	1.49

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	IKH	C01-C02-C03	4.97	123.98	118.80
2	A	801	IKH	C02-C01-C05	-3.67	115.31	119.76
2	A	801	IKH	C20-N16-C15	-2.97	122.19	126.39
2	A	801	IKH	C20-N16-C17	2.57	129.72	124.90
2	B	801	IKH	C26-C21-C22	2.46	122.08	117.68
2	B	801	IKH	C22-C21-C18	-2.31	117.63	121.28
2	C	801	IKH	C17-N16-C15	-2.27	106.14	108.60
2	A	801	IKH	C02-C03-N04	-2.26	120.01	123.07
2	D	801	IKH	C02-C03-N04	-2.15	120.17	123.07
2	A	801	IKH	C18-C17-N16	-2.03	105.78	107.91

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	801	IKH	3	0
2	A	801	IKH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	312/343 (90%)	-0.15	1 (0%) 90 92	24, 46, 73, 104	1 (0%)
1	B	319/343 (93%)	-0.18	3 (0%) 81 83	34, 45, 78, 124	0
1	C	312/343 (90%)	-0.26	3 (0%) 79 82	26, 44, 69, 104	1 (0%)
1	D	316/343 (92%)	0.31	4 (1%) 74 78	47, 64, 90, 137	0
All	All	1259/1372 (91%)	-0.07	11 (0%) 81 83	24, 50, 83, 137	2 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	456	TRP	3.7
1	A	459	LEU	3.5
1	B	771	THR	3.5
1	D	452	THR	3.3
1	B	456	TRP	3.1
1	D	459	LEU	2.8
1	C	459	LEU	2.7
1	C	460	MET	2.5
1	B	454	GLU	2.3
1	C	458	GLY	2.2
1	D	727	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	D	509	10/11	0.81	0.15	63,79,103,112	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	CME	B	509	10/11	0.90	0.12	47,64,84,94	0
1	CME	C	509	10/11	0.92	0.11	44,52,84,86	0
1	CME	A	509	10/11	0.93	0.10	48,60,89,93	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	IKH	D	801	26/26	0.94	0.09	49,58,65,67	0
2	IKH	B	801	26/26	0.96	0.07	37,41,45,50	0
2	IKH	C	801	26/26	0.96	0.07	35,45,49,54	0
2	IKH	A	801	26/26	0.96	0.07	35,41,48,51	0
4	MG	B	803	1/1	0.99	0.02	32,32,32,32	0
4	MG	C	803	1/1	0.99	0.02	33,33,33,33	0
4	MG	D	803	1/1	0.99	0.04	47,47,47,47	0
3	ZN	D	802	1/1	1.00	0.03	54,54,54,54	0
4	MG	A	803	1/1	1.00	0.02	33,33,33,33	0
3	ZN	A	802	1/1	1.00	0.02	41,41,41,41	0
3	ZN	B	802	1/1	1.00	0.01	38,38,38,38	0
3	ZN	C	802	1/1	1.00	0.01	38,38,38,38	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around IKH D 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around IKH B 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around IKH C 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around IKH A 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.