

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 8, 2024 – 08:26 AM EDT

PDB ID	:	5SJP
Title	:	CRYSTAL STRUCTURE OF HUMAN PHOSPHODIESTERASE 10 IN
		COMPLEX WITH $N1(CCC1)C(=O)c2c(n(nc2)C)C(=O)Nc3cc(nc(c3)C)$
		)NC(NC(OCC)=O)=S, micromolar IC50=0.402697
Authors	:	Joseph, C.; Benz, J.; Flohr, A.; Lerner, C.; Rudolph, M.G.
Deposited on	:	2022-02-01
Resolution	:	2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motric	Whole archive	Similar resolution		
IVIETIC	$(\# { m Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$		
$R_{free}$	164625	1763 (2.26-2.26)		
Clashscore	180529	1919 (2.26-2.26)		
Ramachandran outliers	177936	1884 (2.26-2.26)		
Sidechain outliers	177891	1885 (2.26-2.26)		
RSRZ outliers	164620	1763 (2.26-2.26)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	А	343	% • 77%	12%	•	9%
1	В	343	% • 79%	11%	•	8%
1	С	343	% • 80%	9%	•	9%
1	D	343	% 81%	9%		10%



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10771 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	212	Total	С	Ν	Ο	$\mathbf{S}$	0	1	0
1	Л	515	2549	1629	435	461	24	0		0
1	В	215	Total	С	Ν	Ο	S	0	1	0
1	I D	313	2559	1635	437	463	24	0		
1	С	313	Total	С	Ν	0	S	0	2	0
			2557	1634	438	461	24	0	2	U
1 D	210	Total	С	Ν	0	S	0	0	0	
	I D	310	2519	1612	429	454	24	U	U	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	447	GLY	-	expression tag	UNP Q9Y233
А	448	SER	-	expression tag	UNP Q9Y233
В	447	GLY	-	expression tag	UNP Q9Y233
В	448	SER	-	expression tag	UNP Q9Y233
С	447	GLY	-	expression tag	UNP Q9Y233
С	448	SER	-	expression tag	UNP Q9Y233
D	447	GLY	-	expression tag	UNP Q9Y233
D	448	SER	-	expression tag	UNP Q9Y233

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Zn 1 1	0	0
2	В	1	Total Zn 1 1	0	0
2	С	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0
3	С	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

• Molecule 4 is ethyl [(4-{[4-(azetidine-1-carbonyl)-1-methyl-1H-pyrazole-5-carbonyl]am ino}-6-methylpyridin-2-yl)carbamothioyl]carbamate (three-letter code: K6O) (formula:  $C_{19}H_{23}N_7O_4S$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	Δ	1	Total	С	Ν	0	$\mathbf{S}$	0	0
4	Π	T	31	19	7	4	1	0	0
4	В	1	Total	С	Ν	0	S	0	0
4	D	I	31	19	7	4	1	0	0
4	С	Y 1	Total	С	Ν	Ο	S	0	0
4	U	L	31	19	7	4	1	0	0
4	Л	1	Total	С	Ν	0	S	0	0
4	D	L	31	19	7	4	1	0	0

• Molecule 5 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	124	Total O 124 124	0	0
5	В	131	Total O 131 131	0	0
5	С	137	Total O 137 137	0	0
5	D	63	$\begin{array}{cc} \text{Total} & \text{O} \\ 63 & 63 \end{array}$	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



# 



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	Н 3	Depositor
Cell constants	135.30Å 135.30Å 234.70Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
<b>D</b> ecolution $(\hat{\lambda})$	43.52 - 2.25	Depositor
Resolution (A)	43.52 - 2.25	EDS
% Data completeness	95.3 (43.52-2.25)	Depositor
(in resolution range)	95.3(43.52-2.25)	EDS
R <sub>merge</sub>	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.40 (at 2.24 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
P. P.	0.173 , $0.234$	Depositor
$n, n_{free}$	0.180 , $0.236$	DCC
$R_{free}$ test set	3788 reflections $(4.97%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	40.7	Xtriage
Anisotropy	0.184	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33, $32.3$	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49, < L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	10771	wwPDB-VP
Average B, all atoms $(Å^2)$	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.73% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CME, MG, K6O

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Unain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.81	0/2603	0.91	2/3521~(0.1%)	
1	В	0.78	0/2613	0.91	1/3535~(0.0%)	
1	С	0.79	0/2614	0.94	2/3535~(0.1%)	
1	D	0.83	0/2570	0.89	0/3478	
All	All	0.80	0/10400	0.91	5/14069~(0.0%)	

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	С	467	ARG	NE-CZ-NH1	8.16	124.38	120.30
1	В	642	ARG	NE-CZ-NH2	-6.94	116.83	120.30
1	С	642	ARG	NE-CZ-NH2	-5.76	117.42	120.30
1	А	642	ARG	NE-CZ-NH2	-5.56	117.52	120.30
1	А	553	ARG	CB-CA-C	5.55	121.50	110.40

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2549	0	2524	32	0
1	В	2559	0	2528	25	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	С	2557	0	2537	17	0
1	D	2519	0	2496	17	0
2	А	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
3	А	1	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
4	А	31	0	0	1	0
4	В	31	0	0	1	0
4	С	31	0	0	2	0
4	D	31	0	0	0	0
5	А	124	0	0	3	0
5	В	131	0	0	4	0
5	С	137	0	0	3	0
5	D	63	0	0	2	0
All	All	10771	0	10085	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:C:469:CYS:SG	5:C:1022:HOH:O	2.15	1.04
1:B:509:CME:HB2	5:B:1008:HOH:O	1.69	0.91
1:D:469:CYS:SG	5:D:961:HOH:O	2.32	0.87
1:B:470:LYS:HE2	1:D:746:PRO:HG3	1.61	0.83
1:A:549:THR:OG1	1:A:552:GLU:HG3	1.85	0.77
1:D:766:ILE:HG22	1:D:766:ILE:O	1.87	0.74
1:B:576:GLN:HB2	5:B:1010:HOH:O	1.88	0.74
1:C:509:CME:HB2	5:C:1002:HOH:O	1.87	0.72
1:A:461:GLN:HE22	1:A:500:THR:HG21	1.59	0.68
1:A:461:GLN:NE2	1:A:461:GLN:HA	2.07	0.68
1:A:722:VAL:HB	1:A:723:PRO:HD3	1.77	0.64
1:A:493:MET:O	1:A:497:SER:HB2	1.98	0.64
1:D:469:CYS:SG	5:D:951:HOH:O	2.56	0.61
1:A:492:TYR:CZ	1:A:496:ARG:HD2	2.36	0.60
1:C:568[A]:ARG:NH1	5:C:904:HOH:O	2.36	0.59



Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:735:ILE:HB	1:A:736:PRO:HD3	1.85	0.58
1:D:730:TYR:HA	1:D:734:ALA:HB3	1.83	0.58
1:D:493:MET:O	1:D:497:SER:HB3	2.03	0.58
4:C:803:K6O:O3	4:C:803:K6O:N11	2.37	0.57
1:A:469:CYS:SG	5:A:1008:HOH:O	2.16	0.56
1:D:766:ILE:O	1:D:766:ILE:CG2	2.52	0.55
1:B:497:SER:O	1:B:553:ARG:HD2	2.07	0.55
1:C:611:PHE:HB3	1:C:614:LEU:HD22	1.89	0.55
1:C:727:LEU:HD21	1:C:763:GLU:HG3	1.89	0.55
1:B:493:MET:O	1:B:497:SER:HB2	2.08	0.53
1:A:461:GLN:NE2	1:A:461:GLN:CA	2.71	0.53
1:B:700:GLY:HA3	1:B:714:MET:O	2.09	0.53
1:B:727:LEU:CD2	1:B:759:LEU:HD11	2.39	0.52
1:D:492:TYR:CZ	1:D:496:ARG:HD2	2.44	0.52
1:A:497:SER:O	1:A:553:ARG:HD2	2.09	0.52
1:C:749:GLU:HB3	1:C:750:PRO:HD3	1.92	0.52
1:B:727:LEU:HD23	1:B:759:LEU:CD1	2.40	0.51
1:A:749:GLU:N	1:A:750:PRO:CD	2.74	0.51
1:D:627:ARG:O	1:D:631:ILE:HG12	2.11	0.51
1:D:740:THR:HA	1:D:743:GLN:HE21	1.75	0.51
1:A:648:MET:HE1	5:A:1010:HOH:O	2.10	0.50
1:A:638:TYR:OH	1:A:642:ARG:HD3	2.12	0.50
1:A:602:ILE:HA	1:A:605:LEU:HD22	1.94	0.49
1:A:719:LYS:O	1:A:722:VAL:HG23	2.12	0.49
1:A:507:LYS:NZ	5:A:905:HOH:O	2.45	0.48
1:A:711:ILE:CD1	1:A:713:MET:HE3	2.42	0.48
1:C:711:ILE:HD11	1:C:713:MET:CE	2.43	0.48
1:D:493:MET:SD	1:D:535:HIS:HA	2.54	0.48
1:C:486:TRP:N	1:C:487:PRO:CD	2.77	0.48
1:A:461:GLN:CA	1:A:461:GLN:HE21	2.28	0.47
1:B:724:GLN:NE2	5:B:905:HOH:O	2.39	0.47
1:B:548:PHE:O	1:B:553:ARG:NH1	2.48	0.46
1:D:564:ASP:O	1:D:567:HIS:HB2	2.15	0.46
1:A:477:PHE:HB3	1:A:535:HIS:CE1	2.51	0.46
1:B:546:THR:HG21	1:D:767:ARG:NH2	2.31	0.46
1:B:752:LEU:HD21	1:B:756[B]:ARG:NH2	2.31	0.46
1:A:730:TYR:HA	1:A:734:ALA:HB3	1.98	0.46
4:C:803:K6O:O16	4:C:803:K6O:C21	2.63	0.46
1:A:542:GLN:NE2	1:A:542:GLN:HA	2.31	0.46
1:B:611:PHE:HB3	1:B:614:LEU:HD22	1.98	0.46
1:B:638:TYR:CD1	1:B:671:THR:HG21	2.51	0.46



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:B:627:ARG:O	1:B:631:ILE:HG12	2.17	0.45
1:C:727:LEU:HG	1:C:766:ILE:CD1	2.47	0.45
1:A:461:GLN:HA	1:A:461:GLN:HE21	1.80	0.45
1:B:769:GLU:O	1:B:769:GLU:HG2	2.16	0.45
1:C:707:GLY:O	1:D:652:GLY:HA3	2.15	0.45
1:C:700:GLY:HA3	1:C:714:MET:O	2.17	0.45
1:A:461:GLN:HE22	1:A:500:THR:CG2	2.29	0.45
1:A:486:TRP:N	1:A:487:PRO:CD	2.80	0.44
1:B:767:ARG:NH2	1:B:769:GLU:OE1	2.51	0.44
1:A:644:GLN:HE21	1:A:644:GLN:HA	1.83	0.44
1:A:662:ARG:NH1	1:A:662:ARG:HG2	2.32	0.44
1:B:697:TRP:CZ2	1:B:719:LYS:HG2	2.53	0.43
1:B:492:TYR:CZ	1:B:496:ARG:HD2	2.54	0.43
1:B:646:GLU:HG2	5:B:932:HOH:O	2.18	0.43
4:B:803:K6O:O16	4:B:803:K6O:C21	2.64	0.43
1:C:638:TYR:CD1	1:C:671:THR:HG21	2.53	0.43
1:C:467:ARG:O	1:C:471:GLU:HB2	2.20	0.42
1:B:498:CYS:HB3	1:B:553:ARG:HB3	2.00	0.42
1:A:735:ILE:N	1:A:736:PRO:CD	2.82	0.42
1:B:646:GLU:HA	1:B:646:GLU:OE1	2.20	0.42
1:D:605:LEU:HB3	1:D:608:HIS:ND1	2.35	0.42
1:D:535:HIS:O	1:D:538:TYR:HB3	2.20	0.41
1:B:727:LEU:HD23	1:B:727:LEU:HA	1.81	0.41
1:A:716:ARG:O	1:A:719:LYS:HG3	2.20	0.41
1:C:682:TRP:HB3	1:C:683:PRO:HD3	2.03	0.41
1:D:622:VAL:HA	1:D:625:ILE:HD12	2.02	0.41
1:C:697:TRP:CH2	1:C:719:LYS:HD3	2.56	0.41
1:C:746:PRO:N	1:C:747:PRO:CD	2.84	0.41
1:A:489:ILE:O	1:A:493:MET:HG3	2.21	0.41
1:C:557:LEU:HD23	1:C:557:LEU:HA	1.91	0.40
4:A:803:K6O:O29	4:A:803:K6O:N23	2.54	0.40
1:B:493:MET:SD	1:B:535:HIS:HA	2.62	0.40
1:A:662:ARG:CG	1:A:662:ARG:HH11	2.34	0.40
1:A:746:PRO:HB2	1:A:747:PRO:HD3	2.04	0.40
1:B:467:ARG:O	1:B:471:GLU:HB2	2.22	0.40
1:A:662:ARG:NH1	1:A:662:ARG:CG	2.81	0.40

There are no symmetry-related clashes.



#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	311/343~(91%)	298~(96%)	12 (4%)	1 (0%)	37	41
1	В	313/343~(91%)	300~(96%)	13~(4%)	0	100	100
1	С	312/343~(91%)	302~(97%)	10 (3%)	0	100	100
1	D	307/343~(90%)	291 (95%)	16 (5%)	0	100	100
All	All	1243/1372 (91%)	1191 (96%)	51 (4%)	1 (0%)	48	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	768	GLY

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	282/305~(92%)	268~(95%)	14~(5%)	20	22
1	В	282/305~(92%)	265~(94%)	17~(6%)	16	15
1	С	283/305~(93%)	268~(95%)	15~(5%)	19	19
1	D	279/305~(92%)	268~(96%)	11 (4%)	27	33
All	All	1126/1220~(92%)	1069 (95%)	57 (5%)	20	21

All (57) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	А	461	GLN
1	А	471	GLU
1	А	497	SER
1	А	501	SER
1	А	504	GLU
1	А	568	ARG
1	А	575	LEU
1	А	587	SER
1	А	605	LEU
1	А	606	GLU
1	А	622	VAL
1	А	644	GLN
1	А	677	SER
1	А	713	MET
1	В	459	LEU
1	В	470	LYS
1	В	471	GLU
1	В	497	SER
1	В	504	GLU
1	В	575	LEU
1	В	576	GLN
1	В	605	LEU
1	В	617	SER
1	В	638	TYR
1	В	650	GLN
1	В	660	SER
1	В	662	ARG
1	В	677	SER
1	В	713	MET
1	В	727	LEU
1	В	760	SER
1	С	463	THR
1	С	471	GLU
1	С	517	LYS
1	С	547	LEU
1	С	568[A]	ARG
1	С	568[B]	ARG
1	С	576	GLN
1	С	605	LEU
1	С	616	SER
1	С	617	SER
1	С	638	TYR
1	С	711	ILE



Mol	Chain	Res	Type
1	С	713	MET
1	С	727	LEU
1	С	760	SER
1	D	500	THR
1	D	504	GLU
1	D	542	GLN
1	D	575	LEU
1	D	605	LEU
1	D	617	SER
1	D	623	LEU
1	D	644	GLN
1	D	651	THR
1	D	660	SER
1	D	709	GLN

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such side chains are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	А	461	GLN
1	А	484	ASN
1	А	542	GLN
1	А	604	GLN
1	А	644	GLN
1	А	724	GLN
1	А	743	GLN
1	В	484	ASN
1	В	495	HIS
1	В	542	GLN
1	В	604	GLN
1	С	542	GLN
1	С	604	GLN
1	С	743	GLN
1	D	542	GLN
1	D	604	GLN
1	D	621	GLN
1	D	644	GLN
1	D	650	GLN
1	D	743	GLN



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Chain	Dog	Tiple	B	Bond lengths			Bond angles		
IVIOI	туре	Unain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
1	CME	А	509	1	8,9,10	0.45	0	6,9,11	1.17	1 (16%)
1	CME	В	509	1	8,9,10	0.44	0	6,9,11	0.82	0
1	CME	D	509	1	8,9,10	0.36	0	6,9,11	0.86	0
1	CME	С	509	1	8,9,10	0.36	0	6,9,11	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	А	509	1	-	1/5/8/10	-
1	CME	В	509	1	-	2/5/8/10	-
1	CME	D	509	1	-	1/5/8/10	-
1	CME	С	509	1	-	3/5/8/10	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	509	CME	CZ-CE-SD	-2.05	106.53	113.39

There are no chirality outliers.

All (7) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
1	В	509	CME	SD-CE-CZ-OH
1	С	509	CME	CE-SD-SG-CB
1	С	509	CME	SD-CE-CZ-OH
1	А	509	CME	CZ-CE-SD-SG
1	С	509	CME	CZ-CE-SD-SG
1	В	509	CME	CZ-CE-SD-SG
1	D	509	CME	CZ-CE-SD-SG

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	В	509	CME	1	0
1	С	509	CME	1	0

#### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Iol Type Chain Re		Dec		В	Bond lengths			Bond angles		
INIOI	туре	Chain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
4	K6O	D	803	-	30,33,33	2.78	16 (53%)	38,46,46	2.81	16 (42%)	
4	K6O	В	803	-	30,33,33	2.82	16 (53%)	38,46,46	<mark>3.37</mark>	18 (47%)	
4	K6O	С	803	-	30,33,33	2.90	14 (46%)	38,46,46	<mark>3.90</mark>	18 (47%)	
4	K6O	А	803	-	30,33,33	<mark>3.00</mark>	18 (60%)	38,46,46	<mark>3.66</mark>	15 (39%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol

4

4

4

4

Type

K6O

K6O

K6O

K6O

-

-

8/20/33/33

10/20/33/33

0/3/3/3

0/3/3/3

All	(64)	$\operatorname{bond}$	length	outliers	are	listed	below:	

803

803

-

-

С

А

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	А	803	K6O	C24-N23	7.15	1.47	1.35
4	D	803	K6O	C24-N23	6.45	1.46	1.35
4	С	803	K6O	C24-N25	6.40	1.49	1.39
4	С	803	K6O	C24-N23	6.29	1.46	1.35
4	А	803	K6O	C24-N25	6.27	1.49	1.39
4	В	803	K6O	C24-N23	5.92	1.45	1.35
4	С	803	K6O	C27-N25	5.85	1.47	1.37
4	В	803	K6O	C27-N25	5.73	1.47	1.37
4	D	803	K6O	C24-N25	5.61	1.48	1.39
4	В	803	K6O	C24-N25	5.59	1.48	1.39
4	А	803	K6O	C27-N25	5.32	1.46	1.37
4	С	803	K6O	C2-N1	5.18	1.45	1.34
4	D	803	K6O	C27-N25	5.02	1.46	1.37
4	В	803	K6O	C2-N1	4.43	1.44	1.34
4	А	803	K6O	C2-N1	4.00	1.43	1.34
4	В	803	K6O	O28-C27	3.99	1.43	1.35
4	А	803	K6O	C21-C22	3.94	1.46	1.38
4	D	803	K6O	C2-N1	3.92	1.43	1.34
4	С	803	K6O	O28-C27	3.92	1.43	1.35
4	А	803	K6O	C14-N23	3.76	1.49	1.40
4	А	803	K6O	O28-C27	3.75	1.42	1.35
4	В	803	K6O	C4-C2	3.48	1.55	1.50
4	В	803	K6O	C22-N15	3.48	1.41	1.34
4	D	803	K6O	C14-N23	3.47	1.48	1.40
4	D	803	K6O	C21-C22	3.41	1.45	1.38
4	В	803	K6O	C21-C22	3.31	1.45	1.38
4	D	803	K6O	C4-C2	3.27	1.55	1.50
4	А	803	K6O	C19-C18	-3.22	1.46	1.54
4	D	803	K6O	C19-C20	-3.17	1.47	1.54
4	С	803	K6O	C21-C22	3.15	1.44	1.38
4	А	803	K6O	C19-C20	-3.11	1.47	1.54
4	С	803	K6O	C19-C20	-3.07	1.47	1.54
4	А	803	K6O	C22-N15	3.07	1.40	1.34
4	В	803	K6O	C14-N23	3.01	1.47	1.40



Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
4	С	803	K6O	C19-C18	-2.99	1.47	1.54
4	С	803	K6O	C9-N7	2.97	1.53	1.47
4	D	803	K6O	O28-C27	2.91	1.41	1.35
4	В	803	K6O	C19-C18	-2.91	1.47	1.54
4	С	803	K6O	C22-N15	2.91	1.40	1.34
4	С	803	K6O	C14-N23	2.90	1.47	1.40
4	D	803	K6O	C19-C18	-2.86	1.47	1.54
4	С	803	K6O	C10-N11	2.85	1.44	1.35
4	В	803	K6O	C13-C12	2.76	1.44	1.39
4	А	803	K6O	C14-N15	2.69	1.39	1.34
4	D	803	K6O	C10-N11	2.65	1.43	1.35
4	В	803	K6O	C13-C14	2.58	1.46	1.39
4	А	803	K6O	C20-N1	-2.54	1.45	1.47
4	В	803	K6O	C9-N7	2.49	1.52	1.47
4	В	803	K6O	C19-C20	-2.49	1.48	1.54
4	А	803	K6O	C13-C12	2.48	1.43	1.39
4	А	803	K6O	C10-N11	2.47	1.42	1.35
4	D	803	K6O	C22-N15	2.47	1.39	1.34
4	А	803	K6O	C12-N11	2.47	1.46	1.41
4	D	803	K6O	C13-C14	2.45	1.45	1.39
4	С	803	K6O	C14-N15	2.43	1.39	1.34
4	А	803	K6O	C13-C14	2.36	1.45	1.39
4	В	803	K6O	C14-N15	2.31	1.39	1.34
4	D	803	K6O	C12-N11	2.22	1.46	1.41
4	В	803	K6O	C5-N6	2.16	1.39	1.33
4	А	803	K6O	C4-C2	2.14	1.53	1.50
4	А	803	K6O	C4-C8	2.13	1.44	1.40
4	С	803	K6O	C13-C14	2.12	1.45	1.39
4	D	803	K6O	C17-C22	2.08	1.55	1.50
4	D	803	K6O	C8-N7	2.04	1.40	1.36

Continued from previous page...

All (67) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	С	803	K6O	C24-N25-C27	-13.26	117.38	127.40
4	А	803	K6O	C24-N25-C27	-11.65	118.60	127.40
4	С	803	K6O	O28-C27-N25	11.35	125.78	108.83
4	В	803	K6O	O28-C27-N25	9.51	123.03	108.83
4	А	803	K6O	O28-C27-N25	9.38	122.84	108.83
4	В	803	K6O	O3-C2-N1	-7.81	110.04	122.35
4	D	803	K6O	O28-C27-N25	7.14	119.50	108.83
4	А	803	K6O	O3-C2-N1	-6.98	111.35	122.35



Mol	Chain	Res	Type	Atoms	Ζ	Observed(°)	$Ideal(^{o})$
4	С	803	K6O	O29-C27-N25	-6.97	115.85	125.47
4	В	803	K6O	C24-N25-C27	-6.68	122.35	127.40
4	А	803	K6O	O29-C27-N25	-6.63	116.31	125.47
4	D	803	K6O	O3-C2-N1	-6.22	112.55	122.35
4	D	803	K6O	C24-N25-C27	-6.16	122.74	127.40
4	С	803	K6O	C30-O28-C27	6.13	127.52	116.06
4	В	803	K6O	C8-N7-N6	-6.08	106.94	112.85
4	В	803	K6O	O29-C27-N25	-5.96	117.23	125.47
4	D	803	K6O	C8-N7-N6	-5.74	107.27	112.85
4	А	803	K6O	C5-C4-C8	5.29	108.74	103.81
4	С	803	K6O	C8-N7-N6	-5.24	107.76	112.85
4	С	803	K6O	C9-N7-C8	-5.04	124.22	129.47
4	В	803	K6O	C5-N6-N7	5.01	109.45	104.23
4	А	803	K6O	C5-N6-N7	4.93	109.36	104.23
4	С	803	K6O	C17-C22-N15	4.65	123.59	116.56
4	D	803	K6O	O29-C27-N25	-4.54	119.20	125.47
4	D	803	K6O	O3-C2-C4	4.38	128.66	120.06
4	D	803	K6O	C5-N6-N7	4.35	108.76	104.23
4	D	803	K6O	C14-N15-C22	4.28	124.09	118.15
4	А	803	K6O	C9-N7-C8	-4.23	125.06	129.47
4	А	803	K6O	C4-C2-N1	4.19	124.75	118.24
4	С	803	K6O	O3-C2-N1	-4.12	115.85	122.35
4	В	803	K6O	C4-C2-N1	4.06	124.54	118.24
4	А	803	K6O	C14-N15-C22	3.99	123.69	118.15
4	В	803	K6O	C14-N15-C22	3.82	123.45	118.15
4	В	803	K6O	C21-C12-C13	3.74	124.67	119.59
4	А	803	K6O	C8-N7-N6	-3.71	109.24	112.85
4	С	803	K6O	C14-N15-C22	3.57	123.10	118.15
4	А	803	K6O	C30-O28-C27	3.36	122.35	116.06
4	В	803	K6O	C14-N23-C24	-3.33	127.16	130.74
4	С	803	K6O	C14-N23-C24	-3.08	127.42	130.74
4	С	803	K6O	O28-C27-O29	-3.08	118.35	124.26
4	В	803	K6O	C13-C14-N15	-3.00	117.13	123.35
4	А	803	K6O	O16-C10-N11	-2.99	116.15	123.75
4	С	803	K6O	C5-C4-C8	2.95	106.56	103.81
4	В	803	K6O	N23-C24-N25	2.83	119.55	115.31
4	В	803	K6O	C30-O28-C27	2.76	121.23	116.06
4	В	803	K6O	O3-C2-C4	2.73	125.42	120.06
4	А	803	K6O	C13-C14-N15	-2.70	117.75	123.35
4	D	803	K6O	C20-C19-C18	2.57	91.84	88.69
4	С	803	K6O	C5-N6-N7	2.54	106.88	104.23
4	С	803	K6O	C21-C12-C13	2.53	123.02	119.59



Mol	Chain	$\mathbf{Res}$	Type	Atoms	$\mathbf{Z}$	$Observed(^{o})$	$Ideal(^{o})$
4	В	803	K6O	C5-C4-C8	2.51	106.15	103.81
4	С	803	K6O	C21-C22-N15	-2.50	118.02	121.50
4	D	803	K6O	C30-O28-C27	2.49	120.72	116.06
4	А	803	K6O	C19-C18-N1	-2.47	86.66	88.21
4	D	803	K6O	C13-C14-N15	-2.45	118.26	123.35
4	А	803	K6O	C5-C4-C2	-2.39	122.10	128.56
4	С	803	K6O	N23-C14-N15	2.39	119.11	113.33
4	В	803	K6O	O28-C27-O29	-2.36	119.73	124.26
4	С	803	K6O	C17-C22-C21	-2.31	118.38	121.80
4	С	803	K6O	C13-C14-N15	-2.18	118.82	123.35
4	D	803	K6O	C5-C4-C8	2.15	105.82	103.81
4	D	803	K6O	C17-C22-N15	2.12	119.76	116.56
4	D	803	K6O	C8-C4-C2	2.12	131.10	127.04
4	D	803	K6O	C21-C22-N15	-2.09	118.59	121.50
4	B	803	K6O	C5-C4-C2	-2.05	123.02	128.56
4	D	803	K6O	C5-C4-C2	-2.03	123.06	128.56
4	В	803	K6O	S26-C24-N23	-2.03	118.44	124.33

There are no chirality outliers.

All (32) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	803	K6O	N23-C24-N25-C27
4	А	803	K6O	S26-C24-N25-C27
4	А	803	K6O	N25-C27-O28-C30
4	А	803	K6O	O29-C27-O28-C30
4	А	803	K6O	O3-C2-C4-C5
4	В	803	K6O	N25-C24-N23-C14
4	В	803	K6O	S26-C24-N23-C14
4	В	803	K6O	N25-C27-O28-C30
4	В	803	K6O	O29-C27-O28-C30
4	В	803	K6O	O3-C2-C4-C5
4	С	803	K6O	N25-C24-N23-C14
4	С	803	K6O	S26-C24-N23-C14
4	С	803	K6O	N23-C24-N25-C27
4	С	803	K6O	S26-C24-N25-C27
4	С	803	K6O	N25-C27-O28-C30
4	С	803	K6O	O29-C27-O28-C30
4	С	803	K6O	O3-C2-C4-C5
4	D	803	K6O	O28-C27-N25-C24
4	D	803	K6O	O29-C27-N25-C24
4	D	803	K6O	N25-C27-O28-C30



Mol	Chain	Res	Type	Atoms
4	D	803	K6O	O29-C27-O28-C30
4	D	803	K6O	O3-C2-C4-C5
4	А	803	K6O	O28-C27-N25-C24
4	А	803	K6O	O29-C27-N25-C24
4	D	803	K6O	C13-C14-N23-C24
4	D	803	K6O	N15-C14-N23-C24
4	В	803	K6O	O29-C27-N25-C24
4	А	803	K6O	C13-C14-N23-C24
4	А	803	K6O	N15-C14-N23-C24
4	С	803	K6O	O28-C27-N25-C24
4	А	803	K6O	C31-C30-O28-C27
4	D	803	K6O	C31-C30-O28-C27

Continued from previous page...

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	803	K6O	1	0
4	С	803	K6O	2	0
4	А	803	K6O	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	312/343~(90%)	-0.33	4 (1%) 74 76	24, 44, 73, 107	1 (0%)
1	В	314/343~(91%)	-0.31	4 (1%) 74 76	26, 44, 74, 105	1 (0%)
1	С	312/343~(90%)	-0.38	3 (0%) 79 80	24, 43, 70, 103	2 (0%)
1	D	309/343~(90%)	0.05	3 (0%) 79 80	44, 60, 83, 109	0
All	All	1247/1372~(90%)	-0.24	14 (1%) 77 79	24, 49, 79, 109	4 (0%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	459	LEU	5.4
1	С	459	LEU	4.2
1	В	771	THR	3.8
1	В	458	GLY	3.5
1	В	457	GLN	3.3
1	А	459	LEU	3.0
1	А	458	GLY	2.9
1	С	458	GLY	2.7
1	В	459	LEU	2.5
1	А	770	GLU	2.4
1	С	460	MET	2.3
1	D	768	GLY	2.1
1	А	643	LYS	2.0
1	D	460	MET	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B} ext{-factors}({ m \AA}^2)$	Q < 0.9
1	CME	D	509	10/11	0.87	0.13	54,78,101,103	0
1	CME	В	509	10/11	0.88	0.14	44,60,99,99	0
1	CME	С	509	10/11	0.91	0.12	43,51,88,92	0
1	CME	А	509	10/11	0.94	0.09	44,61,84,90	0

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
4	K6O	D	803	31/31	0.80	0.16	62,83,104,108	0
4	K6O	С	803	31/31	0.82	0.15	45,65,102,105	0
4	K6O	А	803	31/31	0.86	0.16	44,68,103,104	0
4	K6O	В	803	31/31	0.87	0.13	48,62,85,88	0
3	MG	С	802	1/1	0.99	0.02	$31,\!31,\!31,\!31$	0
3	MG	D	802	1/1	0.99	0.03	44,44,44,44	0
2	ZN	А	801	1/1	1.00	0.01	$38,\!38,\!38,\!38$	0
2	ZN	В	801	1/1	1.00	0.01	$38,\!38,\!38,\!38$	0
2	ZN	С	801	1/1	1.00	0.01	40,40,40,40	0
2	ZN	D	801	1/1	1.00	0.02	52,52,52,52	0
3	MG	А	802	1/1	1.00	0.02	31,31,31,31	0
3	MG	В	802	1/1	1.00	0.01	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.













## 6.5 Other polymers (i)

There are no such residues in this entry.

