



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2024 – 09:30 AM EDT

PDB ID : 5T9Y  
Title : Crystal structure of the infectious salmon anemia virus (ISAV) hemagglutinin-esterase protein  
Authors : Cook, J.D.; Sultana, A.; Lee, J.E.  
Deposited on : 2016-09-09  
Resolution : 1.80 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	 2% 89% 5% 6%
1	B	342	 2% 88% 5% 7%
1	C	342	 2% 87% 6% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FMT	B	407	-	-	X	-
4	FMT	B	408	-	-	X	-

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 15896 atoms, of which 7247 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HE protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	321	4867	1539	2406	414	490	18	0	6	0
1	B	319	4796	1523	2365	410	479	19	0	7	0
1	C	319	4873	1538	2413	419	484	19	0	6	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	333	ASP	ASN	conflict	UNP Q9J0Y0
A	354	ARG	-	expression tag	UNP Q9J0Y0
A	355	LEU	-	expression tag	UNP Q9J0Y0
A	356	VAL	-	expression tag	UNP Q9J0Y0
A	357	PRO	-	expression tag	UNP Q9J0Y0
A	358	ARG	-	expression tag	UNP Q9J0Y0
B	333	ASP	ASN	conflict	UNP Q9J0Y0
B	354	ARG	-	expression tag	UNP Q9J0Y0
B	355	LEU	-	expression tag	UNP Q9J0Y0
B	356	VAL	-	expression tag	UNP Q9J0Y0
B	357	PRO	-	expression tag	UNP Q9J0Y0
B	358	ARG	-	expression tag	UNP Q9J0Y0
C	333	ASP	ASN	conflict	UNP Q9J0Y0
C	354	ARG	-	expression tag	UNP Q9J0Y0
C	355	LEU	-	expression tag	UNP Q9J0Y0
C	356	VAL	-	expression tag	UNP Q9J0Y0
C	357	PRO	-	expression tag	UNP Q9J0Y0
C	358	ARG	-	expression tag	UNP Q9J0Y0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Mg 2 2	0	0
2	B	1	Total Mg 1 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C H O 14 3 8 3	0	0
3	B	1	Total C H O 14 3 8 3	0	0
3	B	1	Total C H O 14 3 8 3	0	0

- Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula: CH<sub>2</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			4	1	1	2		
4	A	1	Total	C	H	O	0	0
			4	1	1	2		
4	A	1	Total	C	H	O	0	0
			4	1	1	2		
4	A	1	Total	C	H	O	0	0
			4	1	1	2		
4	B	1	Total	C	H	O	0	0
			4	1	1	2		
4	B	1	Total	C	H	O	0	0
			4	1	1	2		
4	B	1	Total	C	H	O	0	0
			4	1	1	2		
4	B	1	Total	C	H	O	0	0
			4	1	1	2		
4	B	1	Total	C	H	O	0	0
			4	1	1	2		
4	B	1	Total	C	H	O	0	0
			4	1	1	2		
4	B	1	Total	C	H	O	0	0
			4	1	1	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	C	1	Total 4	C 1	H 1	O 2	0	0
4	C	1	Total 4	C 1	H 1	O 2	0	0
4	C	1	Total 4	C 1	H 1	O 2	0	0
4	C	1	Total 4	C 1	H 1	O 2	0	0
4	C	1	Total 4	C 1	H 1	O 2	0	0
4	C	1	Total 4	C 1	H 1	O 2	0	0
4	C	1	Total 4	C 1	H 1	O 2	0	0
4	C	1	Total 4	C 1	H 1	O 2	0	0
4	C	1	Total 4	C 1	H 1	O 2	0	0

- Molecule 5 is POLYETHYLENE GLYCOL (N=34) (three-letter code: 15P) (formula:  $C_{69}H_{140}O_{35}$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
5	C	1	Total 36	C 12	H 16	O 8	0	1

- Molecule 6 is water.


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	A	383	Total 383	O 383	0	0
6	B	408	Total 408	O 408	0	0
6	C	396	Total 396	O 396	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HE protein



- Molecule 1: HE protein



- Molecule 1: HE protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	142.42Å 93.29Å 96.72Å 90.00° 122.01° 90.00°	Depositor
Resolution (Å)	47.72 – 1.80 47.72 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.72-1.80) 96.7 (47.72-1.80)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.50 (at 1.79Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.137 , 0.168 0.141 , 0.170	Depositor DCC
$R_{free}$ test set	2000 reflections (2.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.2	Xtrriage
Anisotropy	0.508	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	15896	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 15P, MG, GOL, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/2524	0.66	1/3435 (0.0%)
1	B	0.43	0/2495	0.65	1/3394 (0.0%)
1	C	0.43	0/2523	0.66	1/3427 (0.0%)
All	All	0.44	0/7542	0.66	3/10256 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	197	LEU	CA-CB-CG	5.55	128.06	115.30
1	C	197	LEU	CA-CB-CG	5.38	127.68	115.30
1	A	35	ASP	CB-CG-OD1	5.30	123.07	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2461	2406	2403	10	0
1	B	2431	2365	2363	15	0
1	C	2460	2413	2422	12	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	6	8	8	0	0
3	B	12	16	16	1	0
4	A	15	5	5	0	0
4	B	27	9	9	8	0
4	C	27	9	9	1	0
5	C	20	16	24	2	0
6	A	383	0	0	2	0
6	B	408	0	0	2	0
6	C	396	0	0	2	0
All	All	8649	7247	7259	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:25:ASP:OD2	4:B:408:FMT:H	1.80	0.82
1:C:237[A]:MET:SD	6:C:772:HOH:O	2.46	0.72
1:B:25:ASP:HB3	4:B:408:FMT:H	1.86	0.57
1:B:169[B]:ILE:HD11	1:B:192:TYR:CD1	2.40	0.57
1:B:169[B]:ILE:HD11	1:B:192:TYR:HD1	1.71	0.56
1:B:25:ASP:OD2	4:B:408:FMT:C	2.54	0.54
1:C:139:ARG:O	5:C:410[B]:15P:H26	2.08	0.54
1:A:326:MET:HE1	6:A:854:HOH:O	2.08	0.54
1:B:293:GLU:OE1	1:B:293:GLU:OE2	2.30	0.50
1:C:137:VAL:HG22	1:C:151:LEU:CD2	2.41	0.50
1:B:25:ASP:CB	4:B:408:FMT:H	2.41	0.50
4:B:404:FMT:O2	6:B:501:HOH:O	2.18	0.49
1:B:293:GLU:OE1	1:B:293:GLU:CG	2.60	0.49
1:C:89:ASN:HD21	4:C:407:FMT:C	2.26	0.49
1:B:25:ASP:CG	4:B:408:FMT:H	2.33	0.48
3:B:403:GOL:O1	6:B:502:HOH:O	2.19	0.48
1:B:168:GLY:N	4:B:407:FMT:H	2.30	0.46
1:A:326:MET:HE3	1:C:327:ILE:HG21	1.97	0.46
1:A:63[A]:LEU:HD23	1:A:113:LEU:CD1	2.46	0.45
1:A:132:ALA:HA	5:C:410[B]:15P:H28	1.98	0.45
1:A:66:MET:CE	1:A:113:LEU:HD21	2.46	0.45
1:C:164[A]:ASN:OD1	6:C:501:HOH:O	2.21	0.45
1:A:24:PRO:HA	1:A:52:GLY:HA2	2.00	0.44
1:C:332:ARG:C	1:C:334:ILE:H	2.20	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:GLU:HB3	1:A:169:ILE:HD11	1.99	0.44
1:B:112:GLU:HB3	1:B:169[A]:ILE:HD11	1.99	0.43
1:A:330:LEU:HD21	1:C:331:GLN:HG2	2.00	0.43
1:A:300:MET:HE3	6:A:649:HOH:O	2.18	0.43
1:B:24:PRO:HA	1:B:52:GLY:HA2	2.00	0.43
1:B:168:GLY:H	4:B:407:FMT:H	1.83	0.43
1:C:137:VAL:HG22	1:C:151:LEU:HD21	2.00	0.42
1:A:334:ILE:CG1	1:A:335:THR:N	2.82	0.42
1:C:112:GLU:HB3	1:C:169:ILE:HD11	2.02	0.42
1:C:332:ARG:C	1:C:334:ILE:N	2.74	0.41
1:C:115:LEU:HD21	1:C:242:LEU:HD22	2.03	0.41
1:B:37:SER:HB2	1:B:38:ARG:HG3	2.02	0.41
1:B:66:MET:HE1	1:B:170:TYR:HA	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/342 (95%)	318 (98%)	7 (2%)	0	100	100
1	B	324/342 (95%)	320 (99%)	4 (1%)	0	100	100
1	C	323/342 (94%)	317 (98%)	6 (2%)	0	100	100
All	All	972/1026 (95%)	955 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/298 (94%)	279 (99%)	2 (1%)	81	79
1	B	272/298 (91%)	267 (98%)	5 (2%)	54	45
1	C	281/298 (94%)	276 (98%)	5 (2%)	54	45
All	All	834/894 (93%)	822 (99%)	12 (1%)	60	56

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	215	ARG
1	A	221	CYS
1	B	93	ASP
1	B	215	ARG
1	B	221	CYS
1	B	323	HIS
1	B	328	SER
1	C	93	ASP
1	C	215	ARG
1	C	221	CYS
1	C	329	LYS
1	C	335	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	203	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 31 ligands modelled in this entry, 3 are monoatomic - leaving 28 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	FMT	A	408	-	2,2,2	0.72	0	1,1,1	0.09	0
4	FMT	B	408	-	2,2,2	0.30	0	1,1,1	0.41	0
4	FMT	C	408	-	2,2,2	0.64	0	1,1,1	0.18	0
4	FMT	A	407	-	2,2,2	0.69	0	1,1,1	0.39	0
4	FMT	C	405	-	2,2,2	0.69	0	1,1,1	0.40	0
4	FMT	C	404	-	2,2,2	0.64	0	1,1,1	0.28	0
4	FMT	B	406	-	2,2,2	0.74	0	1,1,1	0.41	0
4	FMT	A	406	-	2,2,2	0.76	0	1,1,1	0.43	0
3	GOL	A	403	-	5,5,5	0.39	0	5,5,5	0.35	0
3	GOL	B	402	-	5,5,5	0.36	0	5,5,5	0.96	0
3	GOL	B	403	-	5,5,5	0.36	0	5,5,5	0.56	0
4	FMT	B	407	-	2,2,2	0.77	0	1,1,1	0.28	0
4	FMT	C	402	-	2,2,2	0.70	0	1,1,1	0.41	0
4	FMT	A	404	-	2,2,2	0.73	0	1,1,1	0.30	0
4	FMT	C	406	-	2,2,2	0.87	0	1,1,1	0.41	0
4	FMT	B	404	-	2,2,2	0.74	0	1,1,1	0.41	0
4	FMT	B	412	-	2,2,2	0.58	0	1,1,1	0.21	0
4	FMT	A	405	-	2,2,2	0.70	0	1,1,1	0.44	0
5	15P	C	410[A]	-	9,9,103	0.72	0	8,8,102	0.27	0
4	FMT	B	405	-	2,2,2	0.62	0	1,1,1	0.30	0
4	FMT	B	409	-	2,2,2	0.66	0	1,1,1	0.36	0
4	FMT	C	409	-	2,2,2	0.73	0	1,1,1	0.44	0
4	FMT	C	403	-	2,2,2	0.69	0	1,1,1	0.39	0
5	15P	C	410[B]	-	9,9,103	0.72	0	8,8,102	0.24	0
4	FMT	B	410	-	2,2,2	0.68	0	1,1,1	0.41	0
4	FMT	C	407	-	2,2,2	0.70	0	1,1,1	0.25	0
4	FMT	B	411	-	2,2,2	0.75	0	1,1,1	0.40	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	FMT	C	401	-	2,2,2	0.74	0	1,1,1	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	402	-	-	2/4/4/4	-
3	GOL	B	403	-	-	2/4/4/4	-
5	15P	C	410[B]	-	-	5/7/7/101	-
5	15P	C	410[A]	-	-	4/7/7/101	-
3	GOL	A	403	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	402	GOL	O1-C1-C2-C3
3	B	403	GOL	O1-C1-C2-C3
5	C	410[B]	15P	O3-C7-C8-O4
5	C	410[B]	15P	O2-C5-C6-O3
5	C	410[A]	15P	O2-C5-C6-O3
5	C	410[B]	15P	O1-C3-C4-O2
3	B	403	GOL	O1-C1-C2-O2
5	C	410[A]	15P	O3-C7-C8-O4
3	B	402	GOL	O1-C1-C2-O2
5	C	410[A]	15P	C3-C4-O2-C5
5	C	410[B]	15P	C3-C4-O2-C5
5	C	410[B]	15P	C5-C6-O3-C7
5	C	410[A]	15P	C5-C6-O3-C7

There are no ring outliers.

6 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	408	FMT	5	0


*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	403	GOL	1	0
4	B	407	FMT	2	0
4	B	404	FMT	1	0
5	C	410[B]	15P	2	0
4	C	407	FMT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	321/342 (93%)	-0.60	8 (2%) 58 57	11, 26, 67, 156	6 (1%)
1	B	319/342 (93%)	-0.57	8 (2%) 58 57	11, 27, 79, 113	7 (2%)
1	C	319/342 (93%)	-0.60	7 (2%) 62 60	10, 27, 71, 135	6 (1%)
All	All	959/1026 (93%)	-0.59	23 (2%) 59 58	10, 27, 73, 156	19 (1%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	335	THR	6.1
1	B	335	THR	5.6
1	B	322	LEU	4.9
1	A	328	SER	4.6
1	C	334	ILE	4.5
1	C	309	LEU	4.4
1	A	330	LEU	4.3
1	C	327	ILE	3.5
1	A	315	LEU	3.2
1	A	337	VAL	3.1
1	A	329	LYS	3.1
1	B	334	ILE	2.9
1	B	325	GLU	2.8
1	B	330	LEU	2.8
1	B	327	ILE	2.6
1	C	308	SER	2.6
1	C	332	ARG	2.6
1	C	307	THR	2.6
1	A	334	ILE	2.2
1	B	329	LYS	2.1
1	B	333	ASP	2.1
1	A	304	PHE	2.0
1	A	335	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	403	6/6	0.73	0.25	85,102,105,105	0
5	15P	C	410[A]	10/104	0.76	0.24	66,68,81,81	18
5	15P	C	410[B]	10/104	0.76	0.24	66,68,81,81	18
4	FMT	B	406	3/3	0.78	0.18	72,72,72,87	0
4	FMT	B	404	3/3	0.79	0.23	86,86,86,104	0
4	FMT	B	410	3/3	0.79	0.24	63,65,67,78	0
4	FMT	B	408	3/3	0.80	0.29	72,76,78,91	0
3	GOL	B	402	6/6	0.81	0.18	41,63,75,82	0
4	FMT	C	409	3/3	0.81	0.38	92,93,94,112	0
4	FMT	B	409	3/3	0.82	0.14	50,50,51,61	0
4	FMT	A	407	3/3	0.84	0.15	43,49,53,58	0
4	FMT	B	407	3/3	0.84	0.18	20,29,34,36	4
4	FMT	C	408	3/3	0.86	0.25	46,47,47,57	0
4	FMT	A	404	3/3	0.86	0.14	39,44,48,53	0
4	FMT	A	405	3/3	0.87	0.19	55,60,64,72	0
4	FMT	C	403	3/3	0.88	0.14	59,61,63,74	0
4	FMT	C	405	3/3	0.88	0.18	62,64,65,77	0
4	FMT	B	411	3/3	0.88	0.26	80,81,82,97	0
4	FMT	C	401	3/3	0.89	0.12	41,41,41,50	0
4	FMT	C	402	3/3	0.89	0.16	53,58,63,70	0
4	FMT	C	406	3/3	0.90	0.23	37,48,58,58	0
4	FMT	C	407	3/3	0.91	0.15	40,47,53,57	0
4	FMT	B	405	3/3	0.91	0.11	36,38,40,46	0
4	FMT	C	404	3/3	0.91	0.14	30,30,31,36	4
3	GOL	A	403	6/6	0.91	0.13	29,46,62,64	0
4	FMT	A	406	3/3	0.91	0.11	30,35,38,43	0
4	FMT	B	412	3/3	0.93	0.12	28,35,37,42	0


*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	FMT	A	408	3/3	0.93	0.18	30,34,36,41	0
2	MG	A	401	1/1	0.94	0.08	42,42,42,42	0
2	MG	A	402	1/1	0.94	0.07	58,58,58,58	0
2	MG	B	401	1/1	0.99	0.06	40,40,40,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.