



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2023 – 02:21 PM EDT

PDB ID : 5U3Q
Title : Human PPARdelta ligand-binding domain in complexed with specific agonist 1
Authors : Wu, C.-C.; Baiga, T.J.; Downes, M.; La Clair, J.J.; Atkins, A.R.; Richard, S.B.; Stockley-Noel, T.A.; Bowman, M.E.; Evans, R.M.; Noel, J.P.
Deposited on : 2016-12-03
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	 4% 92% 5% . .
1	B	272	 3% 94% 5%

2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 10671 atoms, of which 5165 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisome proliferator-activated receptor delta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	265	Total	C	H	N	O	S	0	33	0
			4869	1559	2456	404	439	11			
1	B	271	Total	C	H	N	O	S	0	20	0
			4717	1509	2385	394	418	11			

- Molecule 2 is 6-(2-[[[(1,1'-biphenyl)-4-carbonyl](propan-2-yl)amino]methyl]phenoxy)hexanoic acid (three-letter code: 7UJ) (formula: C₂₉H₃₃NO₄).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	Total	C	H	N	O	0	1
			132	58	64	2	8		
2	B	1	Total	C	H	N	O	0	0
			66	29	32	1	4		

- Molecule 3 is heptyl beta-D-glucopyranoside (three-letter code: B7G) (formula: C₁₃H₂₆O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	45	13	26	6	0	0
3	B	1	45	13	26	6	0	0

- Molecule 4 is S-1,2-PROPANEDIOL (three-letter code: PGO) (formula: $C_3H_8O_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	13	3	8	2	0	0
4	A	1	13	3	8	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			13	3	8	2		
4	A	1	Total	C	H	O	0	0
			13	3	8	2		
4	A	1	Total	C	H	O	0	1
			26	6	16	4		
4	B	1	Total	C	H	O	0	1
			26	6	16	4		
4	B	1	Total	C	H	O	0	1
			26	6	16	4		
4	B	1	Total	C	H	O	0	1
			26	6	16	4		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			17	4	10	3		
5	A	1	Total	C	H	O	0	0
			17	4	10	3		
5	A	1	Total	C	H	O	0	0
			17	4	10	3		
5	B	1	Total	C	H	O	0	1
			34	8	20	6		
5	B	1	Total	C	H	O	0	0
			17	4	10	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
5	B	1	Total	C	H	O	0	0
			17	4	10	3		
5	B	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 6 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	K	0	0
			1	1		
6	B	1	Total	K	0	0
			1	1		

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	2	Total	Cl	0	0
			2	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	238	Total	O	0	0
			238	238		
8	B	263	Total	O	0	0
			263	263		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor delta



- Molecule 1: Peroxisome proliferator-activated receptor delta



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	39.46Å 94.35Å 96.73Å 90.00° 98.36° 90.00°	Depositor
Resolution (Å)	47.85 – 1.50 47.85 – 1.50	Depositor EDS
% Data completeness (in resolution range)	82.8 (47.85-1.50) 82.9 (47.85-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 1.50Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.159 , 0.189 0.160 , 0.190	Depositor DCC
R_{free} test set	1662 reflections (1.79%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtrriage
Anisotropy	0.503	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 52.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10671	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, B7G, CL, K, 7UJ, PGO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.47	1/2464 (0.0%)	0.58	0/3327
1	B	0.51	0/2397	0.65	3/3239 (0.1%)
All	All	0.49	1/4861 (0.0%)	0.62	3/6566 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	251	CYS	CB-SG	-5.13	1.73	1.81

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	286	VAL	CG1-CB-CG2	8.94	125.21	110.90
1	B	410	VAL	CG1-CB-CG2	5.39	119.52	110.90
1	B	286	VAL	CA-CB-CG2	5.21	118.71	110.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	205	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2413	2456	2427	9	1
1	B	2332	2385	2365	8	1
2	A	68	64	0	0	0
2	B	34	32	0	0	0
3	A	19	26	26	0	0
3	B	19	26	26	1	0
4	A	30	48	48	1	0
4	B	30	48	48	1	0
5	A	21	30	30	1	0
5	B	35	50	50	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	B	2	0	0	0	0
8	A	238	0	0	2	0
8	B	263	0	0	2	0
All	All	5506	5165	5020	19	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:GLY:HA3	4:A:508:PGO:H33	1.89	0.53
1:A:429:LEU:HB3	8:A:611:HOH:O	2.12	0.49
1:A:235:LEU:HD21	1:A:251:CYS:SG	2.53	0.49
1:B:191:ASN:HB2	1:B:259:GLU:OE2	2.13	0.48
1:B:230:GLN:HB2	1:B:251[B]:CYS:SG	2.54	0.47
1:A:269[B]:SER:OG	1:A:369:ARG:NH2	2.49	0.45
1:B:207[B]:HIS:ND1	8:B:605:HOH:O	2.36	0.45
8:A:749:HOH:O	3:B:502:B7G:H131	2.17	0.44
1:B:230:GLN:CB	1:B:251[B]:CYS:SG	3.05	0.44
1:B:206:SER:OG	1:B:208:THR:N	2.43	0.44
1:B:357[B]:LEU:O	1:B:374[B]:GLN:HB2	2.19	0.42
1:A:327[B]:ILE:HD12	1:A:416:MET:HB3	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:192:MET:HG3	1:B:196[B]:LYS:HD2	2.02	0.41
1:A:219:LEU:HD12	1:A:241:ILE:CD1	2.50	0.41
4:B:507[B]:PGO:H31	8:B:640:HOH:O	2.20	0.41
1:A:327[A]:ILE:HD11	1:A:413:HIS:CE1	2.56	0.41
1:A:220:TRP:CZ2	1:A:224:LYS:HD2	2.56	0.41
1:B:207[B]:HIS:CD2	1:B:208:THR:HG23	2.57	0.40
1:A:238:TYR:CE1	5:A:505:PEG:H41	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:TYR:HH	1:B:382[B]:GLU:OE2[2_858]	1.53	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/272 (108%)	292 (99%)	2 (1%)	0	100	100
1	B	289/272 (106%)	282 (98%)	7 (2%)	0	100	100
All	All	583/544 (107%)	574 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/240 (111%)	265 (99%)	2 (1%)	84	69
1	B	259/240 (108%)	258 (100%)	1 (0%)	91	82
All	All	526/480 (110%)	523 (99%)	3 (1%)	86	74

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	219	LEU
1	A	393	GLN
1	B	286	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 29 ligands modelled in this entry, 4 are monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	PEG	A	509	-	6,6,6	0.64	0	5,5,5	0.33	0
4	PGO	B	506[B]	-	3,4,4	0.31	0	1,4,4	0.53	0
4	PGO	B	507[B]	-	3,4,4	0.52	0	1,4,4	0.15	0
4	PGO	B	506[A]	-	3,4,4	0.39	0	1,4,4	0.17	0
4	PGO	B	507[A]	-	3,4,4	0.27	0	1,4,4	0.01	0
4	PGO	A	507	-	3,4,4	0.33	0	1,4,4	0.02	0
4	PGO	A	510[B]	-	3,4,4	0.23	0	1,4,4	0.14	0
2	7UJ	A	501[B]	-	36,36,36	0.53	0	46,47,47	1.41	5 (10%)
2	7UJ	A	501[A]	-	36,36,36	0.52	0	46,47,47	1.49	7 (15%)
5	PEG	B	511	-	6,6,6	0.44	0	5,5,5	0.38	0
5	PEG	A	505	-	6,6,6	0.56	0	5,5,5	0.31	0
4	PGO	A	510[A]	-	3,4,4	0.25	0	1,4,4	0.05	0
4	PGO	A	503	-	3,4,4	0.31	0	1,4,4	0.04	0
5	PEG	B	504	-	6,6,6	0.54	0	5,5,5	0.52	0
3	B7G	B	502	-	19,19,19	0.80	0	24,24,24	1.12	1 (4%)
5	PEG	B	505	-	6,6,6	0.57	0	5,5,5	0.30	0
4	PGO	A	506	-	3,4,4	0.28	0	1,4,4	0.13	0
5	PEG	A	504	-	6,6,6	0.55	0	5,5,5	0.38	0
4	PGO	B	508[B]	-	3,4,4	0.26	0	1,4,4	0.06	0
4	PGO	B	508[A]	-	3,4,4	0.32	0	1,4,4	0.21	0
5	PEG	B	503[B]	-	6,6,6	0.56	0	5,5,5	0.29	0
4	PGO	A	508	-	3,4,4	0.34	0	1,4,4	0.12	0
5	PEG	B	503[A]	-	6,6,6	0.57	0	5,5,5	0.27	0
2	7UJ	B	501	-	36,36,36	0.61	0	46,47,47	1.36	8 (17%)
3	B7G	A	502	-	19,19,19	0.79	0	24,24,24	1.06	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PEG	A	509	-	-	2/4/4/4	-
4	PGO	B	506[B]	-	-	1/2/2/2	-
4	PGO	B	507[B]	-	-	1/2/2/2	-
4	PGO	B	506[A]	-	-	2/2/2/2	-
4	PGO	B	507[A]	-	-	0/2/2/2	-
4	PGO	A	507	-	-	0/2/2/2	-
4	PGO	A	510[B]	-	-	0/2/2/2	-
2	7UJ	A	501[B]	-	-	4/29/29/29	0/3/3/3
2	7UJ	A	501[A]	-	-	5/29/29/29	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PEG	B	511	-	-	3/4/4/4	-
5	PEG	A	505	-	-	3/4/4/4	-
4	PGO	A	510[A]	-	-	2/2/2/2	-
4	PGO	A	503	-	-	2/2/2/2	-
5	PEG	B	504	-	-	4/4/4/4	-
3	B7G	B	502	-	-	2/10/30/30	0/1/1/1
5	PEG	B	505	-	-	2/4/4/4	-
4	PGO	A	506	-	-	2/2/2/2	-
5	PEG	A	504	-	-	1/4/4/4	-
4	PGO	B	508[B]	-	-	0/2/2/2	-
4	PGO	B	508[A]	-	-	2/2/2/2	-
5	PEG	B	503[B]	-	-	2/4/4/4	-
4	PGO	A	508	-	-	0/2/2/2	-
5	PEG	B	503[A]	-	-	1/4/4/4	-
2	7UJ	B	501	-	-	6/29/29/29	0/3/3/3
3	B7G	A	502	-	-	1/10/30/30	0/1/1/1

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501[B]	7UJ	C14-C13-N	-3.97	105.64	111.01
2	A	501[B]	7UJ	C15-C13-N	3.59	115.88	111.01
2	A	501[B]	7UJ	C6-N-C16	3.47	126.62	116.97
2	A	501[A]	7UJ	C6-N-C16	3.44	126.54	116.97
2	A	501[A]	7UJ	C15-C13-N	3.43	115.65	111.01
2	A	501[A]	7UJ	C14-C13-N	-3.39	106.42	111.01
2	A	501[A]	7UJ	O3-C16-C17	-3.17	114.05	120.23
2	A	501[B]	7UJ	O3-C16-C17	-3.10	114.19	120.23
2	A	501[A]	7UJ	C10-C11-C12	-3.01	106.88	114.47
2	B	501	7UJ	O3-C16-C17	-3.00	114.39	120.23
2	B	501	7UJ	C15-C13-N	2.77	114.77	111.01
2	B	501	7UJ	C6-N-C16	2.66	124.38	116.97
2	B	501	7UJ	C17-C16-N	2.60	123.94	118.77
2	A	501[B]	7UJ	C17-C16-N	2.49	123.72	118.77
2	A	501[A]	7UJ	C17-C16-N	2.44	123.62	118.77
2	A	501[A]	7UJ	O1-C12-C11	-2.36	115.51	123.08
2	B	501	7UJ	O1-C12-C11	-2.30	115.70	123.08
2	B	501	7UJ	C10-C11-C12	-2.28	108.72	114.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	B7G	O5-C5-C4	2.25	113.78	109.69
2	B	501	7UJ	C22-C18-C17	-2.25	118.16	120.78
3	A	502	B7G	O5-C5-C4	2.18	113.66	109.69
2	B	501	7UJ	C14-C13-N	-2.06	108.22	111.01

There are no chirality outliers.

All (48) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	506	PGO	O1-C1-C2-O2
4	A	510[A]	PGO	O1-C1-C2-O2
4	B	506[A]	PGO	O1-C1-C2-C3
4	B	506[A]	PGO	O1-C1-C2-O2
4	B	508[A]	PGO	O1-C1-C2-O2
5	B	511	PEG	O1-C1-C2-O2
5	B	505	PEG	O1-C1-C2-O2
2	A	501[A]	7UJ	C9-C10-C11-C12
5	B	504	PEG	O1-C1-C2-O2
5	B	503[B]	PEG	O2-C3-C4-O4
5	A	504	PEG	O2-C3-C4-O4
5	B	504	PEG	O2-C3-C4-O4
5	B	511	PEG	O2-C3-C4-O4
4	A	503	PGO	O1-C1-C2-O2
2	B	501	7UJ	C15-C13-N-C16
5	B	503[B]	PEG	O1-C1-C2-O2
3	B	502	B7G	C9-C10-C11-C12
5	A	505	PEG	O2-C3-C4-O4
5	A	509	PEG	C4-C3-O2-C2
4	A	503	PGO	O1-C1-C2-C3
4	A	506	PGO	O1-C1-C2-C3
4	A	510[A]	PGO	O1-C1-C2-C3
4	B	508[A]	PGO	O1-C1-C2-C3
5	B	505	PEG	C1-C2-O2-C3
2	A	501[B]	7UJ	C15-C13-N-C6
5	B	503[A]	PEG	C4-C3-O2-C2
2	B	501	7UJ	C9-C10-C11-C12
5	A	509	PEG	O2-C3-C4-O4
2	B	501	7UJ	C15-C13-N-C6
3	A	502	B7G	C7-C8-C9-C10
5	B	504	PEG	C1-C2-O2-C3
2	A	501[B]	7UJ	C15-C13-N-C16
3	B	502	B7G	C10-C11-C12-C13

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
5	B	511	PEG	C1-C2-O2-C3
5	A	505	PEG	C1-C2-O2-C3
2	B	501	7UJ	C10-C11-C12-O1
2	A	501[A]	7UJ	C10-C11-C12-O1
2	A	501[A]	7UJ	C10-C11-C12-O2
2	B	501	7UJ	C10-C11-C12-O2
5	A	505	PEG	O1-C1-C2-O2
5	B	504	PEG	C4-C3-O2-C2
2	A	501[B]	7UJ	C10-C11-C12-O2
4	B	506[B]	PGO	O1-C1-C2-O2
4	B	507[B]	PGO	O1-C1-C2-O2
2	A	501[A]	7UJ	C15-C13-N-C16
2	A	501[A]	7UJ	C14-C13-N-C16
2	A	501[B]	7UJ	C14-C13-N-C16
2	B	501	7UJ	C14-C13-N-C16



There are no ring outliers.


4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	507[B]	PGO	1	0
5	A	505	PEG	1	0
3	B	502	B7G	1	0
4	A	508	PGO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/272 (97%)	-0.04	10 (3%) 40 44	14, 26, 47, 85	0
1	B	271/272 (99%)	-0.10	8 (2%) 50 55	11, 21, 42, 81	0
All	All	536/544 (98%)	-0.07	18 (3%) 45 49	11, 24, 46, 85	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	393	GLN	5.5
1	A	208	THR	5.1
1	B	208	THR	4.7
1	B	205	ALA	4.4
1	A	173	ALA	3.9
1	B	229	LYS	3.2
1	A	203	GLY	3.2
1	B	204	LYS	3.1
1	B	233[B]	ASN	2.8
1	A	441	TYR	2.6
1	A	426[A]	GLU	2.2
1	A	227	VAL	2.2
1	B	232	VAL	2.2
1	A	390	PRO	2.1
1	B	207[A]	HIS	2.0
1	A	394	TYR	2.0
1	B	439	ASP	2.0
1	A	392	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PGO	A	510[A]	5/5	0.56	0.25	41,58,67,70	13
4	PGO	A	510[B]	5/5	0.56	0.25	49,59,68,68	13
6	K	B	512	1/1	0.63	0.19	77,77,77,77	0
5	PEG	B	505	7/7	0.73	0.23	42,73,90,90	0
4	PGO	A	507	5/5	0.73	0.18	44,58,82,82	0
5	PEG	B	503[A]	7/7	0.75	0.19	58,70,83,83	17
5	PEG	B	503[B]	7/7	0.75	0.19	54,70,82,84	17
4	PGO	B	508[A]	5/5	0.77	0.17	38,50,54,64	13
4	PGO	B	508[B]	5/5	0.77	0.17	21,41,50,54	13
4	PGO	B	506[A]	5/5	0.82	0.19	29,40,49,49	13
4	PGO	B	506[B]	5/5	0.82	0.19	30,38,48,50	13
5	PEG	B	511	7/7	0.83	0.28	32,59,89,89	0
4	PGO	A	506	5/5	0.83	0.20	41,69,80,85	0
5	PEG	A	509	7/7	0.84	0.11	42,52,64,66	0
3	B7G	B	502	19/19	0.84	0.11	31,47,63,68	0
4	PGO	B	507[A]	5/5	0.85	0.16	23,35,43,46	13
5	PEG	A	504	7/7	0.85	0.11	42,60,72,72	0
4	PGO	B	507[B]	5/5	0.85	0.16	24,34,40,42	13
4	PGO	A	508	5/5	0.85	0.18	46,61,76,81	0
4	PGO	A	503	5/5	0.86	0.20	54,68,93,97	0
3	B7G	A	502	19/19	0.89	0.10	26,41,56,61	0
5	PEG	B	504	7/7	0.91	0.20	31,63,78,78	0
5	PEG	A	505	7/7	0.92	0.14	48,59,71,81	0
2	7UJ	A	501[B]	34/34	0.96	0.07	16,22,30,34	66
2	7UJ	A	501[A]	34/34	0.96	0.07	15,22,33,34	66
2	7UJ	B	501	34/34	0.97	0.08	11,17,31,31	0
7	CL	B	510	1/1	0.97	0.05	38,38,38,38	0
6	K	A	511	1/1	0.98	0.04	31,31,31,31	0
7	CL	B	509	1/1	0.99	0.06	24,24,24,24	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around 7UJ A 501 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 7UJ A 501 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.