



Full wwPDB NMR Structure Validation Report ⓘ

Dec 25, 2024 – 05:30 AM EST

PDB ID : 5WXE
BMRB ID : 36041
Title : Highly disulfide-constrained antifeedant jasmintides from Jasminum sambac flowers
Authors : Kumari, G.; Wong, K.H.; Serra, A.; Shin, J.; Yoon, H.S.; Sze, S.K.; James, P.T.
Deposited on : 2017-01-07

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 51%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	28	<div> <div>43%</div> <div>46%</div> <div>7%</div> <div>.</div> </div>

2 Ensemble composition and analysis

This entry contains 20 models. Model 19 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:28 (27)	0.35	19

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 6 single-model clusters were found.

Cluster number	Models
1	8, 9, 12, 18, 19
2	1, 4, 7, 20
3	3, 6, 10
4	5, 17
Single-model clusters	2; 11; 13; 14; 15; 16

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 429 atoms, of which 209 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called jasmintide js3.


Mol	Chain	Residues	Atoms						Trace
1	A	28	Total	C	H	N	O	S	0
			429	133	209	39	42	6	

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: jasmintide js3




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: jasmintide js3



4.2.2 Score per residue for model 2

- Molecule 1: jasmintide js3



4.2.3 Score per residue for model 3

- Molecule 1: jasmintide js3




4.2.4 Score per residue for model 4

- Molecule 1: jasmintide js3




4.2.5 Score per residue for model 5

- Molecule 1: jasmintide js3




4.2.6 Score per residue for model 6

- Molecule 1: jasmintide js3



4.2.7 Score per residue for model 7

- Molecule 1: jasmintide js3




4.2.8 Score per residue for model 8

- Molecule 1: jasmintide js3




4.2.9 Score per residue for model 9

- Molecule 1: jasmintide js3



4.2.10 Score per residue for model 10

- Molecule 1: jasmintide js3




4.2.11 Score per residue for model 11

- Molecule 1: jasmintide js3




4.2.12 Score per residue for model 12

- Molecule 1: jasmintide js3




4.2.13 Score per residue for model 13

- Molecule 1: jasmintide js3




4.2.14 Score per residue for model 14

- Molecule 1: jasmintide js3




4.2.15 Score per residue for model 15

- Molecule 1: jasmintide js3




4.2.16 Score per residue for model 16

- Molecule 1: jasmintide js3




4.2.17 Score per residue for model 17

- Molecule 1: jasmintide js3



4.2.18 Score per residue for model 18

- Molecule 1: jasmintide js3




4.2.19 Score per residue for model 19 (medoid)

- Molecule 1: jasmintide js3



4.2.20 Score per residue for model 20

- Molecule 1: jasmintide js3



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure calculation	1.2
CNS	refinement	1.2

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	186
Number of shifts mapped to atoms	186
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	51%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	211	199	199	9±3
All	All	4220	3980	3980	188

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:4:LEU:HD11	1:A:17:TRP:CE3	0.78	2.13	6	11
1:A:2:LEU:HD23	1:A:17:TRP:CZ2	0.77	2.14	3	3
1:A:8:THR:HG22	1:A:11:ASP:HB2	0.74	1.60	16	6
1:A:19:VAL:HG11	1:A:28:SER:OXT	0.71	1.84	9	2
1:A:2:LEU:HD11	1:A:4:LEU:HD23	0.68	1.64	3	1
1:A:2:LEU:HD23	1:A:17:TRP:CH2	0.67	2.25	10	2
1:A:2:LEU:HD11	1:A:4:LEU:CD2	0.65	2.22	3	1
1:A:2:LEU:HD12	1:A:3:CYS:N	0.65	2.07	10	2
1:A:5:LEU:O	1:A:5:LEU:HD12	0.64	1.93	14	1
1:A:16:ILE:HD13	1:A:17:TRP:CD2	0.64	2.28	7	6
1:A:4:LEU:HD12	1:A:25:CYS:SG	0.62	2.34	15	2
1:A:8:THR:HG23	1:A:11:ASP:OD1	0.61	1.94	5	3
1:A:19:VAL:HG21	1:A:28:SER:HB3	0.61	1.70	16	1
1:A:16:ILE:HG23	1:A:17:TRP:CD2	0.61	2.30	16	4

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:14:TYR:O	1:A:18:THR:HG22	0.59	1.97	10	2
1:A:16:ILE:HD11	1:A:17:TRP:CE3	0.58	2.33	18	7
1:A:15:ILE:HG23	1:A:16:ILE:HG23	0.58	1.75	18	1
1:A:16:ILE:HD13	1:A:17:TRP:CE3	0.58	2.34	7	6
1:A:15:ILE:HG13	1:A:16:ILE:HG23	0.55	1.79	15	2
1:A:8:THR:HG23	1:A:10:ARG:H	0.54	1.63	1	7
1:A:4:LEU:HD11	1:A:17:TRP:CZ3	0.54	2.37	6	3
1:A:19:VAL:HG11	1:A:28:SER:CB	0.54	2.33	15	2
1:A:15:ILE:CD1	1:A:16:ILE:HG23	0.54	2.33	7	1
1:A:15:ILE:HD12	1:A:16:ILE:HG23	0.52	1.80	7	1
1:A:4:LEU:O	1:A:25:CYS:N	0.52	2.42	7	17
1:A:21:ARG:O	1:A:23:GLY:N	0.52	2.42	3	20
1:A:4:LEU:O	1:A:24:CYS:HA	0.51	2.06	6	3
1:A:16:ILE:HD11	1:A:17:TRP:CH2	0.51	2.40	14	2
1:A:16:ILE:HD11	1:A:17:TRP:CZ3	0.50	2.42	2	1
1:A:16:ILE:CD1	1:A:17:TRP:CE3	0.50	2.95	4	6
1:A:2:LEU:HD13	1:A:17:TRP:CD1	0.50	2.41	14	1
1:A:2:LEU:HD12	1:A:2:LEU:C	0.49	2.27	3	3
1:A:16:ILE:HD11	1:A:17:TRP:CZ2	0.48	2.44	14	1
1:A:2:LEU:HD12	1:A:4:LEU:CD2	0.47	2.38	16	1
1:A:4:LEU:HD11	1:A:17:TRP:CD2	0.47	2.44	14	2
1:A:16:ILE:HG23	1:A:17:TRP:CE3	0.47	2.44	16	4
1:A:4:LEU:CD1	1:A:17:TRP:CE3	0.46	2.99	13	2
1:A:16:ILE:HD12	1:A:27:ILE:HD13	0.45	1.87	16	2
1:A:15:ILE:HG23	1:A:16:ILE:N	0.45	2.26	20	2
1:A:15:ILE:HD12	1:A:16:ILE:N	0.45	2.26	7	1
1:A:16:ILE:O	1:A:17:TRP:CD1	0.45	2.70	7	16
1:A:8:THR:HG22	1:A:11:ASP:CB	0.45	2.39	7	2
1:A:12:CYS:HB2	1:A:14:TYR:CD2	0.44	2.46	14	2
1:A:4:LEU:CD1	1:A:17:TRP:CD2	0.44	3.00	14	1
1:A:20:CYS:HA	1:A:24:CYS:O	0.44	2.12	8	2
1:A:27:ILE:HG23	1:A:28:SER:N	0.44	2.28	14	3
1:A:2:LEU:CD2	1:A:17:TRP:CZ2	0.43	2.98	10	2
1:A:16:ILE:H	1:A:16:ILE:HD13	0.43	1.74	14	1
1:A:2:LEU:HG	1:A:17:TRP:CZ2	0.43	2.49	8	2
1:A:19:VAL:HG21	1:A:27:ILE:O	0.42	2.14	11	1
1:A:12:CYS:HB3	1:A:14:TYR:CD1	0.42	2.50	5	2
1:A:12:CYS:HB3	1:A:14:TYR:CD2	0.42	2.49	12	1
1:A:12:CYS:HB2	1:A:14:TYR:CD1	0.41	2.51	2	1
1:A:2:LEU:CD1	1:A:4:LEU:CD2	0.41	2.98	8	3
1:A:12:CYS:O	1:A:14:TYR:CD2	0.41	2.74	3	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:12:CYS:O	1:A:14:TYR:CD1	0.41	2.74	19	1
1:A:2:LEU:CD1	1:A:4:LEU:HD21	0.41	2.46	16	1
1:A:19:VAL:HG11	1:A:28:SER:O	0.40	2.16	1	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	26/28 (93%)	20±2 (75±6%)	5±2 (21±6%)	1±0 (4±0%)	4	31
All	All	520/560 (93%)	392 (75%)	108 (21%)	20 (4%)	4	31

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	22	ASP	20

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	26/27 (96%)	20±2 (76±6%)	6±2 (24±6%)	2	26
All	All	520/540 (96%)	397 (76%)	123 (24%)	2	26

All 19 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	2	LEU	18
1	A	16	ILE	16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	3	CYS	13
1	A	24	CYS	12
1	A	21	ARG	10
1	A	22	ASP	7
1	A	25	CYS	7
1	A	10	ARG	7
1	A	11	ASP	6
1	A	18	THR	5
1	A	12	CYS	5
1	A	28	SER	5
1	A	9	SER	3
1	A	26	ASN	3
1	A	15	ILE	2
1	A	8	THR	1
1	A	6	CYS	1
1	A	5	LEU	1
1	A	27	ILE	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 51% for the well-defined parts and 50% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *js3.bmr2*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	186
Number of shifts mapped to atoms	186
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 51%, i.e. 180 atoms were assigned a chemical shift out of a possible 355. 0 out of 4 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	55/136 (40%)	55/55 (100%)	0/54 (0%)	0/27 (0%)
Sidechain	116/198 (59%)	116/129 (90%)	0/60 (0%)	0/9 (0%)
Aromatic	9/21 (43%)	9/10 (90%)	0/10 (0%)	0/1 (0%)
Overall	180/355 (51%)	180/194 (93%)	0/124 (0%)	0/37 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 50%, i.e. 185 atoms were assigned a chemical shift out of a possible 370. 0 out of 4 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	56/141 (40%)	56/57 (98%)	0/56 (0%)	0/28 (0%)
Sidechain	120/208 (58%)	120/135 (89%)	0/63 (0%)	0/10 (0%)
Aromatic	9/21 (43%)	9/10 (90%)	0/10 (0%)	0/1 (0%)
Overall	185/370 (50%)	185/202 (92%)	0/129 (0%)	0/39 (0%)


7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis [i](#)

8.1 Conformationally restricting restraints [i](#)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	517
Intra-residue ($ i-j =0$)	138
Sequential ($ i-j =1$)	145
Medium range ($ i-j >1$ and $ i-j <5$)	48
Long range ($ i-j \geq 5$)	172
Inter-chain	0
Hydrogen bond restraints	14
Disulfide bond restraints	0
Total dihedral-angle restraints	23
Number of unmapped restraints	0
Number of restraints per residue	19.3
Number of long range restraints per residue ¹	6.5

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations [i](#)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model [i](#)

Distance violations less than 0.1 Å are not included in the calculation. There are no distance violations

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis ⓘ


9.1 Summary of distance violations ⓘ

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue ($i-j =0$)	138	26.7	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	26	5.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	85	16.4	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	27	5.2	0	0.0	0.0	0	0.0	0.0
Sequential ($i-j =1$)	145	28.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	58	11.2	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	77	14.9	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	10	1.9	0	0.0	0.0	0	0.0	0.0
Medium range ($i-j >1$ & $i-j <5$)	48	9.3	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	16	3.1	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	23	4.4	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	9	1.7	0	0.0	0.0	0	0.0	0.0
Long range ($i-j \geq 5$)	172	33.3	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	38	7.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	67	13.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	67	13.0	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	14	2.7	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	517	100.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	138	26.7	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	266	51.5	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	113	21.9	0	0.0	0.0	0	0.0	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

No violations found

9.3 Distance violation statistics for the ensemble [i](#)

No violations found

9.4 Most violated distance restraints in the ensemble [i](#)

No violations found

9.5 All violated distance restraints [i](#)

No violations found

10 Dihedral-angle violation analysis [i](#)


10.1 Summary of dihedral-angle violations [i](#)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
PHI	23	100.0	0	0.0	0.0	0	0.0	0.0
Total	23	100.0	0	0.0	0.0	0	0.0	0.0

¹ percentage calculated with respect to total number of dihedral-angle restraints, ² percentage calculated with respect to number of restraints in a particular dihedral-angle type, ³ violated in at least one model, ⁴ violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model [i](#)

No violations found

10.3 Dihedral-angle violation statistics for the ensemble [i](#)

No violations found

10.4 Most violated dihedral-angle restraints in the ensemble [i](#)

No violations found

10.5 All violated dihedral-angle restraints [i](#)

No violations found