



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 13, 2023 – 11:43 AM JST

PDB ID : 5X1M
Title : Vanillate/3-O-methylgallate O-demethylase, LigM, protocatechuate-tetrahydrofolate complex form
Authors : Harada, A.; Senda, T.
Deposited on : 2017-01-26
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	474	 91% 5% .
1	B	474	 2% 88% 6% 6%
1	C	474	 9% 86% 9% . .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11022 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Vanillate/3-O-methylgallate O-demethylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	456	3584	2302	608	661	13	0	6	0
1	B	447	3491	2239	594	645	13	0	2	0
1	C	455	3475	2234	587	640	14	0	1	0

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP G2IQS7
A	-1	SER	-	expression tag	UNP G2IQS7
A	0	SER	-	expression tag	UNP G2IQS7
B	-2	GLY	-	expression tag	UNP G2IQS7
B	-1	SER	-	expression tag	UNP G2IQS7
B	0	SER	-	expression tag	UNP G2IQS7
C	-2	GLY	-	expression tag	UNP G2IQS7
C	-1	SER	-	expression tag	UNP G2IQS7
C	0	SER	-	expression tag	UNP G2IQS7

- Molecule 2 is 3,4-DIHYDROXYBENZOIC ACID (three-letter code: DHB) (formula: C₇H₆O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	7	4		
2	A	1	Total	C	O	0	0
			11	7	4		
2	B	1	Total	C	O	0	0
			11	7	4		
2	C	1	Total	C	O	0	0
			11	7	4		

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	4	1	3		
3	A	1	Total	C	N	O	0	0
			8	4	1	3		
3	B	1	Total	C	N	O	0	0
			8	4	1	3		
3	C	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 4 is (6S)-5,6,7,8-TETRAHYDROFOLATE (three-letter code: THG) (formula: $C_{19}H_{23}N_7O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			32	19	7	6		
4	C	1	Total	C	N	O	0	0
			32	19	7	6		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	173	Total O 173 173	0	0
6	B	112	Total O 112 112	0	0
6	C	43	Total O 43 43	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Vanillate/3-O-methylgallate O-demethylase



- Molecule 1: Vanillate/3-O-methylgallate O-demethylase



- Molecule 1: Vanillate/3-O-methylgallate O-demethylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	103.80Å 118.37Å 132.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.53 – 1.90 47.92 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.53-1.90) 87.7 (47.92-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 1.90Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.208 , 0.242 0.209 , 0.244	Depositor DCC
R_{free} test set	6310 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	22.2	Xtrriage
Anisotropy	0.860	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11022	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: THG, EDO, TRS, DHB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/3701	0.57	0/5047
1	B	0.36	0/3595	0.55	0/4904
1	C	0.31	0/3577	0.51	0/4889
All	All	0.36	0/10873	0.54	0/14840

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3584	0	3469	13	0
1	B	3491	0	3338	19	0
1	C	3475	0	3271	27	0
2	A	22	0	10	1	0
2	B	11	0	5	2	0
2	C	11	0	5	1	0
3	A	16	0	24	2	0
3	B	8	0	12	0	0
3	C	8	0	12	0	0
4	B	32	0	21	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	32	0	21	3	0
5	B	4	0	6	0	0
6	A	173	0	0	0	0
6	B	112	0	0	1	0
6	C	43	0	0	1	0
All	All	11022	0	10194	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:452:TYR:O	6:B:601:HOH:O	2.10	0.69
1:C:108:ILE:HD11	4:C:501:THG:HC5	1.76	0.68
1:B:61[B]:MET:HE3	4:B:501:THG:HC71	1.76	0.67
1:C:256:TRP:CH2	4:C:501:THG:HC6	2.33	0.64
1:C:256:TRP:CZ2	4:C:501:THG:HC6	2.34	0.63
1:A:75:LEU:HB3	1:A:79:MET:CE	2.30	0.61
1:B:256:TRP:CH2	4:B:501:THG:HC6	2.35	0.60
1:C:414:PRO:HD2	1:C:417:THR:HG21	1.85	0.58
1:A:291:SER:O	2:A:502:DHB:H5	2.03	0.58
1:B:80:ILE:HD12	1:B:313:VAL:HG22	1.88	0.55
1:C:378:ASP:HB2	1:C:391:SER:HB3	1.89	0.54
1:B:61[B]:MET:HE2	2:B:502:DHB:H2	1.89	0.54
1:A:56:ASP:OD2	1:A:227[A]:ARG:NH1	2.41	0.53
1:C:414:PRO:O	1:C:417:THR:HG23	2.09	0.52
1:A:108:ILE:HG12	3:A:504:TRS:H32	1.94	0.49
1:C:45:GLN:HG2	1:C:48:TRP:CH2	2.48	0.49
1:A:336:LYS:HE3	1:A:450:VAL:HG21	1.93	0.49
1:B:45:GLN:HG2	1:B:48:TRP:CH2	2.47	0.49
1:C:386:ASN:O	1:C:388:VAL:HG13	2.13	0.48
1:C:119:TYR:OH	1:C:125:ALA:HB3	2.14	0.48
1:C:449:PRO:HD3	1:C:455:THR:HG21	1.96	0.48
1:C:353:LEU:HD23	6:C:640:HOH:O	2.14	0.47
1:B:108:ILE:HD12	4:B:501:THG:C2	2.44	0.47
1:C:63:ASP:OD2	1:C:163:ARG:NH1	2.47	0.47
1:C:261:LEU:HB2	1:C:375:THR:HG21	1.97	0.47
1:C:80:ILE:HD12	1:C:313:VAL:HG22	1.98	0.46
1:B:205:ARG:HA	1:B:214:LEU:HD23	1.99	0.45
1:C:61[A]:MET:CE	2:C:502:DHB:H2	2.46	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:120:VAL:HG11	3:A:504:TRS:H22	1.98	0.45
1:B:61[B]:MET:CE	2:B:502:DHB:H2	2.47	0.45
1:C:99:PRO:HD2	1:C:270:LEU:HD21	1.98	0.44
1:C:76:SER:OG	1:C:83:PRO:HG3	2.18	0.44
1:C:363:LYS:HB2	1:C:438:HIS:CD2	2.53	0.44
1:C:96:PRO:HB2	1:C:104:ILE:HD12	2.00	0.44
1:C:98:THR:HG21	1:C:263:ALA:O	2.18	0.44
1:A:362:TYR:CD1	1:A:441:MET:HB2	2.53	0.43
1:C:172:TRP:CZ2	1:C:185:LYS:HG2	2.53	0.43
1:C:119:TYR:CE2	1:C:126:ALA:HB2	2.53	0.43
1:C:290:GLY:O	1:C:425:GLU:HG2	2.18	0.43
1:B:362:TYR:CD1	1:B:441:MET:HB2	2.54	0.43
1:B:186:LEU:O	1:B:205:ARG:NE	2.51	0.43
1:B:52:ALA:O	1:B:239:LEU:HA	2.18	0.43
1:A:110:PHE:HB2	1:A:118[B]:VAL:HG22	2.01	0.42
1:C:258:PRO:HD2	1:C:393:PHE:CD1	2.54	0.42
1:B:48:TRP:HD1	1:B:49:ARG:HG2	1.85	0.42
1:B:298:GLU:HG2	1:B:301:TYR:CE2	2.55	0.42
1:B:61[B]:MET:CE	4:B:501:THG:HC71	2.48	0.42
1:A:32:PRO:HB2	1:A:33:VAL:HG23	2.02	0.41
1:A:296:ASN:HB3	1:A:298:GLU:OE2	2.20	0.41
1:C:302:VAL:HG21	1:C:307:ILE:HG23	2.03	0.41
1:B:255:GLY:HA3	1:B:450:VAL:HG23	2.03	0.41
1:A:172:TRP:CZ2	1:A:185:LYS:HG2	2.55	0.41
1:C:82:SER:O	1:C:92:LYS:HE3	2.21	0.41
1:B:89:ASN:OD1	1:B:202:ARG:NH2	2.53	0.41
1:C:206:HIS:HB3	1:C:207:GLY:H	1.68	0.41
1:A:64[A]:LEU:HD12	1:A:144:HIS:HA	2.04	0.40
1:B:256:TRP:CZ2	4:B:501:THG:HC6	2.56	0.40
1:A:303:ASN:HB2	1:A:304:PRO:HD2	2.03	0.40
1:B:40:ASN:O	1:B:44:GLU:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	460/474 (97%)	452 (98%)	8 (2%)	0	100	100
1	B	445/474 (94%)	435 (98%)	10 (2%)	0	100	100
1	C	454/474 (96%)	440 (97%)	14 (3%)	0	100	100
All	All	1359/1422 (96%)	1327 (98%)	32 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/382 (96%)	366 (100%)	1 (0%)	92	93
1	B	354/382 (93%)	350 (99%)	4 (1%)	73	73
1	C	339/382 (89%)	333 (98%)	6 (2%)	59	55
All	All	1060/1146 (92%)	1049 (99%)	11 (1%)	76	76

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	111	TYR
1	B	111	TYR
1	B	153	MET
1	B	163	ARG
1	B	254	SER
1	C	5	THR
1	C	48	TRP
1	C	111	TYR
1	C	163	ARG
1	C	205	ARG
1	C	206	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are

no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TRS	C	503	-	7,7,7	0.32	0	9,9,9	0.52	0
2	DHB	C	502	-	11,11,11	1.67	2 (18%)	15,15,15	1.00	1 (6%)
3	TRS	A	504	-	7,7,7	0.31	0	9,9,9	1.09	0
3	TRS	A	503	-	7,7,7	0.31	0	9,9,9	0.63	0
4	THG	B	501	-	32,34,34	2.43	9 (28%)	39,47,47	3.51	20 (51%)
5	EDO	B	504	-	3,3,3	0.38	0	2,2,2	0.92	0
2	DHB	A	501	-	11,11,11	1.62	3 (27%)	15,15,15	1.15	1 (6%)
3	TRS	B	503	-	7,7,7	0.18	0	9,9,9	0.66	0
2	DHB	B	502	-	11,11,11	1.76	3 (27%)	15,15,15	0.87	0
2	DHB	A	502	-	11,11,11	1.62	4 (36%)	15,15,15	0.98	0
4	THG	C	501	-	32,34,34	2.33	9 (28%)	39,47,47	1.93	8 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	C	503	-	-	3/9/9/9	-
2	DHB	C	502	-	-	0/4/4/4	0/1/1/1
3	TRS	A	504	-	-	3/9/9/9	-
3	TRS	A	503	-	-	3/9/9/9	-
4	THG	B	501	-	-	12/22/31/31	0/3/3/3
5	EDO	B	504	-	-	0/1/1/1	-
2	DHB	A	501	-	-	0/4/4/4	0/1/1/1
3	TRS	B	503	-	-	0/9/9/9	-
2	DHB	B	502	-	-	0/4/4/4	0/1/1/1
2	DHB	A	502	-	-	0/4/4/4	0/1/1/1
4	THG	C	501	-	-	6/22/31/31	0/3/3/3

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	501	THG	C4A-N5	7.58	1.53	1.38
4	C	501	THG	C4A-N5	7.53	1.53	1.38
4	B	501	THG	C11-N	6.03	1.47	1.34
4	C	501	THG	C11-N	5.50	1.46	1.34
4	B	501	THG	C2-N2	5.45	1.44	1.33
4	C	501	THG	C2-N2	5.44	1.44	1.33
2	B	502	DHB	O4-C4	3.88	1.44	1.36
2	C	502	DHB	O4-C4	3.81	1.44	1.36
4	C	501	THG	C7-N8	-3.74	1.38	1.44
4	B	501	THG	C7-N8	-3.64	1.38	1.44
2	A	501	DHB	O4-C4	3.30	1.43	1.36
2	A	502	DHB	O4-C4	3.30	1.43	1.36
4	B	501	THG	C4A-C4	-3.23	1.37	1.41
4	C	501	THG	C4A-C4	-3.19	1.37	1.41
4	B	501	THG	C4'-N10	2.48	1.45	1.38
2	A	501	DHB	O2-C	-2.39	1.23	1.30
2	C	502	DHB	O2-C	-2.38	1.23	1.30
2	B	502	DHB	O2-C	-2.30	1.23	1.30
4	C	501	THG	C2-N3	2.28	1.39	1.35
2	B	502	DHB	O1-C	2.16	1.29	1.22
2	A	502	DHB	O2-C	-2.15	1.24	1.30
2	A	501	DHB	O1-C	2.13	1.29	1.22
4	C	501	THG	C4'-N10	2.10	1.44	1.38
2	A	502	DHB	O1-C	2.09	1.29	1.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	501	THG	O4-C4	-2.07	1.19	1.24
4	B	501	THG	O4-C4	-2.04	1.19	1.24
4	B	501	THG	C2-N3	2.04	1.39	1.35
4	B	501	THG	O11-C11	-2.03	1.19	1.23
4	C	501	THG	O11-C11	-2.02	1.19	1.23
2	A	502	DHB	O3-C3	2.01	1.40	1.36

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	501	THG	C1'-C11-N	10.73	137.63	117.06
4	B	501	THG	CA-N-C11	10.21	146.58	121.60
4	B	501	THG	C8A-C4A-C4	7.13	120.90	114.57
4	C	501	THG	C8A-C4A-C4	7.07	120.85	114.57
4	B	501	THG	O11-C11-N	-6.13	111.17	122.45
4	B	501	THG	O11-C11-C1'	-5.48	111.17	120.94
4	B	501	THG	C4-C4A-N5	4.28	122.71	119.12
4	C	501	THG	C4-C4A-N5	4.15	122.61	119.12
4	B	501	THG	CG-CB-CA	-3.82	106.01	113.16
4	C	501	THG	C6-C7-N8	3.60	117.44	110.68
4	B	501	THG	CB-CA-N	3.56	118.07	110.88
4	B	501	THG	C-CA-N	-3.47	102.34	110.55
4	B	501	THG	C4A-N5-C6	-2.88	113.91	121.48
2	C	502	DHB	O2-C-C1	2.71	121.88	114.85
4	C	501	THG	N1-C2-N3	-2.55	121.41	125.42
4	C	501	THG	N2-C2-N3	2.53	121.19	117.25
4	B	501	THG	N1-C2-N3	-2.47	121.55	125.42
4	B	501	THG	OE2-CD-CG	2.41	121.78	114.03
4	B	501	THG	C6'-C1'-C11	-2.40	112.84	120.62
4	B	501	THG	C6-C7-N8	2.33	115.05	110.68
4	B	501	THG	C7-C6-C9	-2.32	108.28	112.64
2	A	501	DHB	C6-C5-C4	-2.30	118.14	120.50
4	C	501	THG	C4A-N5-C6	-2.29	115.45	121.48
4	B	501	THG	C2-N1-C8A	2.27	119.63	114.54
4	C	501	THG	C2-N1-C8A	2.25	119.58	114.54
4	B	501	THG	C2'-C1'-C11	2.24	127.86	120.62
4	C	501	THG	C2-N3-C4	2.22	119.46	115.93
4	B	501	THG	C2-N3-C4	2.21	119.44	115.93
4	B	501	THG	OE2-CD-OE1	-2.17	117.90	123.30
4	B	501	THG	OX2-C-OX1	-2.04	119.46	124.09

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	503	TRS	C2-C-C1-O1
3	A	503	TRS	C3-C-C1-O1
3	A	504	TRS	C1-C-C3-O3
3	A	504	TRS	C2-C-C3-O3
3	A	504	TRS	N-C-C3-O3
3	C	503	TRS	N-C-C3-O3
4	B	501	THG	C7-C6-C9-N10
4	B	501	THG	N5-C6-C9-N10
4	B	501	THG	C1'-C11-N-CA
4	B	501	THG	O11-C11-N-CA
4	B	501	THG	CB-CA-N-C11
4	C	501	THG	N-CA-CB-CG
4	C	501	THG	C-CA-CB-CG
3	C	503	TRS	C2-C-C3-O3
4	B	501	THG	OX2-C-CA-N
4	B	501	THG	CA-CB-CG-CD
3	A	503	TRS	N-C-C1-O1
4	C	501	THG	OX2-C-CA-CB
4	B	501	THG	C5'-C4'-N10-C9
4	B	501	THG	OX1-C-CA-N
4	C	501	THG	OX1-C-CA-CB
4	B	501	THG	OE1-CD-CG-CB
4	B	501	THG	C3'-C4'-N10-C9
4	B	501	THG	OE2-CD-CG-CB
3	C	503	TRS	C1-C-C3-O3
4	C	501	THG	OE2-CD-CG-CB
4	C	501	THG	OE1-CD-CG-CB


There are no ring outliers.


6 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	502	DHB	1	0
3	A	504	TRS	2	0
4	B	501	THG	5	0
2	B	502	DHB	2	0
2	A	502	DHB	1	0
4	C	501	THG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	456/474 (96%)	-0.04	2 (0%) 92 93	19, 27, 39, 60	0
1	B	447/474 (94%)	0.06	10 (2%) 62 64	22, 32, 53, 64	0
1	C	455/474 (95%)	0.80	43 (9%) 8 9	27, 47, 68, 73	0
All	All	1358/1422 (95%)	0.28	55 (4%) 37 40	19, 33, 62, 73	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	209	ALA	7.2
1	C	210	GLY	6.1
1	C	456	ALA	5.0
1	C	276	TRP	4.8
1	C	3	ALA	4.7
1	B	453	SER	4.4
1	C	327	ALA	4.1
1	C	267	GLY	4.1
1	C	14	GLY	4.0
1	C	315	PHE	3.8
1	C	133	ALA	3.7
1	C	186	LEU	3.7
1	C	279	ALA	3.6
1	C	5	THR	3.6
1	B	206	HIS	3.4
1	C	131	TYR	3.3
1	B	188	PHE	3.3
1	C	4	PRO	3.2
1	C	13	ALA	3.0
1	A	2	SER	2.9
1	C	273	TYR	2.9
1	B	182	THR	2.9
1	C	12	ALA	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	211	ALA	2.8
1	C	386	ASN	2.7
1	C	292	PHE	2.7
1	C	99	PRO	2.7
1	C	312	PHE	2.7
1	C	100	TYR	2.6
1	B	174	VAL	2.6
1	B	185	LYS	2.6
1	C	384	ALA	2.6
1	C	272	ALA	2.6
1	C	135	THR	2.6
1	C	31	TYR	2.5
1	B	205	ARG	2.5
1	B	212	PRO	2.5
1	C	328	ILE	2.5
1	C	137	GLY	2.5
1	C	289	GLY	2.5
1	C	455	THR	2.5
1	C	246	ALA	2.4
1	C	457	ARG	2.4
1	C	367	LEU	2.4
1	C	11	LEU	2.3
1	C	331	ALA	2.3
1	B	178	LEU	2.2
1	C	325	LEU	2.2
1	C	85	GLY	2.2
1	C	29	TYR	2.2
1	C	302	VAL	2.2
1	B	211	ALA	2.1
1	C	323	ASP	2.0
1	A	114	GLU	2.0
1	C	263	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DHB	A	502	11/11	0.69	0.25	42,51,62,66	0
4	THG	C	501	32/32	0.71	0.27	52,62,82,84	0
4	THG	B	501	32/32	0.81	0.35	37,46,63,67	0
5	EDO	B	504	4/4	0.84	0.15	38,42,43,44	0
3	TRS	A	504	8/8	0.88	0.23	27,36,40,42	0
3	TRS	A	503	8/8	0.88	0.17	23,30,36,39	0
3	TRS	C	503	8/8	0.89	0.16	41,47,49,50	0
3	TRS	B	503	8/8	0.93	0.10	34,37,41,44	0
2	DHB	C	502	11/11	0.94	0.17	36,40,43,44	0
2	DHB	A	501	11/11	0.97	0.12	16,19,23,24	0
2	DHB	B	502	11/11	0.98	0.10	26,29,32,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.