

Full wwPDB X-ray Structure Validation Report (i)

Nov 3, 2024 – 01:34 PM JST

PDB ID	:	5Y2L
Title	:	Crystal structure of a group 2 HA binding antibody AF4H1K1 Fab in complex
		with the 1968 H3N2 pandemic $(H3-AC/68)$ hemagglutinin
Authors	:	Xiao, H.; Qi, J.; Gao, F.G.
Deposited on	:	2017-07-26
Resolution	:	2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	164625	2335 (2.90-2.90)
Clashscore	180529	2564 (2.90-2.90)
Ramachandran outliers	177936	2514 (2.90-2.90)
Sidechain outliers	177891	2516 (2.90-2.90)
RSRZ outliers	164620	2337 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	329	% • 80%	16% ••
2	D	176	2%	
	D	170	<u> </u>	24% •• 7%
3	Ι	233	73%	21% • •
4	J	220	% 73%	19% 5% ·
5	С	2	50%	50%
5	D	2	50%	50%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 7189 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	А	317	Total 2446	C 1532	N 429	0 472	S 13	0	0	0

• Molecule 2 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	В	163	Total 1335	C 823	N 236	0 270	${ m S}{ m 6}$	0	0	0

• Molecule 3 is a protein called a group 2 HA binding antibody AF4H1K1 Fab heavy chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
3	Ι	224	Total 1694	C 1077	N 286	O 325	S 6	0	0	0

• Molecule 4 is a protein called a group 2 HA binding antibody AF4H1K1 Fab light chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
4	J	212	Total	С	Ν	Ο	S	0	0	0
-	, v		1630	1018	278	329	5		0	Ŭ

• Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	С	2	Total 28	C 16	N 2	O 10	0	0	0



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	D	2	Total 28	C 16	N 2	O 10	0	0	0

• Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	А	1	Total 14	C 8	N 1	O 5	0	0
6	А	1	Total 14	C 8	N 1	O 5	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Hemagglutinin



D23 124 124 125 126 127 128 129 553 554 555 555 556 557 553 554 555 554 555 556 557 558 559 559 550 551 551 551 552 553

• Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:	50%	50%	
NAG2 NAG2			
• Molecule 5: 2 opyranose	-acetamido-2-deoxy-beta-I	D-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain D:	50%	50%	I.

NAG1 NAG2



4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants	157.48Å 157.48Å 355.34Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Bosolution(A)	47.33 - 2.90	Depositor
Resolution (A)	47.33 - 2.90	EDS
% Data completeness	99.6 (47.33-2.90)	Depositor
(in resolution range)	99.8 (47.33-2.90)	EDS
R_{merge}	0.10	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.11 (at 2.91 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
B B.	0.212 , 0.244	Depositor
II, II, <i>free</i>	0.215 , 0.247	DCC
R_{free} test set	1887 reflections (4.99%)	wwPDB-VP
Wilson B-factor $(Å^2)$	74.1	Xtriage
Anisotropy	0.710	Xtriage
Bulk solvent $k_{sol}(e/A^3)$, $B_{sol}(A^2)$	0.33 , 50.3	EDS
L-test for $twinning^2$	$ < L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7189	wwPDB-VP
Average B, all atoms $(Å^2)$	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles	
Moi Chain		RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.37	0/2502	0.53	0/3410
2	В	0.51	0/1357	0.69	2/1825~(0.1%)
3	Ι	0.40	0/1739	0.55	0/2372
4	J	0.41	0/1665	0.55	0/2258
All	All	0.41	0/7263	0.57	2/9865~(0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	11	GLU	N-CA-CB	-12.34	88.39	110.60
2	В	42	GLN	CA-CB-CG	5.99	126.59	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2446	0	2393	32	0
2	В	1335	0	1246	50	1
3	Ι	1694	0	1644	36	0
4	J	1630	0	1581	34	0
5	С	28	0	25	2	0



• • • • • •	J					
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	28	0	25	0	0
6	А	28	0	26	1	0
All	All	7189	0	6940	137	1

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (137) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:B:58:LYS:HG2	2:B:59:THR:H	1.29	0.94
3:I:58:GLN:HG3	3:I:64:LEU:HD23	1.54	0.90
2:B:42:GLN:HA	2:B:42:GLN:HE21	1.36	0.88
4:J:51:VAL:HG23	4:J:52:SER:H	1.37	0.88
1:A:18:HIS:HD2	2:B:21:TRP:HA	1.37	0.88
2:B:168:ASN:OD1	2:B:169:ASN:ND2	2.07	0.87
4:J:26:MET:HE3	4:J:113:GLN:HB3	1.59	0.84
4:J:52:SER:O	4:J:53:SER:OG	1.94	0.84
2:B:39:LYS:HD2	2:B:39:LYS:H	1.40	0.84
1:A:77:ASP:OD2	1:A:141:ARG:NH1	2.10	0.83
2:B:28:ASN:ND2	2:B:146:ASN:OD1	2.16	0.79
1:A:185:PRO:HB3	1:A:190:GLU:HG2	1.68	0.76
3:I:229:ILE:HG12	3:I:244:ARG:HG3	1.67	0.75
4:J:211:HIS:O	4:J:233:ARG:NH2	2.20	0.74
4:J:51:VAL:HG23	4:J:52:SER:N	2.03	0.73
2:B:58:LYS:CG	2:B:59:THR:H	2.02	0.71
1:A:285:ASN:HD22	6:A:606:NAG:H83	1.56	0.70
2:B:57:GLU:O	2:B:58:LYS:O	2.10	0.68
4:J:60:GLN:NE2	4:J:68:ARG:HE	1.91	0.67
1:A:18:HIS:CD2	2:B:21:TRP:HA	2.26	0.65
3:I:30:VAL:HG21	3:I:181:PRO:HG3	1.78	0.65
4:J:25:VAL:H	4:J:48:SER:HB3	1.61	0.65
2:B:42:GLN:HA	2:B:42:GLN:NE2	2.10	0.65
1:A:129:GLY:HA3	1:A:162:PRO:HG3	1.79	0.65
2:B:119:PHE:HE2	2:B:123:ARG:HH21	1.44	0.64
2:B:12:ASN:N	2:B:12:ASN:OD1	2.31	0.63
2:B:42:GLN:NE2	3:I:129:VAL:HG13	2.14	0.63
4:J:54:SER:O	4:J:56:LEU:N	2.32	0.63
3:I:75:GLY:HA2	5:C:2:NAG:H82	1.81	0.62
1:A:74:PRO:HA	1:A:141:ARG:HH12	1.64	0.62
2:B:56:ILE:O	2:B:58:LYS:N	2.33	0.62



			Clash	
Atom-1	Atom-2	distance (\AA)	overlap (Å)	
3:I:160:PRO:HG3	3:I:172:LEU:HB3	1.83	0.61	
2:B:58:LYS:HG2	2:B:59:THR:N	2.10	0.60	
2:B:39:LYS:HD2	2:B:39:LYS:N	2.15	0.60	
2:B:126:LEU:O	2:B:127:ARG:C	2.40	0.60	
1:A:9:SER:N	2:B:143:LYS:HZ3	1.98	0.60	
3:I:58:GLN:OE1	4:J:61:GLN:NE2	2.27	0.60	
1:A:53:ASN:HD21	1:A:276:THR:HG23	1.66	0.60	
3:I:25:GLU:OE1	3:I:139:GLN:N	2.35	0.59	
4:J:53:SER:O	4:J:54:SER:HB2	2.03	0.59	
1:A:84:TRP:CE2	1:A:116:GLY:HA2	2.37	0.59	
1:A:89:GLU:HG3	1:A:267:ILE:HD11	1.85	0.58	
3:I:202:ALA:HB2	3:I:212:LEU:HD23	1.85	0.58	
1:A:323:VAL:CG2	2:B:12:ASN:HA	2.35	0.57	
2:B:168:ASN:OD1	2:B:169:ASN:N	2.37	0.57	
4:J:118:PHE:N	4:J:118:PHE:CD2	2.73	0.56	
2:B:128:GLU:HB3	2:B:170:ARG:HH12	1.71	0.56	
3:I:25:GLU:OE1	3:I:138:GLY:HA3	2.06	0.56	
2:B:61:GLU:O	2:B:62:LYS:HG2	2.05	0.55	
2:B:124:ARG:HH11	2:B:124:ARG:HG3	1.70	0.55	
2:B:10:ILE:HD13	2:B:10:ILE:N	2.21	0.55	
3:I:169:THR:HA	3:I:220:SER:HB2	1.89	0.55	
1:A:102:VAL:HG22	1:A:232:ILE:HB	1.89	0.55	
4:J:51:VAL:CG2	4:J:52:SER:H	2.13	0.54	
4:J:169:GLN:HG2	4:J:176:LEU:HD11	1.88	0.54	
3:I:49:SER:O	3:I:72:PHE:HB2	2.08	0.54	
4:J:60:GLN:HB2	4:J:70:LEU:HD11	1.89	0.54	
2:B:11:GLU:OE2	2:B:11:GLU:HA	2.08	0.54	
3:I:184:VAL:HG12	3:I:234:HIS:HB2	1.90	0.54	
3:I:119:PRO:HD3	3:I:135:ASP:HB2	1.89	0.53	
1:A:307:LYS:HE2	2:B:92:TRP:CZ2	2.43	0.53	
2:B:128:GLU:O	2:B:170:ARG:NH1	2.40	0.53	
4:J:168:VAL:HG12	4:J:218:VAL:HG22	1.91	0.52	
1:A:182:ILE:HD11	1:A:215:PRO:HD3	1.92	0.52	
1:A:220:ARG:O	1:A:227:SER:HB2	2.09	0.52	
3:I:91:ARG:HG2	3:I:92:ASP:N	2.25	0.52	
4:J:51:VAL:O	4:J:52:SER:HB2	2.10	0.51	
1:A:97:CYS:O	1:A:224:ARG:NH1	2.44	0.51	
1:A:20:VAL:HG11	5:C:1:NAG:C8	2.41	0.50	
1:A:42:LEU:HD12	2:B:100:VAL:HG12	1.92	0.50	
1:A:314:LEU:HB3	2:B:100:VAL:HG11	1.93	0.49	
1:A:222:TRP:CZ2	1:A:225:GLY:HA2	2.47	0.49	



	loue page	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:230:ILE:HD12	1:A:252:ILE:HG13	1.93	0.49
3:I:73:ASP:HB3	3:I:75:GLY:H	1.75	0.49
2:B:165:GLU:O	2:B:168:ASN:OD1	2.29	0.49
2:B:113:SER:O	2:B:117:LYS:HG2	2.12	0.49
1:A:266:SER:OG	1:A:267:ILE:N	2.46	0.49
4:J:147:LEU:O	4:J:205:LYS:HD2	2.13	0.49
2:B:10:ILE:HG22	2:B:11:GLU:N	2.28	0.48
1:A:290:ASN:OD1	3:I:47:THR:OG1	2.29	0.48
3:I:231:ASN:ND2	3:I:242:ASP:OD1	2.38	0.48
1:A:204:VAL:HG13	1:A:243:LEU:HD11	1.96	0.48
4:J:158:LEU:HB2	4:J:197:LEU:HB3	1.96	0.48
2:B:61:GLU:C	2:B:62:LYS:HG2	2.34	0.48
2:B:170:ARG:HB3	2:B:171:PHE:CD1	2.48	0.48
1:A:301:THR:HG23	1:A:305:CYS:SG	2.54	0.47
4:J:173:ASP:OD2	4:J:211:HIS:HB3	2.14	0.47
3:I:21:VAL:HA	3:I:45:GLY:HA3	1.97	0.47
2:B:14:TRP:HB3	2:B:25:ARG:NH2	2.30	0.47
4:J:60:GLN:HE22	4:J:68:ARG:HE	1.61	0.46
2:B:10:ILE:HG22	2:B:11:GLU:H	1.80	0.46
3:I:43:ALA:O	3:I:96:ASN:ND2	2.49	0.46
3:I:173:GLY:HA2	3:I:188:TRP:CH2	2.51	0.46
3:I:159:ALA:HA	3:I:160:PRO:HD3	1.76	0.45
1:A:141:ARG:NH2	1:A:147:PHE:O	2.49	0.45
2:B:128:GLU:HB3	2:B:170:ARG:NH1	2.32	0.45
4:J:130:ARG:HD2	4:J:192:ASP:O	2.16	0.45
3:I:126:PHE:HB2	3:I:129:VAL:HG23	1.99	0.45
1:A:127:TRP:CZ2	1:A:253:ALA:HB1	2.52	0.44
2:B:45:ILE:HG21	3:I:129:VAL:HG11	2.00	0.44
4:J:104:GLU:H	4:J:104:GLU:HG3	1.60	0.44
4:J:142:PRO:HD3	4:J:154:VAL:HG22	1.99	0.44
4:J:28:GLN:O	4:J:122:GLN:NE2	2.49	0.44
4:J:208:TYR:O	4:J:214:TYR:OH	2.35	0.44
2:B:124:ARG:HH11	2:B:124:ARG:CG	2.31	0.44
3:I:132:GLY:HA2	4:J:114:TYR:CD1	2.53	0.44
4:J:156:CYS:HB2	4:J:170:TRP:CH2	2.52	0.44
2:B:53:ASN:OD1	3:I:122:LYS:HG3	2.17	0.44
3:I:92:ASP:O	3:I:93:ILE:C	2.55	0.44
2:B:131:GLU:OE2	2:B:170:ARG:HG2	2.18	0.44
2:B:170:ARG:HB3	2:B:171:PHE:HD1	1.83	0.43
1:A:291:ASP:OD1	1:A:292:LYS:HG3	2.17	0.43
4:J:147:LEU:H	4:J:147:LEU:HD12	1.82	0.43



A 1 1		Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
2:B:141:TYR:CG	2:B:170:ARG:HG3	2.54	0.43
3:I:73:ASP:HB3	3:I:75:GLY:N	2.34	0.43
3:I:171:ALA:HB2	3:I:217:THR:HG22	2.00	0.43
1:A:62:ILE:HG22	1:A:63:ASP:N	2.34	0.43
2:B:51:LYS:HE3	2:B:107:THR:OG1	2.19	0.42
2:B:162:TYR:O	2:B:166:ALA:N	2.38	0.42
3:I:137:TRP:CZ3	4:J:67:PRO:HG2	2.55	0.42
4:J:167:LYS:HB3	4:J:219:THR:HG23	2.02	0.42
3:I:110:THR:HG23	3:I:144:THR:HA	2.01	0.42
3:I:54:HIS:CD2	3:I:118:ASP:HB2	2.55	0.42
1:A:270:SER:HB2	1:A:284:PRO:HA	2.02	0.41
2:B:168:ASN:CG	2:B:169:ASN:N	2.73	0.41
4:J:39:GLU:O	4:J:101:LEU:HB2	2.20	0.41
1:A:176:LYS:HE2	1:A:257:TYR:CE1	2.56	0.41
3:I:246:GLU:O	3:I:247:PRO:O	2.38	0.41
4:J:140:PHE:HA	4:J:141:PRO:HD2	1.91	0.41
2:B:23:GLY:HA3	2:B:36:ALA:HA	2.03	0.41
2:B:149:ILE:HD12	2:B:150:GLU:N	2.36	0.41
3:I:234:HIS:CD2	3:I:236:PRO:HD2	2.56	0.41
4:J:127:GLU:OE2	4:J:195:TYR:OH	2.30	0.41
3:I:129:VAL:HG12	3:I:130:SER:N	2.36	0.40
2:B:154:ASN:O	2:B:156:THR:HG23	2.21	0.40
2:B:27:GLN:HG3	2:B:32:THR:HG22	2.03	0.40
3:I:200:PHE:CE2	4:J:198:SER:HB2	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:62:LYS:NZ	2:B:86:ASP:OD2[3_655]	1.47	0.73

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	315/329~(96%)	295~(94%)	18~(6%)	2(1%)	22 52
2	В	161/176~(92%)	145~(90%)	13~(8%)	3~(2%)	6 24
3	Ι	220/233~(94%)	200~(91%)	19 (9%)	1 (0%)	25 56
4	J	210/220~(96%)	193~(92%)	13~(6%)	4(2%)	6 24
All	All	906/958~(95%)	833 (92%)	63 (7%)	10 (1%)	12 37

analysed, and the total number of residues.

All (10) Ramachandran outliers are listed below:

Mol	Chain	\mathbf{Res}	Type
2	В	57	GLU
2	В	58	LYS
2	В	127	ARG
3	Ι	247	PRO
4	J	51	VAL
4	J	52	SER
4	J	53	SER
4	J	55	TYR
1	А	62	ILE
1	А	324	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	А	279/290~(96%)	273~(98%)	6(2%)	47	78
2	В	142/150~(95%)	136~(96%)	6 (4%)	25	59
3	Ι	187/196~(95%)	180~(96%)	7~(4%)	29	64
4	J	184/192~(96%)	172 (94%)	12~(6%)	14	40
All	All	792/828~(96%)	761 (96%)	31 (4%)	27	62

All (31) residues with a non-rotameric sidechain are listed below:



WIOI	Unam	Res	Type
1	А	20	VAL
1	А	29	ILE
1	А	101	ASP
1	А	186	SER
1	А	212	THR
1	А	301	THR
2	В	11	GLU
2	В	12	ASN
2	В	42	GLN
2	В	59	THR
2	В	72	GLU
2	В	168	ASN
3	Ι	21	VAL
3	Ι	25	GLU
3	Ι	47	THR
3	Ι	48	SER
3	Ι	58	GLN
3	Ι	84	LYS
3	Ι	95	ARG
4	J	25	VAL
4	J	48	SER
4	J	50	SER
4	J	56	LEU
4	J	68	ARG
4	J	92	THR
4	J	104	GLU
4	J	118	PHE
4	J	129	LYS
4	J	147	LEU
4	J	216	CYS
4	J	219	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	18	HIS
1	А	216	ASN
2	В	42	GLN
2	В	60	ASN
2	В	135	ASN
2	В	169	ASN
4	J	60	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	I Type Chain Pos Link		Bond lengths			Bond angles				
	туре	Unain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	С	1	5,1	$14,\!14,\!15$	0.34	0	$17,\!19,\!21$	0.63	0
5	NAG	C	2	5	14,14,15	1.03	1 (7%)	$17,\!19,\!21$	0.88	1 (5%)
5	NAG	D	1	5,1	14,14,15	0.54	0	17,19,21	0.94	1 (5%)
5	NAG	D	2	5	14,14,15	0.78	0	17,19,21	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	С	1	5,1	-	0/6/23/26	0/1/1/1
5	NAG	С	2	5	-	1/6/23/26	0/1/1/1
5	NAG	D	1	5,1	-	2/6/23/26	0/1/1/1
5	NAG	D	2	5	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
5	С	2	NAG	O5-C1	3.45	1.49	1.43



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	D	1	NAG	C1-O5-C5	2.82	116.01	112.19
5	С	2	NAG	C1-O5-C5	2.65	115.78	112.19

All (2) bond angle outliers are listed below:

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	1	NAG	O5-C5-C6-O6
5	D	1	NAG	C4-C5-C6-O6
5	D	2	NAG	C8-C7-N2-C2
5	D	2	NAG	O7-C7-N2-C2
5	С	2	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	С	2	NAG	1	0
5	С	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal Trung Chain D		Dec	Tink	Bo	Bond lengths			Bond angles		
IVIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	А	606	1	14,14,15	0.63	0	17,19,21	0.52	0
6	NAG	А	603	1	14,14,15	0.30	0	17,19,21	0.57	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	А	606	1	-	4/6/23/26	0/1/1/1
6	NAG	А	603	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	А	606	NAG	O5-C5-C6-O6
6	А	603	NAG	O5-C5-C6-O6
6	А	606	NAG	C4-C5-C6-O6
6	А	603	NAG	C4-C5-C6-O6
6	А	606	NAG	C8-C7-N2-C2
6	А	606	NAG	O7-C7-N2-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	А	606	NAG	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	317/329~(96%)	-0.33	3 (0%) 81 76	55, 80, 108, 165	0
2	В	163/176~(92%)	-0.14	4 (2%) 58 52	58, 92, 137, 189	0
3	Ι	224/233~(96%)	-0.22	3 (1%) 74 69	58, 82, 109, 153	0
4	J	212/220~(96%)	-0.18	3 (1%) 73 68	60, 83, 116, 149	0
All	All	916/958~(95%)	-0.23	13 (1%) 73 68	55, 83, 120, 189	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	В	10	ILE	5.7
3	Ι	130	SER	3.5
1	А	9	SER	3.1
3	Ι	169	THR	2.9
2	В	134	GLY	2.6
2	В	132	GLU	2.5
4	J	234	GLY	2.5
2	В	139	LYS	2.3
1	А	11	ALA	2.2
1	А	324	PRO	2.2
3	Ι	170	ALA	2.1
4	J	51	VAL	2.1
4	J	174	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

SUGAR-RSR INFOmissingINFO

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
6	NAG	А	606	14/15	0.68	0.14	108,111,118,119	0
6	NAG	А	603	14/15	0.80	0.09	105,110,116,125	0

6.5 Other polymers (i)

There are no such residues in this entry.

