



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 13, 2023 – 11:01 PM JST

PDB ID : 5YD4  
Title : Crystal structure of the scFv antibody 4B08 with epitope peptide (mutation T6A)  
Authors : Caaveiro, J.M.M.; Miyanabe, K.; Tsumoto, K.  
Deposited on : 2017-09-11  
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1509 (1.38-1.34)
Clashscore	141614	1551 (1.38-1.34)
Ramachandran outliers	138981	1530 (1.38-1.34)
Sidechain outliers	138945	1530 (1.38-1.34)
RSRZ outliers	127900	1487 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	
1	C	251	
1	E	251	
1	G	251	
2	B	9	
2	D	9	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	9	 33% 100%
2	H	9	 56% 67% 11% 22%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8594 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called scFv 4B08.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	232	1874	1188	314	362	10	0	14	0
1	C	229	1856	1180	309	357	10	0	12	0
1	E	230	1824	1158	305	352	9	0	7	0
1	G	226	1776	1125	298	343	10	0	7	0

- Molecule 2 is a protein called Epitope peptide (mutation T6A).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	8	68	44	9	15	0	0	0
2	D	8	68	44	9	15	0	0	0
2	F	9	76	48	10	18	0	0	0
2	H	7	60	38	8	14	0	1	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	1
3	C	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	233	Total O 242 242	0	12
4	B	12	Total O 12 12	0	0
4	C	247	Total O 256 256	0	12
4	D	13	Total O 16 16	0	3

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	E	257	Total 268	O 268	0	13
4	F	19	Total 20	O 20	0	1
4	G	137	Total 138	O 138	0	3



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: scFv 4B08





- Molecule 1: scFv 4B08





- Molecule 1: scFv 4B08




- Molecule 1: scFv 4B08




- Molecule 2: Epitope peptide (mutation T6A)




- Molecule 2: Epitope peptide (mutation T6A)



- Molecule 2: Epitope peptide (mutation T6A)



- Molecule 2: Epitope peptide (mutation T6A)





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.47Å 72.90Å 86.46Å 90.00° 114.15° 90.00°	Depositor
Resolution (Å)	33.10 – 1.35 33.09 – 1.35	Depositor EDS
% Data completeness (in resolution range)	99.1 (33.10-1.35) 99.1 (33.09-1.35)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.44 (at 1.35Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.135 , 0.173 0.141 , 0.178	Depositor DCC
$R_{free}$ test set	3797 reflections (1.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	10.8	Xtrriage
Anisotropy	0.168	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 59.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8594	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.7710e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	0/1958	0.85	1/2649 (0.0%)
1	C	0.78	0/1936	0.91	3/2622 (0.1%)
1	E	0.70	0/1887	0.85	5/2557 (0.2%)
1	G	0.81	3/1833 (0.2%)	0.92	5/2486 (0.2%)
2	B	0.67	0/70	0.66	0/94
2	D	0.77	0/70	0.67	0/94
2	F	0.65	0/78	0.72	0/105
2	H	0.78	0/62	0.73	0/83
All	All	0.74	3/7894 (0.0%)	0.88	14/10690 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	233	CYS	CB-SG	-8.56	1.67	1.82
1	G	233	CYS	CA-CB	5.54	1.66	1.53
1	G	162	SER	CB-OG	-5.16	1.35	1.42

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	144	MET	CG-SD-CE	9.20	114.91	100.20
1	E	195	ARG	NE-CZ-NH1	-7.34	116.63	120.30
1	A	52	ASP	CB-CG-OD2	-6.81	112.17	118.30
1	E	195	ARG	NE-CZ-NH2	6.49	123.55	120.30
1	C	195	ARG	NE-CZ-NH1	-5.66	117.47	120.30
1	C	141	ASP	CB-CG-OD2	-5.51	113.34	118.30
1	E	184	ARG	NE-CZ-NH1	5.51	123.05	120.30
1	E	222	ARG	NE-CZ-NH2	-5.39	117.61	120.30
1	E	52	ASP	CB-CG-OD2	-5.22	113.60	118.30
1	G	52	ASP	CB-CG-OD2	-5.20	113.62	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	52	ASP	CB-CG-OD1	5.18	122.97	118.30
1	G	105	ASN	CB-CA-C	-5.17	100.07	110.40
1	G	227	ASP	CB-CG-OD1	5.09	122.88	118.30
1	C	205	ASP	CB-CG-OD1	-5.08	113.72	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1874	0	1835	6	0
1	C	1856	0	1807	7	0
1	E	1824	0	1776	3	0
1	G	1776	0	1713	20	0
2	B	68	0	57	0	0
2	D	68	0	57	0	0
2	F	76	0	64	0	0
2	H	60	0	46	3	0
3	A	10	0	0	0	0
3	C	15	0	0	0	0
3	E	15	0	0	0	0
4	A	242	0	0	0	0
4	B	12	0	0	0	0
4	C	256	0	0	2	0
4	D	16	0	0	0	0
4	E	268	0	0	2	0
4	F	20	0	0	0	0
4	G	138	0	0	2	0
All	All	8594	0	7355	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:104:THR:HG21	2:H:9[A]:PRO:HG3	1.55	0.87
1:G:144:MET:CE	1:G:169:LEU:HD11	2.09	0.82
1:G:104:THR:CG2	2:H:9[A]:PRO:HG3	2.17	0.74
1:G:144:MET:HE2	1:G:169:LEU:HD11	1.71	0.72
1:G:144:MET:HE1	1:G:169:LEU:HD11	1.77	0.67
1:G:144:MET:HE2	1:G:235:GLN:HB3	1.77	0.67
1:G:104:THR:HG21	2:H:9[A]:PRO:CG	2.24	0.66
1:G:175:ASN:ND2	4:G:301:HOH:O	2.28	0.66
1:G:235:GLN:HE22	1:G:238:GLU:H	1.48	0.62
1:G:175:ASN:OD1	1:G:195[B]:ARG:NE	2.34	0.60
1:G:196[B]:MET:SD	1:G:197:SER:HB3	2.49	0.53
1:A:93[A]:SER:HB3	1:G:222:ARG:HH22	1.75	0.52
1:C:251:LEU:HB3	4:C:435[A]:HOH:O	2.09	0.50
1:G:235:GLN:HE21	1:G:237:LEU:H	1.60	0.50
1:C:158:SER:HB3	1:C:221:SER:O	2.12	0.50
1:E:219[B]:ARG:HG3	4:E:457[B]:HOH:O	2.10	0.49
1:G:235:GLN:NE2	1:G:237:LEU:H	2.11	0.49
1:A:219[B]:ARG:NH1	1:A:221[B]:SER:OG	2.47	0.48
1:G:5:GLN:NE2	1:G:7:GLN:OE1	2.47	0.47
1:C:42[B]:ARG:HD2	4:C:453:HOH:O	2.15	0.46
1:C:5[B]:GLN:HE21	1:C:7:GLN:HE21	1.65	0.45
1:A:251:LEU:HD12	1:G:198:ASN:HD21	1.81	0.45
1:C:5[A]:GLN:HB2	1:C:27:ALA:HB3	1.99	0.44
1:G:176:THR:O	1:G:195[B]:ARG:HA	2.15	0.44
1:A:14:VAL:HG21	1:A:88:LEU:HD13	2.00	0.44
1:E:223:VAL:HG12	1:E:251:LEU:HD21	2.00	0.43
1:A:11:ALA:HB3	1:G:157:GLU:HG2	2.00	0.43
1:C:234[B]:MET:HB2	1:C:243:PHE:CD2	2.54	0.42
1:A:15:ARG:HH11	1:A:118:SER:CB	2.32	0.41
1:E:42:ARG:HD2	4:E:470:HOH:O	2.21	0.41
1:C:42[B]:ARG:HD2	1:C:42[B]:ARG:HH11	1.71	0.40
1:G:164:ARG:NH2	4:G:306:HOH:O	2.50	0.40
1:G:95:ILE:HD13	1:G:114:SER:HA	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	242/251 (96%)	240 (99%)	2 (1%)	0	100	100
1	C	237/251 (94%)	235 (99%)	2 (1%)	0	100	100
1	E	233/251 (93%)	230 (99%)	3 (1%)	0	100	100
1	G	226/251 (90%)	219 (97%)	7 (3%)	0	100	100
2	B	6/9 (67%)	6 (100%)	0	0	100	100
2	D	6/9 (67%)	6 (100%)	0	0	100	100
2	F	7/9 (78%)	7 (100%)	0	0	100	100
2	H	5/9 (56%)	5 (100%)	0	0	100	100
All	All	962/1040 (92%)	948 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/199 (105%)	208 (100%)	1 (0%)	88	74
1	C	206/199 (104%)	206 (100%)	0	100	100
1	E	202/199 (102%)	202 (100%)	0	100	100
1	G	195/199 (98%)	193 (99%)	2 (1%)	76	49
2	B	7/8 (88%)	7 (100%)	0	100	100
2	D	7/8 (88%)	7 (100%)	0	100	100
2	F	8/8 (100%)	8 (100%)	0	100	100
2	H	6/8 (75%)	6 (100%)	0	100	100
All	All	840/828 (101%)	837 (100%)	3 (0%)	91	81

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	226	GLU
1	G	175	ASN
1	G	235	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	C	198	ASN
1	G	5	GLN
1	G	7	GLN
1	G	198	ASN
1	G	235	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	302	-	4,4,4	0.29	0	6,6,6	0.84	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	C	301	-	4,4,4	0.32	0	6,6,6	1.18	0
3	SO4	C	303	-	4,4,4	0.34	0	6,6,6	0.34	0
3	SO4	A	301	-	4,4,4	0.38	0	6,6,6	0.35	0
3	SO4	E	303	-	4,4,4	0.49	0	6,6,6	0.45	0
3	SO4	C	302[A]	-	4,4,4	0.34	0	6,6,6	0.43	0
3	SO4	E	301	-	4,4,4	0.59	0	6,6,6	0.64	0
3	SO4	E	302	-	4,4,4	0.42	0	6,6,6	0.57	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	232/251 (92%)	0.40	7 (3%) 50 57	7, 11, 26, 45	0
1	C	229/251 (91%)	0.43	7 (3%) 49 56	7, 11, 26, 49	0
1	E	230/251 (91%)	0.39	4 (1%) 70 74	9, 14, 27, 46	0
1	G	226/251 (90%)	2.00	90 (39%) 0 0	13, 25, 41, 53	3 (1%)
2	B	8/9 (88%)	1.39	3 (37%) 0 0	12, 14, 28, 37	0
2	D	8/9 (88%)	0.97	1 (12%) 3 4	9, 11, 19, 27	0
2	F	9/9 (100%)	2.34	3 (33%) 0 0	13, 16, 28, 53	0
2	H	7/9 (77%)	4.92	5 (71%) 0 0	30, 35, 63, 68	1 (14%)
All	All	949/1040 (91%)	0.85	120 (12%) 3 3	7, 15, 34, 68	4 (0%)

All (120) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	5	TYR	11.4
2	H	4	TYR	9.7
1	G	14	VAL	9.5
1	G	117	VAL	8.9
2	F	2	ILE	8.8
1	G	103	TYR	8.7
1	G	102	TYR	8.6
1	G	193[B]	ILE	8.6
1	G	203	VAL	8.5
1	G	177	TYR	7.9
1	G	213	GLY	7.7
1	G	13	LEU	7.1
1	G	205	ASP	6.3
1	G	171	HIS	5.9
2	B	2	ILE	5.9
1	G	27	ALA	5.9

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	H	9[A]	PRO	5.9
1	G	143	VAL	5.8
2	F	1	ASP	5.7
1	G	194[B]	TYR	5.4
1	G	169	LEU	5.3
1	G	239	TYR	5.2
1	A	253	VAL	5.1
1	G	59	TYR	5.1
1	G	237	LEU	5.0
1	G	141	ASP	5.0
1	C	251	LEU	5.0
2	D	2	ILE	4.7
1	G	214	THR	4.7
1	G	166	SER	4.6
1	G	11	ALA	4.6
1	G	118	SER	4.4
1	G	17	GLY	4.4
1	G	225	ALA	4.4
1	G	172	ARG	4.3
1	G	204	PRO	4.3
1	G	178	LEU	4.3
1	G	201	SER	4.3
1	G	142	ILE	4.3
2	F	4	TYR	4.2
1	G	180	TRP	4.1
1	G	4	VAL	4.0
1	G	202	GLY	4.0
1	G	156	GLY	4.0
1	G	43	PRO	3.9
1	G	28	GLY	3.8
1	G	165	SER	3.8
1	G	241	TYR	3.8
1	G	18	THR	3.8
1	C	252	LYS	3.8
1	G	186	GLY	3.8
1	G	226	GLU	3.8
1	G	170	LEU	3.8
1	G	163	CYS	3.7
1	G	104	THR	3.6
1	G	164	ARG	3.6
2	H	3	ASN	3.5
1	G	64	GLU	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	E	253	VAL	3.5
1	G	65	LYS	3.5
1	G	93	SER	3.4
1	G	216	PHE	3.3
1	G	90	SER	3.3
1	G	224	GLU	3.3
1	G	116	THR	3.3
1	A	252	LYS	3.3
1	G	16	PRO	3.2
1	E	140	SER	3.2
1	G	167	LYS	3.1
1	G	88	LEU	3.1
1	G	211	GLY	3.1
1	G	200	ALA	3.1
1	G	236	HIS	3.0
1	G	68	GLY	3.0
1	G	12	GLU	2.9
2	B	4	TYR	2.8
1	G	179	PHE	2.8
1	A	139	GLY	2.8
1	C	106[A]	TYR	2.7
1	E	103	TYR	2.7
1	G	15	ARG	2.7
1	G	196[A]	MET	2.7
1	G	198	ASN	2.6
1	G	32	THR	2.6
1	G	89	THR	2.6
1	G	145	THR	2.6
1	G	67	LYS	2.6
1	G	238	GLU	2.6
2	B	5	TYR	2.6
1	G	5	GLN	2.6
1	G	223	VAL	2.6
1	G	34	TYR	2.5
1	G	20	VAL	2.5
1	G	176	THR	2.4
1	C	59[A]	TYR	2.4
1	A	59	TYR	2.4
1	G	86	SER	2.4
1	C	221	SER	2.3
1	G	173	ASN	2.3
1	G	155	PRO	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	222	ARG	2.3
1	G	218	LEU	2.2
1	G	233	CYS	2.2
1	G	154	THR	2.2
1	A	118	SER	2.2
1	G	222	ARG	2.2
1	G	168	SER	2.2
1	A	3	GLU	2.2
1	G	240	PRO	2.2
1	G	195[B]	ARG	2.1
1	E	39	VAL	2.1
1	G	63	ASN	2.1
1	C	118	SER	2.1
1	G	207	PHE	2.1
2	H	6	ALA	2.1
1	G	115	VAL	2.0
1	G	174	GLY	2.0
1	G	66	PHE	2.0
1	A	178	LEU	2.0
1	G	206	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	C	302[A]	5/5	0.87	0.18	30,31,44,47	5
3	SO4	C	303	5/5	0.94	0.18	29,30,40,47	5
3	SO4	E	301	5/5	0.95	0.14	16,17,20,21	5

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	C	301	5/5	0.97	0.14	20,23,32,34	0
3	SO4	A	301	5/5	0.97	0.33	43,44,53,57	0
3	SO4	A	302	5/5	0.98	0.16	20,25,32,34	0
3	SO4	E	302	5/5	0.98	0.07	25,25,27,28	5
3	SO4	E	303	5/5	0.98	0.26	24,28,33,40	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.