



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 13, 2023 – 10:59 PM JST

PDB ID : 5YD8  
Title : Crystal structure of human PCNA in complex with APIM of human ZRANB3  
Authors : Hashimoto, H.; Tagata, R.  
Deposited on : 2017-09-12  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	261	 6% (poor fit), 88% (0-1 outliers), 7% (2 outliers), 5% (3+ outliers)
1	Y	261	 6% (poor fit), 82% (0-1 outliers), 10% (2 outliers), 7% (3+ outliers)
1	Z	261	 8% (poor fit), 78% (0-1 outliers), 15% (2 outliers), 5% (3+ outliers)
2	U	11	 18% (poor fit), 82% (0-1 outliers), 18% (2 outliers)
2	V	11	 27% (poor fit), 55% (0-1 outliers), 9% (2 outliers), 36% (3+ outliers)
2	W	11	 18% (poor fit), 45% (0-1 outliers), 18% (2 outliers), 9% (3 outliers), 27% (3+ outliers)

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5933 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proliferating cell nuclear antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	Y	243	Total 1870	C 1178	N 307	O 369	S 16	0	0	0
1	X	248	Total 1904	C 1199	N 313	O 376	S 16	0	0	0
1	Z	247	Total 1899	C 1197	N 312	O 374	S 16	0	0	0

- Molecule 2 is a protein called ZRANB3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	W	8	Total 66	C 43	N 11	O 12	0	0	0
2	U	11	Total 89	C 57	N 16	O 16	0	0	0
2	V	7	Total 60	C 40	N 10	O 10	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	Y	13	Total 13 O 13	0	0
3	X	16	Total 16 O 16	0	0
3	Z	16	Total 16 O 16	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proliferating cell nuclear antigen




- Molecule 1: Proliferating cell nuclear antigen





- Molecule 1: Proliferating cell nuclear antigen



- Molecule 2: ZRANB3



## ● Molecule 2: ZRANB3



## ● Molecule 2: ZRANB3



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.68Å 83.68Å 194.93Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.30 19.99 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.3 (20.00-2.30) 98.5 (19.99-2.30)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.33 (at 2.30Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.219 , 0.265 0.218 , 0.263	Depositor DCC
$R_{free}$ test set	1771 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.8	Xtrriage
Anisotropy	0.549	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 35.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.039 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5933	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	X	0.55	0/1928	0.74	0/2602
1	Y	0.54	0/1892	0.73	0/2549
1	Z	0.55	0/1923	0.74	0/2595
2	U	0.52	0/89	0.80	0/115
2	V	0.44	0/60	0.61	0/80
2	W	0.67	0/66	0.57	0/88
All	All	0.55	0/5958	0.74	0/8029

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	1904	0	1919	11	0
1	Y	1870	0	1887	22	0
1	Z	1899	0	1916	26	0
2	U	89	0	97	2	0
2	V	60	0	63	0	0
2	W	66	0	68	2	0
3	X	16	0	0	1	0
3	Y	13	0	0	1	0
3	Z	16	0	0	0	0
All	All	5933	0	5950	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Y:16:LEU:HD22	1:Y:79:LEU:HD12	1.37	1.06
1:Y:16:LEU:CD2	1:Y:79:LEU:HD12	2.05	0.85
1:Z:71:ASN:OD1	1:Z:73:THR:HG23	1.77	0.84
1:Y:114:TYR:HE2	1:Z:154:ILE:HD11	1.45	0.79
1:X:185:THR:HG22	1:X:195:VAL:H	1.51	0.76
1:Y:104:GLU:HG3	1:Y:105:ALA:H	1.53	0.72
1:Z:56:GLY:HA3	1:Z:244:MET:HG2	1.70	0.71
1:Z:108:GLN:O	1:Z:108:GLN:HG2	1.93	0.68
1:Z:151:LEU:HA	1:Z:154:ILE:HD12	1.75	0.68
1:X:129:PRO:HA	2:U:1079:LYS:HE2	1.76	0.68
2:W:1073:THR:HA	2:W:1076:LEU:HD12	1.76	0.68
1:Z:5:ARG:NH2	1:Z:87:ILE:HD13	2.10	0.67
1:Y:27:CYS:SG	1:Y:67:ALA:HB1	2.36	0.66
1:Y:56:GLY:HA3	1:Y:244:MET:HG2	1.78	0.66
1:Y:16:LEU:HD22	1:Y:79:LEU:CD1	2.22	0.65
1:Y:114:TYR:CE2	1:Z:154:ILE:HD11	2.31	0.65
1:Z:68:MET:HE2	1:Z:118:LEU:HD11	1.77	0.64
1:Z:5:ARG:NH2	1:Z:87:ILE:HG21	2.14	0.62
1:X:185:THR:CG2	1:X:195:VAL:H	2.12	0.61
1:Y:104:GLU:HG3	1:Y:105:ALA:N	2.16	0.59
1:Y:87:ILE:HB	1:Y:104:GLU:HB3	1.86	0.57
1:Z:5:ARG:CZ	1:Z:87:ILE:HG21	2.34	0.57
1:Z:38:GLN:HB3	1:Z:125:GLN:HE22	1.73	0.53
1:X:109:GLU:HG3	1:X:110:LYS:H	1.74	0.52
1:Z:241:ILE:O	1:Z:241:ILE:HG22	2.08	0.52
1:Z:45:VAL:HG22	2:W:1070:SER:HB2	1.94	0.50
1:X:109:GLU:HG3	1:X:110:LYS:N	2.27	0.50
1:Z:6:LEU:HD21	1:Z:12:LEU:HB2	1.94	0.49
1:Y:78:ILE:HD11	1:Z:175:LEU:HD13	1.95	0.49
1:Z:1:MET:HG2	1:Z:61:ARG:CZ	2.43	0.48
1:Y:188:VAL:HG21	1:Y:194:ALA:HB2	1.96	0.48
1:Z:205:LEU:HD21	1:Z:232:ASP:H	1.77	0.47
1:Y:184:GLN:HE22	1:Y:197:ILE:H	1.63	0.47
1:Y:64:ARG:CD	3:Y:308:HOH:O	2.62	0.47
1:Z:124:GLU:HG2	1:Z:125:GLN:N	2.29	0.47
1:Z:10:SER:HA	1:Z:13:LYS:HD3	1.97	0.47
1:Z:2:PHE:HB3	1:Z:92:ALA:HB3	1.97	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:91:ARG:HB3	1:X:100:ALA:HB3	1.97	0.47
1:Y:207:PHE:CZ	1:Y:235:LEU:HB2	2.50	0.46
1:Y:16:LEU:HD23	1:Y:76:SER:HA	1.98	0.45
1:Y:146:ARG:HD2	1:Y:150:ASP:OD2	2.16	0.45
1:Z:207:PHE:CZ	1:Z:235:LEU:HB2	2.52	0.44
1:Y:114:TYR:HE2	1:Z:154:ILE:CD1	2.24	0.43
1:X:185:THR:HG21	1:X:194:ALA:HA	2.01	0.43
1:Z:23:ILE:HG22	1:Z:41:ASP:HA	2.00	0.43
1:Y:12:LEU:HD11	1:Y:90:LEU:HD11	2.00	0.42
1:X:228:SER:HB2	1:X:236:VAL:HB	2.01	0.42
1:Y:184:GLN:HB2	1:Y:197:ILE:HD12	2.02	0.42
1:X:229:MET:HA	3:X:305:HOH:O	2.19	0.42
1:Y:87:ILE:O	1:Y:87:ILE:HG22	2.20	0.41
1:X:123:VAL:HG11	1:X:125:GLN:HE21	1.85	0.41
1:Y:235:LEU:O	1:Y:250:TYR:HA	2.19	0.41
1:Z:104:GLU:OE2	1:Z:108:GLN:HG3	2.19	0.41
1:X:45:VAL:HG13	2:U:1070:SER:HB3	2.02	0.41
1:Z:28:TRP:CE3	1:Z:35:VAL:HG11	2.56	0.41
1:Z:16:LEU:HG	1:Z:79:LEU:HD12	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	X	242/261 (93%)	235 (97%)	7 (3%)	0	100	100
1	Y	235/261 (90%)	231 (98%)	3 (1%)	1 (0%)	34	42
1	Z	241/261 (92%)	236 (98%)	4 (2%)	1 (0%)	34	42
2	U	9/11 (82%)	8 (89%)	1 (11%)	0	100	100
2	V	5/11 (46%)	4 (80%)	1 (20%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	W	6/11 (54%)	6 (100%)	0	0	100	100
All	All	738/816 (90%)	720 (98%)	16 (2%)	2 (0%)	41	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Z	108	GLN
1	Y	104	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	X	216/228 (95%)	212 (98%)	4 (2%)	57	73
1	Y	212/228 (93%)	204 (96%)	8 (4%)	33	47
1	Z	215/228 (94%)	204 (95%)	11 (5%)	24	33
2	U	10/10 (100%)	10 (100%)	0	100	100
2	V	7/10 (70%)	6 (86%)	1 (14%)	3	3
2	W	8/10 (80%)	7 (88%)	1 (12%)	4	5
All	All	668/714 (94%)	643 (96%)	25 (4%)	34	48

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Y	12	LEU
1	Y	36	ASN
1	Y	58	ASP
1	Y	64	ARG
1	Y	128	ILE
1	Y	177	ASN
1	Y	203	VAL
1	Y	224	THR
1	X	66	LEU

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	X	120	ASP
1	X	162	CYS
1	X	243	ASP
1	Z	20	LYS
1	Z	25	GLU
1	Z	45	VAL
1	Z	65	ASN
1	Z	73	THR
1	Z	93	GLU
1	Z	97	ASP
1	Z	135	CYS
1	Z	156	ASP
1	Z	177	ASN
1	Z	244	MET
2	W	1070	SER
2	V	1074	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	Y	36	ASN
1	Y	38	GLN
1	Y	177	ASN
1	Y	179	ASN
1	Y	184	GLN
1	Y	213	ASN
1	Y	246	HIS
1	X	84	ASN
1	X	125	GLN
1	X	213	ASN
1	Z	38	GLN
1	Z	65	ASN
1	Z	125	GLN
1	Z	177	ASN

### 5.3.3 RNA

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	X	248/261 (95%)	0.27	16 (6%) 18 24	28, 45, 74, 103	0
1	Y	243/261 (93%)	0.36	16 (6%) 18 23	36, 51, 78, 109	0
1	Z	247/261 (94%)	0.53	22 (8%) 9 13	30, 51, 83, 114	0
2	U	11/11 (100%)	0.99	2 (18%) 1 1	44, 55, 78, 86	0
2	V	7/11 (63%)	1.56	3 (42%) 0 0	68, 72, 86, 88	0
2	W	8/11 (72%)	0.68	2 (25%) 0 0	44, 49, 62, 64	0
All	All	764/816 (93%)	0.41	61 (7%) 12 16	28, 50, 80, 114	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Z	123	VAL	7.8
1	Z	122	ASP	6.5
1	Z	95	ASN	4.9
1	X	186	SER	4.8
2	U	1079	LYS	4.7
1	Y	105	ALA	4.6
1	Y	125	GLN	4.2
1	Z	96	ALA	4.1
1	X	96	ALA	4.1
1	X	125	GLN	3.8
1	Y	130	GLU	3.7
1	X	122	ASP	3.7
1	X	93	GLU	3.6
2	V	1071	ASP	3.5
1	Z	124	GLU	3.4
2	V	1077	VAL	3.3
1	Z	84	ASN	3.3
1	X	107	ASN	3.2
2	V	1074	ARG	3.1

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	Y	101	LEU	3.1
1	Z	106	PRO	3.0
1	Y	120	ASP	3.0
1	Z	1	MET	3.0
1	Y	109	GLU	3.0
1	Z	120	ASP	2.9
1	X	165	ASP	2.9
1	X	130	GLU	2.9
1	Z	174	GLU	2.8
1	Z	94	ASP	2.8
1	X	164	LYS	2.8
1	X	251	LEU	2.8
1	Y	191	GLU	2.7
1	Z	108	GLN	2.7
1	Z	130	GLU	2.7
1	Z	235	LEU	2.7
1	Y	95	ASN	2.7
1	X	95	ASN	2.6
2	W	1074	ARG	2.6
1	Y	87	ILE	2.6
1	X	120	ASP	2.6
1	Z	194	ALA	2.6
1	Y	189	ASP	2.5
2	W	1077	VAL	2.4
1	X	235	LEU	2.4
1	Z	126	LEU	2.4
1	Z	236	VAL	2.3
1	Y	82	ALA	2.3
1	Z	107	ASN	2.3
1	Y	156	ASP	2.3
1	Z	97	ASP	2.3
1	Z	125	GLN	2.3
2	U	1078	LYS	2.2
1	X	123	VAL	2.2
1	Y	232	ASP	2.2
1	X	236	VAL	2.2
1	Y	126	LEU	2.1
1	X	193	GLU	2.1
1	Y	243	ASP	2.1
1	Y	190	LYS	2.1
1	Z	227	LEU	2.0
1	Z	255	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.