



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 7, 2023 – 04:11 PM EDT

PDB ID : 6DDP  
Title : Mycobacterium tuberculosis Dihydrofolate Reductase complexed with beta-NADPH and 3'-[(2R)-4-(2,4-diamino-6-ethylpyrimidin-5-yl)but-3-yn-2-yl]-5'-methoxy[1,1'-biphenyl]-4-carboxylic acid  
Authors : Hajian, B.; Wright, D.  
Deposited on : 2018-05-10  
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	 4% (poor fit), 87% (0-1 outliers), 12% (2-3 outliers)
1	B	159	 6% (poor fit), 91% (0-1 outliers), 9% (2-3 outliers)
1	C	159	 6% (poor fit), 92% (0-1 outliers), 8% (2-3 outliers)
1	D	159	 3% (poor fit), 94% (0-1 outliers), 6% (2-3 outliers)

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5652 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydrofolate reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	159	Total 1261	C 794	N 230	O 232	S 5	0	2	0
1	B	159	Total 1260	C 793	N 230	O 232	S 5	0	2	0
1	C	159	Total 1244	C 783	N 228	O 228	S 5	0	0	0
1	D	159	Total 1264	C 794	N 233	O 232	S 5	0	2	0

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C<sub>21</sub>H<sub>30</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	B	1	Total 48	C 21	N 7	O 17	P 3	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is 3'-[(2R)-4-(2,4-diamino-6-ethylpyrimidin-5-yl)but-3-yn-2-yl]-5'-methoxy[1,1'-biphenyl]-4-carboxylic acid (three-letter code: G6Y) (formula: C<sub>24</sub>H<sub>24</sub>N<sub>4</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			31	24	4	3		
3	B	1	Total	C	N	O	0	0
			31	24	4	3		
3	C	1	Total	C	N	O	0	0
			31	24	4	3		
3	D	1	Total	C	N	O	0	0
			31	24	4	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	72	Total	O	0	0
			72	72		
4	B	81	Total	O	0	0
			81	81		
4	C	75	Total	O	0	0
			75	75		

Continued on next page...



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	D	79	Total	O	0	0
			79	79		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dihydrofolate reductase




- Molecule 1: Dihydrofolate reductase



- Molecule 1: Dihydrofolate reductase



- Molecule 1: Dihydrofolate reductase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.77Å 60.55Å 60.64Å 90.04° 90.02° 90.03°	Depositor
Resolution (Å)	34.42 – 1.49 34.42 – 1.46	Depositor EDS
% Data completeness (in resolution range)	83.1 (34.42-1.49) 82.8 (34.42-1.46)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.02	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.54 (at 1.46Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.203 , 0.225 0.209 , 0.228	Depositor DCC
$R_{free}$ test set	2060 reflections (1.74%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.1	Xtriage
Anisotropy	0.611	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 40.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.31$	Xtriage

*Continued on next page...*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

Continued from previous page...

Property	Value	Source
Estimated twinning fraction	0.477 for h,l,-k	Xtriage
	0.477 for h,-l,k	
	0.029 for -l,k,h	
	0.029 for l,k,-h	
	0.034 for k,-h,l	
	0.034 for -k,h,l	
	0.027 for k,l,h	
	0.027 for l,h,k	
	0.028 for l,-h,-k	
	0.028 for -k,-l,h	
	0.034 for -l,-h,k	
	0.034 for -k,l,-h	
	0.036 for k,-l,-h	
	0.036 for -l,h,-k	
	0.487 for h,-k,-l	
	0.075 for -h,k,-l	
	0.076 for -h,-k,l	
	0.037 for k,h,-l	
0.035 for -k,-h,-l		
0.029 for l,-k,h		
0.032 for -l,-k,-h		
0.074 for -h,l,k		
0.076 for -h,-l,-k		
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5652	wwPDB-VP
Average B, all atoms ( $\text{\AA}^2$ )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.32% of the height of the origin peak. No significant pseudotranslation is detected.*



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, G6Y

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/1292	0.58	1/1755 (0.1%)
1	B	0.32	0/1291	0.53	0/1754
1	C	0.32	0/1275	0.52	0/1732
1	D	0.33	0/1295	0.53	0/1758
All	All	0.33	0/5153	0.54	1/6999 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	ARG	C-N-CA	5.10	134.45	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1261	0	1238	14	0
1	B	1260	0	1236	9	0
1	C	1244	0	1223	10	0
1	D	1264	0	1240	6	0
2	A	48	0	26	2	0
2	B	48	0	26	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	48	0	26	1	0
2	D	48	0	26	1	0
3	A	31	0	0	1	0
3	B	31	0	0	0	0
3	C	31	0	0	0	0
3	D	31	0	0	0	0
4	A	72	0	0	0	0
4	B	81	0	0	0	0
4	C	75	0	0	4	0
4	D	79	0	0	0	0
All	All	5652	0	5041	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:53:LYS:NZ	4:C:303:HOH:O	2.20	0.72
1:C:147:SER:O	4:C:301:HOH:O	2.11	0.68
1:A:54:VAL:HG22	1:A:57:LEU:HD23	1.75	0.67
1:C:83:GLU:OE2	4:C:302:HOH:O	2.14	0.66
1:C:67:ARG:NH1	4:C:304:HOH:O	2.23	0.64
1:D:136:ARG:NH1	1:D:137:GLY:O	2.31	0.63
1:A:82:LEU:HD22	1:A:102:LEU:HG	1.79	0.63
1:A:146:ARG:HB2	1:A:147:SER:HB2	1.81	0.61
1:A:19:ASP:OD1	1:A:20:ILE:N	2.33	0.61
1:C:82:LEU:HD22	1:C:102:LEU:HG	1.85	0.58
1:B:82:LEU:HD22	1:B:102:LEU:HG	1.85	0.58
1:A:54:VAL:HG21	3:A:202:G6Y:CAV	2.37	0.55
1:A:14:ILE:O	2:A:201:NDP:H2N	2.06	0.55
1:C:26:GLU:HG3	1:C:146:ARG:HH21	1.73	0.53
1:A:68:GLN:HB3	1:A:70:ASP:OD1	2.09	0.52
1:C:14:ILE:O	2:C:202:NDP:H2N	2.12	0.50
1:A:65[A]:LEU:HG	2:A:201:NDP:N3A	2.27	0.49
1:B:14:ILE:O	2:B:201:NDP:H2N	2.12	0.49
1:D:14:ILE:O	2:D:201:NDP:H2N	2.12	0.49
1:C:47:TRP:CE2	1:C:73:ALA:HB1	2.48	0.48
1:B:65[B]:LEU:HG	2:B:201:NDP:N3A	2.28	0.48
1:B:47:TRP:CZ2	1:B:73:ALA:HB1	2.49	0.47
1:C:47:TRP:CZ2	1:C:73:ALA:HB1	2.51	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:47:TRP:CE2	1:B:73:ALA:HB1	2.51	0.46
1:C:4:LEU:HD13	1:C:107:ALA:HB2	1.98	0.46
1:D:116:ASP:HB3	1:D:150:ARG:HB3	1.99	0.45
1:B:68:GLN:HB2	1:B:71:PHE:HB2	1.99	0.45
1:A:65[B]:LEU:HD23	1:A:79:VAL:HG23	1.99	0.43
1:B:77:GLU:HG3	1:D:106:TYR:CD2	2.53	0.43
1:A:146:ARG:N	1:A:147:SER:HB3	2.33	0.43
1:A:4:LEU:HD13	1:A:107:ALA:HB2	2.00	0.42
1:D:82:LEU:HD22	1:D:102:LEU:HG	2.02	0.42
1:D:22:TRP:CE3	1:D:117:ILE:HD11	2.55	0.42
1:A:47:TRP:CZ2	1:A:73:ALA:HB1	2.55	0.41
1:A:22:TRP:CE3	1:A:117:ILE:HD11	2.56	0.41
1:A:77:GLU:HG3	1:B:106:TYR:CD2	2.55	0.41
1:B:4:LEU:HD13	1:B:107:ALA:HB2	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/159 (100%)	155 (98%)	3 (2%)	1 (1%)	25	7
1	B	159/159 (100%)	157 (99%)	2 (1%)	0	100	100
1	C	157/159 (99%)	155 (99%)	2 (1%)	0	100	100
1	D	159/159 (100%)	156 (98%)	3 (2%)	0	100	100
All	All	634/636 (100%)	623 (98%)	10 (2%)	1 (0%)	47	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	147	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	129/127 (102%)	128 (99%)	1 (1%)	81	66
1	B	129/127 (102%)	127 (98%)	2 (2%)	62	36
1	C	127/127 (100%)	127 (100%)	0	100	100
1	D	129/127 (102%)	129 (100%)	0	100	100
All	All	514/508 (101%)	511 (99%)	3 (1%)	86	74

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	144	PHE
1	B	72	MET
1	B	144	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	G6Y	A	202	-	32,33,33	2.35	6 (18%)	41,46,46	1.70	10 (24%)
2	NDP	C	202	-	45,52,52	2.29	6 (13%)	53,80,80	1.34	5 (9%)
2	NDP	A	201	-	45,52,52	2.23	6 (13%)	53,80,80	1.34	5 (9%)
3	G6Y	C	201	-	32,33,33	2.40	6 (18%)	41,46,46	1.75	10 (24%)
2	NDP	D	201	-	45,52,52	2.22	6 (13%)	53,80,80	1.33	6 (11%)
3	G6Y	B	202	-	32,33,33	2.25	4 (12%)	41,46,46	1.84	11 (26%)
2	NDP	B	201	-	45,52,52	2.18	5 (11%)	53,80,80	1.35	7 (13%)
3	G6Y	D	202	-	32,33,33	2.25	5 (15%)	41,46,46	1.76	8 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	G6Y	A	202	-	-	0/18/21/21	0/3/3/3
2	NDP	C	202	-	-	4/30/77/77	0/5/5/5
2	NDP	A	201	-	-	4/30/77/77	0/5/5/5
3	G6Y	C	201	-	-	0/18/21/21	0/3/3/3
2	NDP	D	201	-	-	4/30/77/77	0/5/5/5
3	G6Y	B	202	-	-	0/18/21/21	0/3/3/3
2	NDP	B	201	-	-	3/30/77/77	0/5/5/5
3	G6Y	D	202	-	-	0/18/21/21	0/3/3/3

All (44) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	202	NDP	P2B-O2B	12.46	1.82	1.59
2	B	201	NDP	P2B-O2B	12.10	1.82	1.59
2	D	201	NDP	P2B-O2B	12.03	1.82	1.59
2	A	201	NDP	P2B-O2B	11.84	1.81	1.59
3	C	201	G6Y	CAL-CAK	-10.39	1.42	1.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	202	G6Y	CAL-CAK	-9.95	1.43	1.52
3	D	202	G6Y	CAL-CAK	-9.49	1.43	1.52
3	B	202	G6Y	CAL-CAK	-9.37	1.43	1.52
3	A	202	G6Y	CBD-C4	-5.62	1.41	1.51
3	B	202	G6Y	CBD-C4	-5.39	1.41	1.51
3	C	201	G6Y	CBD-C4	-5.25	1.42	1.51
3	D	202	G6Y	CBD-C4	-5.07	1.42	1.51
3	A	202	G6Y	CAU-CAZ	-4.28	1.40	1.49
3	D	202	G6Y	CAU-CAZ	-4.26	1.40	1.49
3	B	202	G6Y	CAU-CAZ	-4.07	1.40	1.49
2	A	201	NDP	PN-O5D	4.07	1.75	1.59
2	C	202	NDP	PN-O5D	3.94	1.75	1.59
3	C	201	G6Y	CAU-CAZ	-3.92	1.41	1.49
2	D	201	NDP	PN-O5D	3.78	1.74	1.59
2	B	201	NDP	PN-O5D	3.44	1.73	1.59
2	A	201	NDP	O2B-C2B	-3.19	1.32	1.44
2	C	202	NDP	O2B-C2B	-3.14	1.32	1.44
3	A	202	G6Y	CAR-CAN	-3.05	1.41	1.49
3	B	202	G6Y	CAR-CAN	-3.01	1.41	1.49
3	C	201	G6Y	CAR-CAN	-2.99	1.41	1.49
3	D	202	G6Y	CAR-CAN	-2.96	1.41	1.49
2	D	201	NDP	O2B-C2B	-2.93	1.33	1.44
2	B	201	NDP	O2B-C2B	-2.81	1.33	1.44
3	C	201	G6Y	CAI-CAJ	-2.60	1.16	1.19
2	C	202	NDP	C7N-N7N	2.57	1.40	1.33
2	D	201	NDP	O4B-C1B	2.52	1.44	1.41
2	C	202	NDP	O4B-C1B	2.45	1.44	1.41
2	D	201	NDP	C2A-N1A	2.39	1.38	1.33
2	D	201	NDP	C7N-N7N	2.37	1.39	1.33
2	B	201	NDP	C7N-N7N	2.34	1.39	1.33
2	A	201	NDP	C2A-N1A	2.31	1.38	1.33
3	D	202	G6Y	CAI-CAJ	2.21	1.21	1.19
2	B	201	NDP	C2A-N1A	2.19	1.38	1.33
2	A	201	NDP	O4B-C1B	2.10	1.44	1.41
2	A	201	NDP	C7N-N7N	2.08	1.38	1.33
3	A	202	G6Y	C5-C6	-2.07	1.39	1.42
3	C	201	G6Y	C5-C6	-2.07	1.39	1.42
3	A	202	G6Y	CAI-CAJ	-2.04	1.17	1.19
2	C	202	NDP	C2A-N1A	2.03	1.37	1.33

All (62) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	202	G6Y	C2-N3-C4	7.44	122.44	116.24
3	B	202	G6Y	C2-N3-C4	7.12	122.18	116.24
3	C	201	G6Y	C2-N3-C4	6.59	121.73	116.24
3	A	202	G6Y	C2-N3-C4	6.51	121.66	116.24
3	B	202	G6Y	N3-C2-N1	-3.63	119.73	125.42
2	A	201	NDP	PN-O3-PA	-3.59	120.51	132.83
3	C	201	G6Y	C5-C6-NAH	-3.59	117.48	121.62
2	A	201	NDP	O2B-P2B-O1X	-3.48	95.94	109.39
2	D	201	NDP	PN-O3-PA	-3.48	120.90	132.83
2	C	202	NDP	PN-O3-PA	-3.31	121.48	132.83
2	D	201	NDP	O2B-P2B-O1X	-3.29	96.68	109.39
2	C	202	NDP	O2B-P2B-O1X	-3.29	96.71	109.39
3	D	202	G6Y	N3-C2-N1	-3.28	120.28	125.42
2	B	201	NDP	PN-O3-PA	-3.24	121.72	132.83
3	B	202	G6Y	C5-C6-NAH	-3.24	117.89	121.62
2	B	201	NDP	O2B-P2B-O1X	-3.22	96.97	109.39
3	A	202	G6Y	N3-C2-N1	-3.19	120.42	125.42
3	C	201	G6Y	N3-C2-N1	-3.12	120.53	125.42
3	B	202	G6Y	C2-N1-C6	2.97	120.27	116.99
3	C	201	G6Y	NAH-C6-N1	2.97	121.23	117.03
3	A	202	G6Y	C5-C6-NAH	-2.89	118.28	121.62
3	B	202	G6Y	NAH-C6-N1	2.84	121.05	117.03
3	D	202	G6Y	C5-C6-NAH	-2.77	118.42	121.62
3	A	202	G6Y	C6-C5-CAI	2.77	123.37	118.57
3	A	202	G6Y	C2-N1-C6	2.75	120.03	116.99
3	B	202	G6Y	C6-C5-CAI	2.68	123.21	118.57
2	A	201	NDP	O3X-P2B-O2X	2.62	117.64	107.64
2	C	202	NDP	O3X-P2B-O2X	2.60	117.57	107.64
3	A	202	G6Y	NAH-C6-N1	2.59	120.70	117.03
2	B	201	NDP	O3X-P2B-O2X	2.59	117.52	107.64
2	D	201	NDP	O3X-P2B-O2X	2.55	117.39	107.64
3	D	202	G6Y	C6-C5-CAI	2.55	122.98	118.57
3	C	201	G6Y	C6-C5-CAI	2.51	122.92	118.57
3	B	202	G6Y	NAG-C2-N3	2.50	121.14	117.25
3	B	202	G6Y	CAY-OAX-CAP	-2.43	112.23	117.51
2	D	201	NDP	C2A-N1A-C6A	-2.39	114.66	118.75
3	D	202	G6Y	OBA-CAZ-CAU	2.39	121.05	114.85
3	D	202	G6Y	C2-N1-C6	2.36	119.59	116.99
3	C	201	G6Y	CAY-OAX-CAP	-2.35	112.41	117.51
3	A	202	G6Y	OBA-CAZ-CAU	2.32	120.86	114.85
2	B	201	NDP	O2N-PN-O1N	2.29	123.54	112.24
3	C	201	G6Y	C2-N1-C6	2.26	119.48	116.99
3	D	202	G6Y	NAH-C6-N1	2.25	120.21	117.03

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	201	G6Y	OBA-CAZ-CAU	2.23	120.64	114.85
3	B	202	G6Y	CAL-CAK-CAJ	-2.21	108.93	112.33
3	C	201	G6Y	OBA-CAZ-OB	-2.21	118.45	123.35
3	B	202	G6Y	OBA-CAZ-CAU	2.20	120.55	114.85
2	C	202	NDP	C2A-N1A-C6A	-2.20	115.00	118.75
3	A	202	G6Y	CAL-CAK-CAJ	-2.18	108.97	112.33
2	B	201	NDP	C2A-N1A-C6A	-2.17	115.03	118.75
3	A	202	G6Y	CAY-OAX-CAP	-2.17	112.80	117.51
2	B	201	NDP	C2B-C3B-C4B	-2.17	97.29	101.99
3	D	202	G6Y	CAL-CAK-CAJ	-2.16	109.00	112.33
3	A	202	G6Y	NAG-C2-N3	2.16	120.61	117.25
2	A	201	NDP	PA-O5B-C5B	-2.16	109.04	121.68
3	B	202	G6Y	OBA-CAZ-OB	-2.13	118.63	123.35
2	D	201	NDP	PA-O5B-C5B	-2.12	109.24	121.68
2	A	201	NDP	C2A-N1A-C6A	-2.12	115.13	118.75
2	C	202	NDP	C2B-C3B-C4B	-2.06	97.52	101.99
2	B	201	NDP	PA-O5B-C5B	-2.06	109.63	121.68
2	D	201	NDP	C2B-C3B-C4B	-2.04	97.55	101.99
3	C	201	G6Y	CAL-CAK-CAJ	-2.03	109.21	112.33

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	NDP	PA-O3-PN-O5D
2	B	201	NDP	PA-O3-PN-O5D
2	C	202	NDP	PA-O3-PN-O5D
2	D	201	NDP	PA-O3-PN-O5D
2	A	201	NDP	O4D-C1D-N1N-C2N
2	B	201	NDP	O4D-C1D-N1N-C2N
2	C	202	NDP	O4D-C1D-N1N-C2N
2	D	201	NDP	O4D-C1D-N1N-C2N
2	B	201	NDP	C2D-C1D-N1N-C2N
2	D	201	NDP	C2D-C1D-N1N-C2N
2	A	201	NDP	C2D-C1D-N1N-C2N
2	C	202	NDP	C2D-C1D-N1N-C2N
2	A	201	NDP	C5B-O5B-PA-O1A
2	C	202	NDP	C5B-O5B-PA-O1A
2	D	201	NDP	C5B-O5B-PA-O1A


There are no ring outliers.


5 monomers are involved in 7 short contacts:





Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	202	G6Y	1	0
2	C	202	NDP	1	0
2	A	201	NDP	2	0
2	D	201	NDP	1	0
2	B	201	NDP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






















## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	159/159 (100%)	0.43	7 (4%) 34 38	17, 25, 43, 55	0
1	B	159/159 (100%)	0.46	10 (6%) 20 21	17, 25, 44, 57	0
1	C	159/159 (100%)	0.48	9 (5%) 23 25	17, 25, 43, 57	0
1	D	159/159 (100%)	0.41	4 (2%) 57 62	17, 25, 42, 57	0
All	All	636/636 (100%)	0.45	30 (4%) 31 34	17, 25, 43, 57	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	147	SER	4.1
1	A	144	PHE	4.0
1	C	72	MET	3.9
1	C	146	ARG	3.9
1	C	144	PHE	3.5
1	A	147	SER	3.5
1	D	51	PRO	3.2
1	A	23	ARG	3.2
1	B	71	PHE	3.2
1	B	136	ARG	3.2
1	A	159	SER	3.1
1	C	51	PRO	2.8
1	C	23	ARG	2.7
1	B	72	MET	2.7
1	A	146	ARG	2.6
1	A	148	GLY	2.6
1	B	159	SER	2.6
1	B	69	ALA	2.6
1	B	52	ALA	2.4
1	C	149	LEU	2.4
1	D	52	ALA	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	87	THR	2.2
1	D	70	ASP	2.2
1	B	117	ILE	2.2
1	C	70	ASP	2.2
1	D	149	LEU	2.2
1	A	72	MET	2.1
1	B	144	PHE	2.1
1	B	23	ARG	2.1
1	B	149	LEU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	G6Y	B	202	31/31	0.88	0.10	16,24,35,38	0
3	G6Y	D	202	31/31	0.88	0.11	16,22,31,36	0
3	G6Y	C	201	31/31	0.90	0.10	17,24,33,39	0
3	G6Y	A	202	31/31	0.91	0.10	15,24,33,37	0
2	NDP	B	201	48/48	0.94	0.10	18,24,32,35	0
2	NDP	C	202	48/48	0.94	0.10	18,24,33,35	0
2	NDP	A	201	48/48	0.94	0.10	17,24,32,36	0
2	NDP	D	201	48/48	0.95	0.09	19,24,33,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around G6Y B 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around G6Y D 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around G6Y C 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around G6Y A 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around NDP B 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around NDP C 202:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NDP A 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.