



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 06:24 PM EDT

PDB ID : 6EMY  
Title : Structure of the Tn1549 transposon Integrase (aa 82-397, Y379F) in complex with transposon right end DNA  
Authors : Schulz, E.C.; Rubio-Cosials, A.; Barabas, O.  
Deposited on : 2017-10-04  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	76% 22% ..
1	B	317	79% 20% .
2	C	20	45% 50% 5%
2	E	20	60% 30% 10%
3	D	26	50% 46% .

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	26	 69% 27%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7063 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Int protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	315	2513	1592	445	461	15	0	0	0
1	B	316	2525	1597	449	463	16	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	81	SER	-	expression tag	UNP Q7BP35
A	379	PHE	TYR	engineered mutation	UNP Q7BP35
B	81	SER	-	expression tag	UNP Q7BP35
B	379	PHE	TYR	engineered mutation	UNP Q7BP35

- Molecule 2 is a DNA chain called DNA (20-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	19	385	187	65	114	19	0	0	0
2	E	18	362	177	63	105	17	0	0	0

- Molecule 3 is a DNA chain called DNA (26-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	F	25	511	249	87	151	24	0	0	0
3	D	26	533	259	92	157	25	0	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	81	Total O 81 81	0	0
4	B	75	Total O 75 75	0	0
4	C	13	Total O 13 13	0	0
4	E	14	Total O 14 14	0	0
4	F	22	Total O 22 22	0	0
4	D	29	Total O 29 29	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Int protein

Chain A: 



- Molecule 1: Int protein

Chain B: 




- Molecule 2: DNA (20-MER)

Chain C: 



- Molecule 2: DNA (20-MER)

Chain E: 



- Molecule 3: DNA (26-MER)



- Molecule 3: DNA (26-MER)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.23Å 121.68Å 77.23Å 90.00° 118.12° 90.00°	Depositor
Resolution (Å)	38.60 – 2.50 38.60 – 2.50	Depositor EDS
% Data completeness (in resolution range)	93.0 (38.60-2.50) 93.0 (38.60-2.50)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.43 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.161 , 0.184 0.158 , 0.178	Depositor DCC
$R_{free}$ test set	2034 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	61.4	Xtriage
Anisotropy	0.583	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 45.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.020 for -h-l,k,h 0.020 for l,k,-h-l 0.028 for h,-k,-h-l 0.028 for -h-l,-k,l 0.478 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7063	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/2562	0.42	0/3451
1	B	0.26	0/2575	0.43	0/3470
2	C	0.66	0/430	1.02	0/660
2	E	0.57	0/405	0.96	0/622
3	D	0.75	1/597 (0.2%)	1.08	2/921 (0.2%)
3	F	0.62	0/572	1.08	0/882
All	All	0.41	1/7141 (0.0%)	0.68	2/10006 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	5	DC	C1'-N1	6.10	1.57	1.49

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	22	DT	C5-C4-O4	-5.74	120.89	124.90
3	D	22	DT	N3-C4-O4	5.57	123.24	119.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	305	PHE	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2513	0	2514	54	0
1	B	2525	0	2526	44	0
2	C	385	0	218	12	0
2	E	362	0	207	6	0
3	D	533	0	300	14	0
3	F	511	0	289	7	0
4	A	81	0	0	2	0
4	B	75	0	0	0	0
4	C	13	0	0	0	0
4	D	29	0	0	0	0
4	E	14	0	0	0	0
4	F	22	0	0	0	0
All	All	7063	0	6054	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:347:ARG:NH1	1:A:368:MET:O	1.99	0.93
1:B:242:ARG:NH1	1:B:276:GLU:OE2	2.02	0.92
1:A:252:ARG:NH1	2:C:-9:DT:OP2	2.03	0.90
1:A:298:ASP:OD2	4:A:401:HOH:O	1.98	0.81
1:A:391:GLU:OE1	1:A:394:ARG:NH1	2.12	0.81
1:A:152:LYS:NZ	1:A:175:PHE:O	2.15	0.78
1:B:223:GLY:O	1:B:347:ARG:NH1	2.20	0.74
3:F:1:DT:O4	3:D:4:DT:N3	2.19	0.73
1:B:328:LYS:NZ	2:E:-12:DC:OP1	2.21	0.73
1:B:307:LYS:NZ	2:E:-9:DT:OP2	2.22	0.72
1:A:336:ALA:HB1	1:A:337:LEU:HB2	1.70	0.72
1:A:386:ASP:N	1:A:386:ASP:OD1	2.22	0.71
1:A:284:ARG:O	1:A:288:ASN:ND2	2.22	0.71
1:A:382:HIS:HA	1:B:362:LYS:HD3	1.72	0.71
1:B:139:LYS:NZ	1:B:184:ASP:OD1	2.23	0.71

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:ARG:HH12	2:C:-9:DT:P	2.14	0.69
1:A:163:ILE:O	1:B:169:ARG:NH2	2.26	0.68
1:A:347:ARG:NH2	2:C:-1:DT:OP1	2.25	0.67
1:A:252:ARG:HH11	1:A:307:LYS:NZ	1.93	0.67
1:A:188:LYS:NZ	2:C:-3:DT:OP1	2.29	0.66
1:A:99:LYS:HD2	3:D:8:DA:H3'	1.77	0.66
3:F:16:DG:H2'	3:F:17:DG:C8	2.31	0.65
1:A:377:LEU:HD11	1:B:377:LEU:HD11	1.78	0.64
1:B:91:GLN:HG3	1:B:158:ALA:HA	1.81	0.62
1:A:242:ARG:HB3	1:A:274:MET:HB2	1.82	0.60
1:A:337:LEU:HG	1:A:338:PRO:HD2	1.82	0.59
1:B:116:ASP:OD1	1:B:118:LEU:N	2.25	0.58
1:A:391:GLU:OE2	1:B:360:ASN:N	2.21	0.58
2:C:-16:DA:H2	3:D:23:DT:H3	1.51	0.58
1:B:82:MET:O	1:B:123:ILE:N	2.36	0.58
1:B:153:ARG:HG2	3:F:6:DA:H5''	1.86	0.58
1:B:306:LEU:HA	1:B:312:PRO:HA	1.85	0.58
1:A:293:LYS:HG3	1:A:294:PRO:HD2	1.85	0.57
1:A:238:ASP:O	1:A:242:ARG:N	2.38	0.57
1:B:252:ARG:HE	1:B:314:THR:HG21	1.70	0.56
1:B:162:ALA:HA	1:B:165:ASP:HB3	1.86	0.56
2:E:-9:DT:H2'	2:E:-8:DA:C8	2.40	0.56
1:B:242:ARG:HH12	1:B:276:GLU:CD	2.08	0.55
1:A:342:THR:HG23	1:A:344:HIS:H	1.73	0.54
1:A:276:GLU:OE1	1:A:280:GLN:NE2	2.41	0.54
1:A:388:ALA:HB1	1:B:367:ILE:HD11	1.91	0.52
2:C:-18:DA:H2''	2:C:-17:DA:C8	2.44	0.52
2:C:-19:DT:H3	3:D:25:DG:H22	1.58	0.51
1:A:152:LYS:HG3	1:A:173:PHE:HD2	1.75	0.51
1:B:158:ALA:O	1:B:161:THR:OG1	2.25	0.51
1:B:161:THR:O	1:B:165:ASP:N	2.44	0.51
1:B:226:ILE:HD11	1:B:318:TYR:HD2	1.74	0.51
1:A:327:LYS:O	1:A:331:LYS:HG2	2.11	0.51
1:B:327:LYS:O	1:B:331:LYS:HD3	2.10	0.50
1:A:128:LEU:HD11	1:A:175:PHE:HB3	1.93	0.50
1:B:342:THR:HG21	3:F:11:DT:H3'	1.92	0.50
1:B:208:VAL:HG11	2:E:-12:DC:H5''	1.92	0.50
1:A:156:LYS:HG3	1:A:171:ASN:HD21	1.76	0.49
1:B:83:THR:O	1:B:87:LEU:HB2	2.12	0.49
1:A:238:ASP:OD1	1:A:240:GLU:HG2	2.12	0.49
1:B:153:ARG:NH2	3:F:6:DA:OP1	2.42	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:19:DA:H2''	3:F:20:DT:H5''	1.96	0.48
1:A:250:LEU:HD11	1:A:258:TYR:HD2	1.78	0.48
3:D:12:DA:C8	3:D:12:DA:H5'	2.49	0.48
1:B:260:ILE:HG13	1:B:298:ASP:HB2	1.96	0.47
1:A:252:ARG:HH11	1:A:307:LYS:HZ1	1.59	0.47
1:B:247:SER:HA	1:B:269:VAL:HG13	1.97	0.47
1:B:349:THR:O	1:B:353:ARG:HG2	2.14	0.47
1:A:252:ARG:NH1	1:A:307:LYS:NZ	2.62	0.47
1:B:270:ARG:HD2	1:B:366:TYR:O	2.16	0.46
1:A:163:ILE:HG22	1:A:168:ILE:HG22	1.97	0.46
1:B:157:ALA:O	1:B:161:THR:HG23	2.16	0.46
1:B:163:ILE:HG13	1:B:169:ARG:HB3	1.97	0.46
3:D:16:DG:H2''	3:D:17:DG:H5''	1.97	0.46
1:B:192:THR:OG1	1:B:195:GLN:HB2	2.16	0.46
1:A:394:ARG:NH1	4:A:407:HOH:O	2.49	0.45
2:C:-15:DA:H61	3:D:20:DT:H3	1.63	0.45
1:A:103:LYS:HE3	1:A:106:ARG:HH11	1.82	0.44
1:B:344:HIS:CE1	2:E:-2:DT:H5''	2.52	0.44
3:F:1:DT:H2'	3:F:1:DT:O2	2.16	0.44
1:A:84:VAL:HG23	1:A:121:CYS:O	2.18	0.44
1:A:373:ILE:HD13	1:B:377:LEU:HD13	1.98	0.44
1:B:279:TYR:O	1:B:283:GLN:HG2	2.17	0.44
1:A:336:ALA:HB1	1:A:337:LEU:CB	2.46	0.43
1:A:84:VAL:HB	1:A:119:GLY:HA2	2.00	0.43
1:B:144:SER:O	1:B:148:ILE:HG13	2.17	0.43
3:D:5:DC:H2'	3:D:5:DC:O2	2.17	0.43
1:A:159:PHE:O	1:A:163:ILE:HG23	2.18	0.43
3:D:4:DT:C4'	3:D:5:DC:H5'	2.49	0.43
1:A:372:ASN:ND2	3:D:4:DT:O4	2.52	0.43
3:D:6:DA:H1'	3:D:7:DA:C8	2.54	0.42
2:E:-14:DT:H2''	2:E:-13:DC:OP2	2.18	0.42
2:C:-17:DA:C5	2:C:-16:DA:N1	2.88	0.42
1:A:113:LEU:HD11	1:A:155:LEU:HD21	2.01	0.42
1:A:180:VAL:HG12	1:A:181:LEU:HD12	2.01	0.42
1:A:252:ARG:C	1:A:253:ASN:HD22	2.23	0.42
1:A:272:ILE:HA	1:A:273:PRO:HD3	1.92	0.42
1:B:234:ASP:N	1:B:302:ASN:O	2.43	0.41
2:C:-17:DA:H2	3:D:24:DA:H2	1.67	0.41
2:C:-11:DC:H42	3:D:16:DG:H1	1.66	0.41
1:A:238:ASP:HB3	1:A:243:VAL:HB	2.02	0.41
1:A:329:TYR:HE1	1:A:336:ALA:HB3	1.84	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:231:GLY:O	1:B:248:HIS:NE2	2.53	0.41
1:B:359:MET:O	1:B:380:TYR:OH	2.34	0.41
1:A:208:VAL:HG11	2:C:-12:DC:H5''	2.01	0.41
1:A:377:LEU:HA	1:A:380:TYR:HD1	1.86	0.41
1:A:357:ALA:HB3	1:B:395:LEU:HD23	2.03	0.41
1:A:262:GLU:HA	1:A:263:PRO:HD3	1.97	0.40
1:B:116:ASP:OD1	1:B:117:ARG:N	2.54	0.40
1:B:171:ASN:HA	1:B:172:PRO:HD3	1.78	0.40
1:A:282:PHE:O	1:A:285:VAL:HG12	2.21	0.40
1:A:240:GLU:CD	1:A:240:GLU:H	2.23	0.40
1:B:109:LEU:O	1:B:113:LEU:HG	2.21	0.40
3:D:5:DC:C2	3:D:5:DC:H5''	2.56	0.40
1:A:134:TRP:O	1:A:138:MET:HG2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	311/317 (98%)	301 (97%)	10 (3%)	0	100	100
1	B	314/317 (99%)	296 (94%)	18 (6%)	0	100	100
All	All	625/634 (99%)	597 (96%)	28 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/275 (98%)	257 (96%)	12 (4%)	27	51
1	B	271/275 (98%)	264 (97%)	7 (3%)	46	72
All	All	540/550 (98%)	521 (96%)	19 (4%)	36	62

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	GLN
1	A	140	GLU
1	A	233	THR
1	A	252	ARG
1	A	264	LYS
1	A	282	PHE
1	A	306	LEU
1	A	321	MET
1	A	334	GLU
1	A	337	LEU
1	A	344	HIS
1	A	386	ASP
1	B	83	THR
1	B	124	ASP
1	B	167	CYS
1	B	169	ARG
1	B	266	GLN
1	B	318	TYR
1	B	344	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	315/317 (99%)	-0.52	0 <a href="#">100</a>   <a href="#">100</a>	45, 70, 97, 116	0
1	B	316/317 (99%)	-0.55	1 (0%) <a href="#">94</a>   <a href="#">94</a>	46, 70, 98, 124	0
2	C	19/20 (95%)	-1.03	0 <a href="#">100</a>   <a href="#">100</a>	52, 69, 129, 133	0
2	E	18/20 (90%)	-0.84	0 <a href="#">100</a>   <a href="#">100</a>	60, 80, 132, 149	0
3	D	26/26 (100%)	-0.77	0 <a href="#">100</a>   <a href="#">100</a>	53, 102, 160, 182	0
3	F	25/26 (96%)	-0.89	0 <a href="#">100</a>   <a href="#">100</a>	50, 92, 123, 141	0
All	All	719/726 (99%)	-0.58	1 (0%) <a href="#">95</a>   <a href="#">96</a>	45, 71, 110, 182	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	94	HIS	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.