



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 25, 2024 – 09:47 AM EDT

PDB ID : 6G86
Title : Structure of Cdc14 bound to SIC1 PxL motif
Authors : Mouilleron, S.; Kataria, M.; Uhlmann, F.
Deposited on : 2018-04-07
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3764 (1.76-1.72)
Clashscore	141614	3923 (1.76-1.72)
Ramachandran outliers	138981	3878 (1.76-1.72)
Sidechain outliers	138945	3878 (1.76-1.72)
RSRZ outliers	127900	3705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	
1	B	374	
2	C	16	
2	D	16	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PEG	A	403	-	-	X	-
5	PEG	A	405	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6919 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein phosphatase CDC14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	357	Total 2944	C 1912	N 490	O 530	S 12	0	11	0
1	B	367	Total 3011	C 1955	N 510	O 534	S 12	0	9	0

- Molecule 2 is a protein called Protein SIC1.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	D	10	Total 69	C 48	N 10	O 11	0	0	0
2	C	11	Total 78	C 52	N 12	O 14	0	0	0

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: C₁₂H₂₆O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			16	10	6		
4	B	1	Total	C	O	0	0
			19	12	7		
4	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0
5	B	1	Total C O 7 4 3	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Zn 1 1	0	0
6	B	1	Total Zn 1 1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	353	Total O 353 353	0	3
7	B	349	Total O 349 349	0	3

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	D	4	Total O 4 4	0	0
7	C	2	Total O 2 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tyrosine-protein phosphatase CDC14



- Molecule 1: Tyrosine-protein phosphatase CDC14



- Molecule 2: Protein SIC1



- Molecule 2: Protein SIC1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.09Å 94.32Å 127.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.80 – 1.74 75.86 – 1.74	Depositor EDS
% Data completeness (in resolution range)	99.6 (63.80-1.74) 99.6 (75.86-1.74)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 1.74Å)	Xtrriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.166 , 0.199 0.165 , 0.198	Depositor DCC
R_{free} test set	4627 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	23.6	Xtrriage
Anisotropy	0.090	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6919	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PEG, MES, P6G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/3062	0.63	2/4158 (0.0%)
1	B	0.51	0/3127	0.62	0/4247
2	C	0.40	0/81	0.58	0/111
2	D	0.33	0/72	0.48	0/99
All	All	0.51	0/6342	0.62	2/8615 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	70	LEU	CA-CB-CG	6.43	130.08	115.30
1	A	121	ASP	CB-CG-OD2	5.21	122.99	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	52	PHE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2944	0	2845	24	0
1	B	3011	0	2937	20	0
2	C	78	0	74	1	0
2	D	69	0	65	1	0
3	A	12	0	12	1	0
3	B	12	0	12	1	0
4	A	16	0	21	2	0
4	B	32	0	43	5	0
5	A	28	0	39	12	0
5	B	7	0	10	1	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	353	0	0	8	0
7	B	349	0	0	3	0
7	C	2	0	0	0	0
7	D	4	0	0	0	0
All	All	6919	0	6058	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55:ASN:HD21	5:A:403:PEG:H31	1.43	0.82
1:A:57:GLY:HA3	5:A:403:PEG:H11	1.75	0.68
1:A:167:TYR:HE2	5:A:403:PEG:H22	1.64	0.62
1:A:154:LYS:NZ	4:A:402:P6G:H122	2.18	0.58
1:A:334:ASN:OD1	1:A:337[A]:ARG:NH2	2.36	0.58
1:A:340:LYS:NZ	7:A:503:HOH:O	2.37	0.58
3:A:401:MES:O1S	7:A:501:HOH:O	2.17	0.57
1:B:236:LYS:NZ	7:B:507:HOH:O	2.38	0.56
1:A:168:GLU:HG2	5:A:403:PEG:H12	1.88	0.54
1:B:334:ASN:OD1	1:B:337[A]:ARG:NH2	2.41	0.52
1:B:120[B]:VAL:HG11	1:B:124:PHE:CE2	2.45	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:270:GLU:O	1:B:274:LYS:HG3	2.10	0.51
1:A:154:LYS:HZ1	4:A:402:P6G:H122	1.76	0.51
1:A:57:GLY:CA	5:A:403:PEG:H11	2.40	0.50
1:A:55:ASN:HD21	5:A:403:PEG:C3	2.18	0.50
5:A:405:PEG:H31	7:A:791:HOH:O	2.12	0.49
1:A:255:THR:HG23	5:A:405:PEG:H42	1.93	0.49
1:B:178:PHE:HB3	1:B:188:PHE:HB3	1.95	0.49
1:B:322:VAL:HG11	3:B:401:MES:H22	1.95	0.49
1:B:343:THR:HG22	4:B:403:P6G:H32	1.94	0.49
1:A:174:GLU:HG3	7:A:758:HOH:O	2.14	0.48
1:B:200:LEU:HB3	1:B:202:THR:HG22	1.95	0.47
1:B:302:TYR:HA	4:B:402:P6G:H62	1.96	0.47
1:B:178:PHE:HB3	1:B:188:PHE:CB	2.45	0.47
1:A:120[B]:VAL:HG21	1:A:124:PHE:CE2	2.51	0.46
1:B:327:HIS:HD2	7:B:512:HOH:O	2.01	0.44
1:B:70:LEU:HD21	2:C:58:LEU:HD11	1.98	0.44
1:B:154:LYS:HE3	4:B:402:P6G:H61	2.00	0.44
1:B:327:HIS:CD2	7:B:512:HOH:O	2.70	0.44
1:A:57:GLY:C	5:A:403:PEG:H11	2.38	0.44
1:A:61:ARG:CZ	1:A:164:LEU:HD21	2.48	0.43
5:A:406:PEG:H42	7:A:771:HOH:O	2.18	0.43
1:A:341:TYR:OH	1:B:338:GLU:HG3	2.19	0.42
1:B:154:LYS:NZ	4:B:402:P6G:H82	2.35	0.42
1:B:178:PHE:CB	1:B:188:PHE:HB3	2.50	0.42
1:A:70:LEU:HD13	1:A:80:VAL:HG21	2.01	0.42
1:A:255:THR:CG2	5:A:405:PEG:H42	2.49	0.42
1:A:345:ILE:HD13	7:A:805:HOH:O	2.20	0.42
1:A:337[A]:ARG:NH2	7:A:509:HOH:O	2.47	0.41
1:A:188:PHE:CE2	1:A:282:HIS:HB3	2.55	0.41
7:A:763:HOH:O	4:B:402:P6G:H22	2.20	0.41
1:A:39:GLU:OE2	1:A:84:SER:HB2	2.21	0.41
1:B:99:CYS:SG	1:B:117:LEU:HD11	2.60	0.41
1:B:156:LEU:O	5:B:404:PEG:H31	2.21	0.41
5:A:405:PEG:H42	5:A:405:PEG:H22	1.83	0.41
2:D:52:PHE:HB2	2:D:55:ALA:HB2	2.03	0.41
1:A:99:CYS:SG	1:A:117:LEU:HD11	2.61	0.40
1:B:159:LEU:HD12	1:B:159:LEU:HA	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	364/374 (97%)	354 (97%)	9 (2%)	1 (0%)	41	23
1	B	376/374 (100%)	364 (97%)	11 (3%)	1 (0%)	41	23
2	C	9/16 (56%)	8 (89%)	1 (11%)	0	100	100
2	D	8/16 (50%)	6 (75%)	2 (25%)	0	100	100
All	All	757/780 (97%)	732 (97%)	23 (3%)	2 (0%)	41	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	321	VAL
1	B	321	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	317/325 (98%)	308 (97%)	9 (3%)	43	19
1	B	322/325 (99%)	318 (99%)	4 (1%)	71	56
2	C	8/14 (57%)	8 (100%)	0	100	100
2	D	6/14 (43%)	5 (83%)	1 (17%)	2	0
All	All	653/678 (96%)	639 (98%)	14 (2%)	57	30

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17[A]	ARG
1	A	17[B]	ARG
1	A	24[A]	ASP
1	A	24[B]	ASP
1	A	70	LEU
1	A	188	PHE
1	A	211	PHE
1	A	237	LYS
1	A	369	LYS
1	B	70	LEU
1	B	87	SER
1	B	188	PHE
1	B	211	PHE
2	D	57	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	P6G	B	403	-	12,12,18	0.50	0	11,11,17	0.19	0
3	MES	A	401	-	12,12,12	1.76	2 (16%)	14,16,16	1.62	2 (14%)
5	PEG	A	406	1	6,6,6	0.50	0	5,5,5	0.53	0
4	P6G	B	402	-	18,18,18	0.44	0	17,17,17	0.41	0
5	PEG	A	404	-	6,6,6	0.50	0	5,5,5	0.25	0
5	PEG	B	404	-	6,6,6	0.48	0	5,5,5	0.20	0
5	PEG	A	405	-	6,6,6	0.62	0	5,5,5	0.67	0
4	P6G	A	402	-	15,15,18	0.50	0	14,14,17	0.60	0
5	PEG	A	403	-	6,6,6	0.34	0	5,5,5	0.74	0
3	MES	B	401	-	12,12,12	2.08	1 (8%)	14,16,16	2.43	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	P6G	B	403	-	-	1/10/10/16	-
3	MES	A	401	-	-	2/6/14/14	0/1/1/1
5	PEG	A	406	1	-	3/4/4/4	-
4	P6G	B	402	-	-	7/16/16/16	-
5	PEG	A	404	-	-	1/4/4/4	-
5	PEG	B	404	-	-	1/4/4/4	-
5	PEG	A	405	-	-	2/4/4/4	-
4	P6G	A	402	-	-	4/13/13/16	-
5	PEG	A	403	-	-	1/4/4/4	-
3	MES	B	401	-	-	5/6/14/14	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	MES	C8-S	-6.61	1.68	1.77
3	A	401	MES	C8-S	-5.24	1.70	1.77
3	A	401	MES	O1S-S	2.22	1.51	1.45

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	MES	C5-N4-C3	5.07	120.23	108.83
3	B	401	MES	C5-N4-C3	4.97	120.02	108.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	MES	C6-C5-N4	-3.70	104.50	110.10
3	B	401	MES	C7-N4-C3	3.53	120.27	111.23
3	B	401	MES	C2-C3-N4	-3.32	105.06	110.10
3	B	401	MES	C7-N4-C5	2.71	118.17	111.23
3	B	401	MES	O3S-S-C8	2.34	109.55	105.77
3	A	401	MES	C7-N4-C3	2.14	116.72	111.23

There are no chirality outliers.

All (27) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	401	MES	C8-C7-N4-C3
5	A	405	PEG	C4-C3-O2-C2
4	B	402	P6G	O4-C5-C6-O7
5	A	405	PEG	O1-C1-C2-O2
3	B	401	MES	C7-C8-S-O3S
5	A	406	PEG	O1-C1-C2-O2
3	B	401	MES	C8-C7-N4-C5
4	B	403	P6G	O1-C2-C3-O4
5	B	404	PEG	C4-C3-O2-C2
4	B	402	P6G	C6-C5-O4-C3
4	B	402	P6G	C9-C8-O7-C6
4	A	402	P6G	C6-C5-O4-C3
4	B	402	P6G	C18-C17-O16-C15
5	A	406	PEG	O2-C3-C4-O4
3	B	401	MES	C7-C8-S-O1S
3	B	401	MES	C7-C8-S-O2S
5	A	403	PEG	C4-C3-O2-C2
4	B	402	P6G	O7-C8-C9-O10
4	A	402	P6G	O1-C2-C3-O4
4	A	402	P6G	O13-C14-C15-O16
4	B	402	P6G	O1-C2-C3-O4
3	A	401	MES	C8-C7-N4-C5
4	B	402	P6G	C5-C6-O7-C8
3	A	401	MES	C7-C8-S-O3S
5	A	406	PEG	C1-C2-O2-C3
4	A	402	P6G	O10-C11-C12-O13
5	A	404	PEG	O2-C3-C4-O4


There are no ring outliers.

9 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	403	P6G	1	0
3	A	401	MES	1	0
5	A	406	PEG	1	0
4	B	402	P6G	4	0
5	B	404	PEG	1	0
5	A	405	PEG	4	0
4	A	402	P6G	2	0
5	A	403	PEG	7	0
3	B	401	MES	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	357/374 (95%)	-0.44	1 (0%) 94 95	14, 23, 49, 86	0
1	B	367/374 (98%)	-0.37	5 (1%) 75 81	15, 25, 53, 86	0
2	C	11/16 (68%)	1.58	4 (36%) 0 0	37, 46, 111, 119	0
2	D	10/16 (62%)	1.19	3 (30%) 0 0	35, 43, 94, 100	0
All	All	745/780 (95%)	-0.35	13 (1%) 70 76	14, 24, 56, 119	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	52	PHE	4.5
2	C	53	LYS	4.1
2	D	53	LYS	3.8
1	B	198	GLY	3.0
2	C	51	SER	2.9
1	A	367	LEU	2.8
2	C	54	ASN	2.8
1	B	204	SER	2.7
1	B	203	LYS	2.5
1	B	7	LEU	2.4
2	D	54	ASN	2.4
2	C	52	PHE	2.3
1	B	205	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	PEG	A	406	7/7	0.48	0.29	59,60,61,61	0
5	PEG	B	404	7/7	0.64	0.22	70,71,72,73	0
5	PEG	A	405	7/7	0.74	0.33	46,49,57,58	0
5	PEG	A	404	7/7	0.77	0.18	61,62,62,63	0
5	PEG	A	403	7/7	0.77	0.34	54,59,62,62	0
4	P6G	A	402	16/19	0.85	0.13	47,54,58,60	0
4	P6G	B	403	13/19	0.86	0.13	51,54,57,57	0
4	P6G	B	402	19/19	0.86	0.17	49,53,62,64	0
3	MES	B	401	12/12	0.91	0.16	42,59,66,67	0
3	MES	A	401	12/12	0.95	0.10	25,34,40,41	0
6	ZN	B	405	1/1	0.95	0.06	36,36,36,36	0
6	ZN	A	407	1/1	0.99	0.08	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around P6G A 402:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around P6G B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.