



# Full wwPDB X-ray Structure Validation Report ⓘ

May 4, 2024 – 12:27 pm BST

PDB ID : 6H6S  
Title : Sad phasing on nickel-substituted human carbonic anhydrase II  
Authors : Calderone, V.; Fragai, M.; Silva, J.P.; Luchinat, C.; Ravera, E.; Geraldles, C.F.G.C.; Macedo, A.L.; Cerofolini, L.; Giuntini, S.  
Deposited on : 2018-07-30  
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	259	 2% 86% 12% .
1	B	259	 3% 89% 8% ..

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4779 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbonic anhydrase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	258	2058	1321	355	380	2	0	0	0
1	B	258	2058	1321	355	380	2	0	0	0

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ni	0	0
			2	2		
2	B	2	Total	Ni	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	325	Total	O	0	0
			325	325		
3	B	334	Total	O	0	0
			334	334		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Carbonic anhydrase 2



- Molecule 1: Carbonic anhydrase 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.48Å 41.18Å 84.04Å 90.00° 109.25° 90.00°	Depositor
Resolution (Å)	45.34 – 1.45 45.34 – 1.45	Depositor EDS
% Data completeness (in resolution range)	95.6 (45.34-1.45) 95.5 (45.34-1.45)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.92 (at 1.45Å)	Xtrriage
Refinement program	REFMAC 5.8.0230	Depositor
R, $R_{free}$	0.150 , 0.204 0.159 , 0.158	Depositor DCC
$R_{free}$ test set	4033 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.4	Xtrriage
Anisotropy	0.270	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 35.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4779	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 96.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.3132e-10. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.94	4/2120 (0.2%)	1.04	8/2877 (0.3%)
1	B	1.15	10/2120 (0.5%)	1.19	11/2877 (0.4%)
All	All	1.05	14/4240 (0.3%)	1.12	19/5754 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	4	TRP	C-N	-22.72	0.92	1.33
1	B	3	HIS	CA-C	14.68	1.91	1.52
1	B	3	HIS	C-O	13.87	1.49	1.23
1	A	234	GLU	CD-OE1	10.52	1.37	1.25
1	B	3	HIS	N-CA	-10.46	1.25	1.46
1	A	68	GLU	CD-OE1	10.10	1.36	1.25
1	B	259	LYS	C-O	9.88	1.42	1.23
1	B	212	GLU	CD-OE1	-8.13	1.16	1.25
1	B	3	HIS	C-N	6.94	1.50	1.34
1	B	3	HIS	CA-CB	6.39	1.68	1.53
1	B	4	TRP	N-CA	6.31	1.58	1.46
1	A	234	GLU	CD-OE2	6.17	1.32	1.25
1	A	212	GLU	CD-OE1	-5.93	1.19	1.25
1	B	41	PRO	C-O	5.24	1.33	1.23

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	4	TRP	O-C-N	-20.74	87.95	123.20
1	B	4	TRP	CA-C-N	19.73	155.67	116.20
1	B	4	TRP	C-N-CA	16.44	156.83	122.30
1	A	26	ARG	NE-CZ-NH2	-10.71	114.94	120.30
1	A	57	ARG	NE-CZ-NH2	9.41	125.01	120.30
1	B	3	HIS	O-C-N	9.40	137.74	122.70
1	B	2	HIS	CA-C-N	-9.07	97.26	117.20
1	B	3	HIS	CB-CA-C	-8.55	93.29	110.40
1	A	26	ARG	CG-CD-NE	-8.51	93.94	111.80
1	B	3	HIS	N-CA-CB	7.32	123.78	110.60
1	A	2	HIS	CB-CA-C	6.54	123.47	110.40
1	B	3	HIS	CA-C-O	-5.98	107.54	120.10
1	A	173	ASP	CB-CG-OD1	-5.73	113.14	118.30
1	B	224	PHE	CB-CG-CD1	5.63	124.74	120.80
1	A	88	ARG	NE-CZ-NH2	-5.34	117.63	120.30
1	B	180	ARG	NE-CZ-NH2	-5.22	117.69	120.30
1	A	57	ARG	CD-NE-CZ	-5.07	116.50	123.60
1	A	26	ARG	NH1-CZ-NH2	5.06	124.96	119.40
1	B	224	PHE	CB-CG-CD2	-5.06	117.26	120.80

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	57	ARG	Sidechain
1	B	2	HIS	Mainchain

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2058	0	2008	35	0
1	B	2058	0	2008	37	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	325	0	0	14	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	334	0	0	6	0
All	All	4779	0	4016	69	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:HIS:CA	1:B:3:HIS:C	1.91	1.38
1:B:2:HIS:ND1	1:B:168:LYS:NZ	1.84	1.26
1:A:57:ARG:HB2	1:A:57:ARG:NH1	1.51	1.24
1:A:57:ARG:CB	1:A:57:ARG:HH11	1.52	1.20
1:A:57:ARG:NH1	1:A:57:ARG:CB	2.03	1.20
1:A:57:ARG:HB2	1:A:57:ARG:HH11	0.92	1.07
1:B:2:HIS:CG	1:B:168:LYS:HZ2	1.73	1.06
1:B:2:HIS:CG	1:B:168:LYS:NZ	2.24	1.04
1:A:131:LYS:CE	3:A:436:HOH:O	2.14	0.95
1:B:91:GLN:HE22	1:B:93:HIS:HD1	1.08	0.95
1:B:2:HIS:CE1	3:B:401:HOH:O	2.24	0.90
1:A:131:LYS:HE2	3:A:436:HOH:O	1.72	0.88
1:B:3:HIS:O	1:B:10:ASN:HB2	1.80	0.81
1:B:3:HIS:O	1:B:10:ASN:CB	2.30	0.80
1:A:57:ARG:NH1	1:A:57:ARG:HB3	1.95	0.80
1:B:2:HIS:CE1	1:B:168:LYS:HE3	2.17	0.80
1:A:251:ASN:HA	3:A:573:HOH:O	1.85	0.76
1:B:2:HIS:CG	1:B:168:LYS:HZ1	2.02	0.76
1:A:57:ARG:CB	1:A:57:ARG:CZ	2.63	0.74
1:A:57:ARG:HB3	1:A:57:ARG:CZ	2.16	0.73
1:A:57:ARG:HH11	1:A:57:ARG:CG	2.01	0.73
1:B:44:LYS:HG2	1:B:45:PRO:HD2	1.72	0.71
1:A:14:HIS:HD2	1:A:17:LYS:NZ	1.87	0.71
1:B:2:HIS:NE2	3:B:401:HOH:O	2.25	0.70
1:B:27:GLN:HE21	1:B:244:ARG:HH12	1.41	0.69
1:B:3:HIS:C	1:B:3:HIS:CB	2.61	0.69
1:B:54:THR:HG22	3:B:601:HOH:O	1.93	0.69
1:B:27:GLN:NE2	1:B:244:ARG:HH12	1.91	0.69
1:A:27:GLN:NE2	1:A:244:ARG:HH12	1.93	0.67
1:A:35:HIS:HD2	3:A:437:HOH:O	1.77	0.67
1:B:2:HIS:CE1	1:B:168:LYS:CE	2.78	0.66
1:A:156:GLN:HE22	1:B:176:ASN:HD21	1.44	0.64

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2:HIS:CE1	1:B:168:LYS:NZ	2.63	0.64
1:A:223:LYS:CE	3:A:404:HOH:O	2.45	0.63
1:A:35:HIS:CD2	3:A:437:HOH:O	2.50	0.63
1:A:14:HIS:HD2	1:A:17:LYS:HZ3	1.47	0.61
1:B:230:ASN:ND2	1:B:237:GLU:H	1.99	0.60
1:A:170:LYS:CE	3:A:403:HOH:O	2.48	0.60
1:B:2:HIS:ND1	1:B:168:LYS:CE	2.65	0.58
1:B:40:ASP:C	1:B:40:ASP:OD1	2.39	0.58
1:A:3:HIS:NE2	1:A:18:ASP:OD2	2.37	0.58
1:B:44:LYS:HG2	1:B:45:PRO:CD	2.33	0.58
1:A:27:GLN:HE21	1:A:244:ARG:HH12	1.50	0.57
1:B:3:HIS:C	1:B:3:HIS:CG	2.78	0.57
1:A:160:ASP:OD2	1:B:54:THR:HG21	2.07	0.54
1:B:4:TRP:CZ3	1:B:15:TRP:CZ2	2.99	0.51
1:B:3:HIS:O	1:B:10:ASN:HB3	2.11	0.50
1:A:131:LYS:HE3	3:A:436:HOH:O	1.96	0.49
1:A:16:HIS:HD2	3:A:653:HOH:O	1.96	0.49
1:B:54:THR:CG2	3:B:601:HOH:O	2.55	0.48
1:B:91:GLN:NE2	1:B:93:HIS:HD1	1.92	0.48
1:B:157:LYS:HE3	3:B:653:HOH:O	2.13	0.48
1:B:4:TRP:CE3	1:B:15:TRP:CZ2	3.03	0.46
1:A:19:PHE:CE2	1:A:199:PRO:HB3	2.51	0.46
1:B:16:HIS:HD2	3:B:667:HOH:O	1.99	0.46
1:A:170:LYS:HD3	3:A:403:HOH:O	2.16	0.45
1:A:170:LYS:NZ	3:A:403:HOH:O	2.48	0.45
1:B:2:HIS:CB	1:B:168:LYS:NZ	2.79	0.45
1:B:40:ASP:OD1	1:B:42:SER:N	2.34	0.45
1:B:230:ASN:ND2	1:B:230:ASN:H	2.15	0.45
1:A:118:HIS:CE1	3:A:401:HOH:O	2.70	0.44
1:A:170:LYS:CD	3:A:403:HOH:O	2.65	0.44
1:A:132:ALA:O	1:A:138:GLY:HA3	2.17	0.44
1:A:160:ASP:OD2	1:B:54:THR:CG2	2.65	0.44
1:A:16:HIS:CG	1:A:23:LYS:HE2	2.53	0.43
1:A:219:GLU:O	1:A:223:LYS:HG2	2.19	0.43
1:B:228:ASN:HB3	1:B:230:ASN:ND2	2.33	0.43
1:A:5:GLY:O	1:A:11:GLY:HA2	2.20	0.41
1:A:157:LYS:HG3	3:A:418:HOH:O	2.19	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:508:HOH:O	3:A:627:HOH:O[2_747]	2.16	0.04

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/259 (99%)	248 (97%)	8 (3%)	0	100	100
1	B	256/259 (99%)	245 (96%)	10 (4%)	1 (0%)	34	13
All	All	512/518 (99%)	493 (96%)	18 (4%)	1 (0%)	47	22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	3	HIS

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/224 (100%)	217 (97%)	6 (3%)	44	12
1	B	223/224 (100%)	219 (98%)	4 (2%)	59	26
All	All	446/448 (100%)	436 (98%)	10 (2%)	52	18

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LYS
1	A	51	ASP
1	A	57	ARG
1	A	91	GLN
1	A	131	LYS
1	A	173	ASP
1	B	44	LYS
1	B	51	ASP
1	B	91	GLN
1	B	259	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	HIS
1	A	14	HIS
1	A	16	HIS
1	A	27	GLN
1	A	35	HIS
1	A	228	ASN
1	B	16	HIS
1	B	27	GLN
1	B	91	GLN
1	B	176	ASN
1	B	228	ASN
1	B	230	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	4:TRP	C	5:GLY	N	0.92

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	258/259 (99%)	0.03	5 (1%) 66 68	8, 13, 26, 62	0
1	B	258/259 (99%)	0.10	7 (2%) 54 56	8, 13, 30, 72	0
All	All	516/518 (99%)	0.07	12 (2%) 60 63	8, 13, 28, 72	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	HIS	18.0
1	B	3	HIS	8.1
1	B	4	TRP	8.0
1	B	41	PRO	3.5
1	B	259	LYS	3.4
1	B	251	ASN	3.2
1	B	42	SER	3.1
1	A	2	HIS	2.8
1	A	251	ASN	2.8
1	A	233	GLY	2.4
1	A	259	LYS	2.4
1	A	3	HIS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NI	B	301	1/1	0.84	0.14	51,51,51,51	0
2	NI	A	302	1/1	1.00	0.07	9,9,9,9	0
2	NI	A	301	1/1	1.00	0.21	31,31,31,31	0
2	NI	B	302	1/1	1.00	0.08	9,9,9,9	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.