



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 25, 2024 – 09:45 AM EDT

PDB ID : 6HZL  
Title : Crystal structure of redox-inhibited phosphoribulokinase from *Synechococcus* sp. (strain PCC 6301), osmate derivative  
Authors : Wilson, R.H.; Bracher, A.; Hartl, F.U.; Hayer-Hartl, M.  
Deposited on : 2018-10-23  
Resolution : 2.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	333	
1	B	333	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	OS	A	402	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4957 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoribulokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	329	Total	C	N	O	S	0	0	0
			2605	1647	448	499	11			
1	B	295	Total	C	N	O	S	0	0	0
			2349	1491	405	444	9			

- Molecule 2 is OSMIUM ION (three-letter code: OS) (formula: Os).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Os	0	0
			2	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O	0	0
			1	1		




### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Phosphoribulokinase



#### • Molecule 1: Phosphoribulokinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	141.43Å 141.43Å 206.56Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.77 29.36 – 2.77	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.00-2.77) 100.0 (29.36-2.77)	Depositor EDS
$R_{merge}$	0.23	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.83 (at 2.76Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.235 , 0.259 0.232 , 0.256	Depositor DCC
$R_{free}$ test set	1017 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	79.5	Xtriage
Anisotropy	0.156	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 50.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4957	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	115.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.27	0/2660	0.50	0/3607
1	B	0.26	0/2398	0.51	0/3250
All	All	0.27	0/5058	0.50	0/6857

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2605	0	2547	3	0
1	B	2349	0	2317	4	0
2	A	2	0	0	0	0
3	A	1	0	0	0	0
All	All	4957	0	4864	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:37:THR:HG23	1:B:104:ASN:HD22	1.57	0.69
1:A:37:THR:HG23	1:A:104:ASN:HD22	1.58	0.68
1:A:49:ARG:NH2	1:A:90:HIS:O	2.44	0.51
1:B:49:ARG:NH2	1:B:90:HIS:O	2.45	0.49
1:B:185:MET:HB2	1:B:186:PRO:HD2	1.95	0.47
1:A:52:ARG:NH2	1:A:88:TYR:OH	2.50	0.41
1:B:52:ARG:NH2	1:B:88:TYR:OH	2.51	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/333 (98%)	322 (98%)	5 (2%)	0	100	100
1	B	289/333 (87%)	285 (99%)	4 (1%)	0	100	100
All	All	616/666 (92%)	607 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	279/287 (97%)	279 (100%)	0	100	100
1	B	253/287 (88%)	253 (100%)	0	100	100

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	532/574 (93%)	532 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	104	ASN
1	B	104	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	329/333 (98%)	0.27	13 (3%)	38 33	53, 74, 116, 183	0
1	B	295/333 (88%)	1.57	90 (30%)	0 0	79, 146, 248, 271	0
All	All	624/666 (93%)	0.88	103 (16%)	1 1	53, 95, 229, 271	0

All (103) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	57	VAL	12.5
1	B	331	ALA	11.9
1	B	326	ALA	11.5
1	B	94	LEU	10.6
1	B	63	ARG	10.3
1	B	56	GLY	9.5
1	B	332	SER	7.4
1	B	60	LEU	7.0
1	B	327	ALA	6.6
1	B	329	VAL	6.1
1	B	123	LEU	5.7
1	B	164	ARG	5.6
1	B	14	GLY	5.5
1	B	95	ILE	5.4
1	B	55	ALA	5.3
1	B	90	HIS	4.9
1	B	328	PRO	4.9
1	B	48	ASP	4.6
1	B	54	GLU	4.6
1	B	252	HIS	4.6
1	B	97	PRO	4.5
1	B	330	ALA	4.5
1	B	91	GLU	4.5
1	B	58	THR	4.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	47	LEU	4.3
1	B	52	ARG	4.3
1	A	150	GLY	4.3
1	B	50	LYS	4.3
1	B	74	VAL	4.3
1	B	6	ARG	4.2
1	B	62	PRO	4.1
1	B	35	LEU	4.1
1	B	78	LYS	4.1
1	B	68	ASP	3.9
1	A	4	PRO	3.9
1	B	132	ILE	3.8
1	B	163	ARG	3.8
1	B	92	THR	3.7
1	B	11	GLY	3.7
1	B	125	ASP	3.7
1	B	105	ARG	3.6
1	A	194	THR	3.6
1	B	324	SER	3.6
1	B	4	PRO	3.5
1	B	49	ARG	3.5
1	B	99	GLU	3.5
1	B	46	SER	3.4
1	B	319	TYR	3.4
1	B	325	GLN	3.4
1	B	12	VAL	3.4
1	B	51	GLY	3.4
1	B	197	LYS	3.3
1	A	147	ALA	3.3
1	B	81	GLU	3.3
1	B	111	GLY	3.2
1	B	161	GLU	3.2
1	B	96	ASP	3.2
1	B	103	PRO	3.1
1	B	8	VAL	3.1
1	B	32	GLY	3.1
1	A	145	ASP	3.1
1	B	67	PHE	3.0
1	B	113	HIS	2.9
1	B	114	PRO	2.9
1	B	85	LYS	2.8
1	B	250	TYR	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	332	SER	2.7
1	B	65	ASN	2.7
1	B	88	TYR	2.7
1	B	170	TYR	2.7
1	B	109	ILE	2.7
1	B	84	MET	2.6
1	B	110	GLU	2.6
1	B	322	LEU	2.6
1	A	149	ARG	2.5
1	B	53	LYS	2.5
1	B	166	ASP	2.4
1	B	126	PHE	2.4
1	B	248	THR	2.4
1	B	75	LYS	2.4
1	B	13	ALA	2.3
1	B	89	ASN	2.3
1	A	148	GLU	2.3
1	B	93	GLY	2.3
1	A	192	ASN	2.3
1	B	167	PHE	2.3
1	B	321	ARG	2.3
1	B	311	THR	2.2
1	B	181	VAL	2.2
1	B	127	SER	2.2
1	B	119	ARG	2.2
1	B	79	ASN	2.2
1	B	106	ILE	2.1
1	A	11	GLY	2.1
1	B	168	LYS	2.1
1	B	307	PHE	2.1
1	B	194	THR	2.1
1	A	128	VAL	2.1
1	B	107	ILE	2.1
1	B	249	TYR	2.1
1	A	210	ASP	2.1
1	B	33	THR	2.0
1	A	151	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	OS	A	402	1/1	0.30	0.44	155,155,155,155	1
2	OS	A	401	1/1	0.91	0.27	182,182,182,182	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.