



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 16, 2023 – 04:29 AM JST

PDB ID : 6KWB
Title : AtDAO1(dioxygenase for auxin oxidation 1 from Arabidopsis thaliana) - 2-oxoglutarate binray complex
Authors : Rhee, S.; Jin, S.; Lee, H.
Deposited on : 2019-09-06
Resolution : 2.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5743 (2.54-2.50)
Clashscore	141614	6463 (2.54-2.50)
Ramachandran outliers	138981	6335 (2.54-2.50)
Sidechain outliers	138945	6337 (2.54-2.50)
RSRZ outliers	127900	5630 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	270	 11% 79% 20% .
1	B	270	 11% 80% 18% ..
1	C	270	 24% 61% 23% . 15%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	B	301	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6067 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-oxoglutarate (2OG) and Fe(II)-dependent oxygenase super-family protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	267	2114	1340	357	404	13	0	0	0
1	B	266	2106	1334	356	403	13	0	0	0
1	C	230	1823	1156	303	353	11	0	0	0


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	MET	-	initiating methionine	UNP Q9XI75
B	8	MET	-	initiating methionine	UNP Q9XI75
C	8	MET	-	initiating methionine	UNP Q9XI75

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 2-OXOGLUTARIC ACID (three-letter code: AKG) (formula: C₅H₆O₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 10 5 5	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	5	Total O 5 5	0	0
4	B	7	Total O 7 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2-oxoglutarate (2OG) and Fe(II)-dependent oxygenase superfamily protein




- Molecule 1: 2-oxoglutarate (2OG) and Fe(II)-dependent oxygenase superfamily protein



- Molecule 1: 2-oxoglutarate (2OG) and Fe(II)-dependent oxygenase superfamily protein





4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.88Å 76.80Å 166.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.21 – 2.52 28.21 – 2.52	Depositor EDS
% Data completeness (in resolution range)	98.2 (28.21-2.52) 98.2 (28.21-2.52)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.15 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.225 , 0.282 0.231 , 0.286	Depositor DCC
R_{free} test set	1993 reflections (6.04%)	wwPDB-VP
Wilson B-factor (Å ²)	71.0	Xtrriage
Anisotropy	0.316	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 62.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.024 for k,h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6067	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AKG, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.51	0/2160	0.71	1/2927 (0.0%)
1	B	0.50	0/2152	0.73	1/2916 (0.0%)
1	C	0.43	0/1857	0.71	2/2510 (0.1%)
All	All	0.49	0/6169	0.72	4/8353 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	93	LEU	CA-CB-CG	5.39	127.70	115.30
1	C	93	LEU	CA-CB-CG	5.20	127.26	115.30
1	C	188	ASP	CB-CG-OD1	5.08	122.87	118.30
1	B	21	ILE	CG1-CB-CG2	-5.05	100.29	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2114	0	2077	32	0
1	B	2106	0	2066	39	0
1	C	1823	0	1775	42	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
3	A	10	0	4	0	0
4	A	5	0	0	0	0
4	B	7	0	0	0	0
All	All	6067	0	5922	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (111) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:24:GLN:HE21	1:C:24:GLN:HA	1.25	0.97
1:B:131:THR:HG22	1:B:150:LYS:HE3	1.57	0.86
1:C:24:GLN:HE22	1:C:27:ARG:HD3	1.58	0.68
1:C:24:GLN:HA	1:C:24:GLN:NE2	2.04	0.67
1:C:41:HIS:NE2	1:C:212:ASN:O	2.27	0.67
1:B:225:TRP:HB2	1:B:260:LEU:HD21	1.77	0.67
1:C:77:GLY:O	1:C:157:ARG:NH2	2.27	0.66
1:B:31:GLU:OE2	1:B:141:TYR:OH	2.09	0.66
1:C:197:MET:HB2	1:C:204:PHE:CE1	2.32	0.65
1:A:258:THR:HG22	1:A:259:ASP:H	1.61	0.64
1:B:238:GLN:HB3	1:B:240:LYS:HZ1	1.63	0.64
1:C:225:TRP:HB2	1:C:260:LEU:HD21	1.80	0.63
1:A:77:GLY:O	1:A:157:ARG:NH2	2.31	0.63
1:C:112:SER:HB2	1:C:115:GLN:H	1.63	0.62
1:A:69:ASN:HA	1:A:108:GLN:OE1	2.00	0.62
1:B:164:LYS:HB3	1:B:165:PRO:HD2	1.81	0.61
1:C:104:THR:O	1:C:108:GLN:HG3	2.02	0.60
1:B:27:ARG:NH2	1:B:140:SER:O	2.33	0.59
1:B:238:GLN:HB3	1:B:240:LYS:NZ	2.18	0.59
1:A:161:TYR:HB2	1:A:245:ARG:HB3	1.86	0.58
1:B:166:GLU:HG3	1:B:166:GLU:O	2.04	0.58
1:C:189:GLU:O	1:C:190:ASN:ND2	2.37	0.57
1:A:270:GLU:OE2	1:B:88:TYR:OH	2.23	0.57
1:A:135:ARG:NH1	1:A:147:ASP:OD1	2.37	0.56
1:A:37:ARG:NH1	1:A:208:ASP:O	2.38	0.56
1:B:127:ASN:O	1:B:131:THR:HG23	2.05	0.56
1:B:261:GLU:OE2	1:B:262:PRO:HD2	2.06	0.56
1:C:258:THR:HG22	1:C:259:ASP:H	1.71	0.55
1:C:93:LEU:HD13	1:C:156:PHE:HB3	1.89	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:ILE:HG22	1:A:213:THR:HG22	1.89	0.54
1:C:63:TYR:O	1:C:67:VAL:HG23	2.06	0.54
1:B:144:VAL:HG23	1:B:145:GLU:HG3	1.88	0.54
1:A:122:TYR:CE2	1:A:126:ILE:HG13	2.42	0.54
1:C:135:ARG:HG2	1:C:149:PHE:CE1	2.42	0.54
1:A:54:THR:HG23	1:A:115:GLN:HG2	1.89	0.53
1:C:8:MET:HG2	1:C:9:ILE:N	2.23	0.52
1:B:175:LEU:HD11	1:B:234:LYS:HB3	1.91	0.52
1:B:45:LEU:HD22	1:B:212:ASN:HB3	1.91	0.51
1:C:135:ARG:HG2	1:C:149:PHE:HE1	1.75	0.51
1:C:61:ARG:HE	1:C:110:GLU:HB2	1.75	0.51
1:A:52:LYS:HZ3	1:A:189:GLU:HG3	1.75	0.51
1:C:145:GLU:H	1:C:145:GLU:CD	2.14	0.51
1:B:131:THR:O	1:B:135:ARG:HG3	2.11	0.51
1:B:181:PHE:HD1	1:B:182:LEU:HB2	1.74	0.50
1:A:165:PRO:O	1:A:168:VAL:HG13	2.11	0.50
1:A:93:LEU:HD13	1:A:156:PHE:HB3	1.92	0.50
1:C:46:SER:O	1:C:50:GLU:HG2	2.11	0.50
1:B:171:LEU:HA	1:B:238:GLN:HA	1.94	0.50
1:A:14:LEU:HD12	1:A:41:HIS:HA	1.94	0.49
1:B:149:PHE:CD1	1:B:150:LYS:HD3	2.47	0.49
1:A:69:ASN:OD1	1:A:77:GLY:HA2	2.13	0.49
1:B:149:PHE:O	1:B:152:TRP:HB2	2.13	0.49
1:B:225:TRP:O	1:B:263:PRO:HD2	2.12	0.49
1:C:263:PRO:HG2	1:C:266:PHE:CD2	2.47	0.49
1:C:56:ILE:HG12	1:C:246:TYR:CZ	2.48	0.49
1:A:201:SER:OG	1:A:203:THR:OG1	2.20	0.48
1:C:198:ASP:OD2	1:C:201:SER:HB3	2.14	0.48
1:B:149:PHE:HD1	1:B:150:LYS:HD3	1.79	0.47
1:C:131:THR:O	1:C:135:ARG:HG3	2.14	0.47
1:C:181:PHE:HA	1:C:221:MET:HB2	1.97	0.47
1:C:54:THR:HG23	1:C:115:GLN:HG2	1.97	0.47
1:C:131:THR:HG23	1:C:149:PHE:CE2	2.49	0.47
1:A:52:LYS:NZ	1:A:189:GLU:HG3	2.29	0.47
1:A:68:ARG:O	1:A:70:THR:HG23	2.14	0.47
1:A:51:MET:HE2	1:A:55:VAL:HG23	1.97	0.46
1:C:16:GLU:HG2	1:C:22:LEU:HB2	1.96	0.46
1:B:166:GLU:HA	1:B:168:VAL:HG22	1.98	0.46
1:C:18:SER:HB3	1:C:21:ILE:HD12	1.98	0.46
1:A:58:LEU:HD22	1:A:109:LEU:HD23	1.98	0.46
1:C:117:GLU:O	1:C:121:LYS:HB2	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:107:ASP:OD1	1:A:116:ARG:NH1	2.36	0.46
1:C:41:HIS:CB	1:C:43:VAL:HG22	2.46	0.46
1:A:194:LEU:HD23	1:A:207:ILE:HB	1.98	0.45
1:A:227:ASN:OD1	1:A:274:LEU:HD13	2.16	0.45
1:A:168:VAL:HA	1:A:239:CYS:SG	2.56	0.45
1:A:174:GLN:NE2	1:C:229:ARG:HE	2.14	0.45
1:A:104:THR:O	1:A:108:GLN:HG3	2.17	0.45
1:A:170:LYS:HD3	1:A:170:LYS:HA	1.60	0.44
1:B:135:ARG:HG2	1:B:149:PHE:CE2	2.52	0.44
1:B:144:VAL:HG23	1:B:145:GLU:N	2.32	0.44
1:B:59:PHE:CE1	1:B:89:GLU:HG2	2.52	0.44
1:B:135:ARG:HG2	1:B:149:PHE:CZ	2.51	0.44
1:B:104:THR:O	1:B:108:GLN:HG3	2.18	0.43
1:B:13:ASP:HA	1:B:39:ILE:HG13	2.01	0.43
1:B:58:LEU:HD23	1:B:58:LEU:HA	1.76	0.43
1:C:54:THR:O	1:C:58:LEU:HD13	2.18	0.43
1:B:122:TYR:CE2	1:B:126:ILE:HG13	2.53	0.43
1:C:14:LEU:HD23	1:C:14:LEU:HA	1.83	0.43
1:C:197:MET:HB2	1:C:204:PHE:CD1	2.53	0.43
1:C:196:ALA:HA	1:C:235:HIS:HA	2.01	0.43
1:B:150:LYS:HD2	1:B:150:LYS:HA	1.32	0.43
1:B:69:ASN:HA	1:B:108:GLN:OE1	2.19	0.42
1:B:93:LEU:HG	1:B:156:PHE:HB3	2.02	0.42
1:B:131:THR:CG2	1:B:150:LYS:HE3	2.37	0.42
1:B:14:LEU:HD12	1:B:41:HIS:HA	2.01	0.42
1:B:166:GLU:O	1:B:166:GLU:CG	2.66	0.42
1:C:59:PHE:HA	1:C:66:LYS:HZ1	1.85	0.42
1:A:119:MET:HE3	1:A:119:MET:HB3	1.97	0.41
1:B:23:ASN:HB3	1:B:140:SER:HB3	2.01	0.41
1:B:164:LYS:HB3	1:B:165:PRO:CD	2.49	0.41
1:C:41:HIS:HB3	1:C:43:VAL:HG22	2.02	0.41
1:A:85:ASN:HA	1:A:86:PRO:HD2	1.88	0.41
1:C:47:LEU:HD21	1:C:121:LYS:HB3	2.03	0.41
1:A:189:GLU:HG3	1:A:189:GLU:H	1.60	0.41
1:A:18:SER:HB3	1:A:21:ILE:HD12	2.03	0.40
1:C:267:VAL:O	1:C:268:ASP:OD2	2.40	0.40
1:A:24:GLN:O	1:A:28:GLU:HG2	2.21	0.40
1:C:41:HIS:HB2	1:C:43:VAL:H	1.86	0.40
1:C:133:LEU:HD13	1:C:133:LEU:HA	1.81	0.40
1:C:183:THR:O	1:C:216:ILE:HA	2.21	0.40
1:B:88:TYR:CZ	1:B:90:ALA:HB2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/270 (98%)	254 (96%)	11 (4%)	0	100	100
1	B	264/270 (98%)	251 (95%)	13 (5%)	0	100	100
1	C	218/270 (81%)	215 (99%)	3 (1%)	0	100	100
All	All	747/810 (92%)	720 (96%)	27 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/235 (99%)	227 (98%)	5 (2%)	52	75
1	B	231/235 (98%)	227 (98%)	4 (2%)	60	81
1	C	201/235 (86%)	194 (96%)	7 (4%)	36	60
All	All	664/705 (94%)	648 (98%)	16 (2%)	49	73

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	LEU
1	A	76	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	93	LEU
1	A	168	VAL
1	A	273	ARG
1	B	133	LEU
1	B	150	LYS
1	B	217	ASN
1	B	240	LYS
1	C	20	LYS
1	C	24	GLN
1	C	61	ARG
1	C	93	LEU
1	C	133	LEU
1	C	185	LEU
1	C	217	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	24	GLN
1	C	190	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	AKG	A	302	2	9,9,9	1.32	1 (11%)	11,11,11	1.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AKG	A	302	2	-	1/9/9/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302	AKG	C3-C2	2.12	1.53	1.51

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	302	AKG	C2-C3-C4-C5

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/270 (98%)	0.62	29 (10%) 5 5	50, 69, 103, 112	0
1	B	266/270 (98%)	0.73	29 (10%) 5 5	56, 75, 112, 131	0
1	C	230/270 (85%)	1.50	66 (28%) 0 0	75, 100, 122, 131	0
All	All	763/810 (94%)	0.92	124 (16%) 1 1	50, 80, 113, 131	0

All (124) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	182	LEU	7.2
1	C	203	THR	6.5
1	C	258	THR	6.2
1	B	257	ASP	6.1
1	A	260	LEU	5.8
1	C	250	SER	5.8
1	C	201	SER	5.6
1	C	216	ILE	5.4
1	C	175	LEU	5.3
1	C	257	ASP	5.1
1	C	267	VAL	4.9
1	C	211	PRO	4.7
1	A	172	GLY	4.6
1	C	17	VAL	4.6
1	A	258	THR	4.6
1	B	183	THR	4.6
1	C	184	ILE	4.6
1	C	113	ALA	4.6
1	B	166	GLU	4.5
1	C	256	MET	4.5
1	C	234	LYS	4.4
1	C	83	GLU	4.4
1	A	240	LYS	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	252	LEU	4.1
1	C	202	GLY	4.1
1	B	273	ARG	4.1
1	C	45	LEU	4.1
1	C	188	ASP	4.1
1	C	208	ASP	4.1
1	C	261	GLU	4.0
1	B	260	LEU	3.9
1	C	209	PRO	3.8
1	C	44	SER	3.8
1	B	169	GLY	3.7
1	C	162	HIS	3.7
1	C	65	VAL	3.6
1	B	261	GLU	3.6
1	B	181	PHE	3.5
1	C	183	THR	3.5
1	B	182	LEU	3.5
1	B	172	GLY	3.5
1	C	47	LEU	3.4
1	B	168	VAL	3.4
1	C	260	LEU	3.4
1	C	21	ILE	3.3
1	C	251	PHE	3.3
1	C	156	PHE	3.3
1	A	249	ALA	3.3
1	C	268	ASP	3.3
1	C	198	ASP	3.3
1	C	197	MET	3.3
1	C	161	TYR	3.3
1	C	74	LEU	3.3
1	A	261	GLU	3.3
1	B	8	MET	3.3
1	B	165	PRO	3.2
1	C	246	TYR	3.2
1	C	176	HIS	3.2
1	C	204	PHE	3.0
1	B	185	LEU	3.0
1	B	259	ASP	3.0
1	B	217	ASN	3.0
1	A	250	SER	2.9
1	A	8	MET	2.9
1	A	158	ILE	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	184	ILE	2.8
1	C	60	GLN	2.8
1	C	114	ASP	2.8
1	B	249	ALA	2.8
1	B	216	ILE	2.7
1	C	49	ALA	2.7
1	A	168	VAL	2.7
1	C	249	ALA	2.7
1	C	206	PRO	2.7
1	B	173	VAL	2.7
1	A	74	LEU	2.7
1	C	8	MET	2.7
1	A	169	GLY	2.6
1	A	185	LEU	2.6
1	A	193	GLY	2.6
1	C	130	ALA	2.6
1	B	240	LYS	2.6
1	C	255	PRO	2.6
1	A	162	HIS	2.6
1	C	92	GLY	2.6
1	A	259	ASP	2.6
1	A	247	SER	2.6
1	C	20	LYS	2.6
1	C	190	ASN	2.6
1	C	212	ASN	2.5
1	C	133	LEU	2.5
1	C	120	VAL	2.5
1	C	207	ILE	2.4
1	A	87	TYR	2.4
1	C	189	GLU	2.4
1	A	130	ALA	2.4
1	C	52	LYS	2.4
1	A	182	LEU	2.4
1	C	195	GLU	2.4
1	B	145	GLU	2.4
1	C	63	TYR	2.4
1	A	134	ALA	2.4
1	C	154	SER	2.4
1	A	257	ASP	2.3
1	C	185	LEU	2.3
1	C	259	ASP	2.3
1	C	217	ASN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	159	ASN	2.3
1	A	184	ILE	2.3
1	A	183	THR	2.3
1	B	162	HIS	2.2
1	A	248	ILE	2.2
1	B	218	LEU	2.2
1	A	91	LEU	2.2
1	B	256	MET	2.1
1	C	134	ALA	2.1
1	A	164	LYS	2.1
1	B	269	ALA	2.1
1	B	144	VAL	2.1
1	C	177	THR	2.1
1	C	28	GLU	2.1
1	B	251	PHE	2.0
1	A	137	LEU	2.0
1	B	158	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	B	301	1/1	0.69	0.47	84,84,84,84	0
3	AKG	A	302	10/10	0.87	0.24	76,83,85,86	0
2	MG	A	301	1/1	0.90	0.26	75,75,75,75	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers


as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around AKG A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.