



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 11:07 PM EDT

PDB ID : 6LIY  
Title : SeMet CRL Protein of Arabidopsis  
Authors : Wang, F.F.; Guan, K.L.; Sun, P.K.; Xing, W.M.  
Deposited on : 2019-12-13  
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 67%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 28%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 10px;">3%      67%      5%      28%</p>
1	B	269	<div style="display: flex; align-items: center;"> <div style="width: 6%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 64%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 27%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 10px;">6%      64%      9%      27%</p>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3412 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chromophore lyase CRL, chloroplastic.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	195	Total	C	N	O	S	Se	0	0	0
			1555	972	279	293	8	3			
1	B	196	Total	C	N	O	S	Se	0	0	0
			1549	967	278	293	8	3			

- Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	139	Total 139	O 139	0	0
3	B	145	Total 145	O 145	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.29Å 73.54Å 67.15Å 90.00° 97.16° 90.00°	Depositor
Resolution (Å)	32.19 – 1.76 32.19 – 1.76	Depositor EDS
% Data completeness (in resolution range)	97.1 (32.19-1.76) 97.1 (32.19-1.76)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.96 (at 1.76Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.207 , 0.244 0.207 , 0.244	Depositor DCC
$R_{free}$ test set	1844 reflections (5.18%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.7	Xtrriage
Anisotropy	0.544	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 44.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3412	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/1592	0.56	0/2152
1	B	0.44	0/1587	0.57	0/2149
All	All	0.44	0/3179	0.57	0/4301

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1555	0	1463	14	0
1	B	1549	0	1445	21	0
2	A	12	0	12	2	0
2	B	12	0	12	1	0
3	A	139	0	0	6	0
3	B	145	0	0	5	1
All	All	3412	0	2932	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:101:ARG:NH1	3:B:401:HOH:O	1.91	0.95
2:A:301:MES:O1S	3:A:401:HOH:O	1.98	0.82
1:B:143:ASP:OD1	3:B:402:HOH:O	2.04	0.75
1:B:133:ARG:CD	1:B:158:ASP:O	2.35	0.75
1:B:133:ARG:NE	1:B:159:ILE:O	2.22	0.71
1:B:133:ARG:HD3	1:B:158:ASP:O	1.90	0.71
1:A:139:LYS:HE3	3:A:438:HOH:O	1.92	0.68
1:B:117:CYS:SG	1:B:123:CYS:HB2	2.33	0.67
1:A:176:ARG:O	3:A:402:HOH:O	2.13	0.66
1:A:206:LYS:HD3	3:A:402:HOH:O	1.96	0.65
1:A:129:SER:HG	1:A:164:THR:HG1	1.41	0.64
1:A:86:MSE:SE	1:A:156:ILE:HD12	2.49	0.61
1:B:133:ARG:HD2	1:B:158:ASP:O	2.00	0.61
1:B:86:MSE:HG3	1:B:90:SER:HB2	1.87	0.56
1:B:54:ARG:HH22	1:B:82:PRO:HG3	1.72	0.55
1:B:133:ARG:HE	1:B:160:GLY:HA2	1.71	0.54
1:A:206:LYS:HD3	1:A:206:LYS:H	1.71	0.54
1:A:90:SER:OG	1:A:113:MSE:HE3	2.09	0.53
1:B:92:VAL:HG22	1:B:113:MSE:HG2	1.89	0.53
1:B:53:SER:O	1:B:57:THR:HG23	2.10	0.52
1:A:92:VAL:HG22	1:A:113:MSE:HG2	1.92	0.51
1:A:206:LYS:HE2	3:A:421:HOH:O	2.13	0.49
1:B:87:VAL:O	3:B:403:HOH:O	2.20	0.49
1:A:86:MSE:HB2	1:A:90:SER:HB3	1.94	0.48
1:A:53:SER:O	1:A:57:THR:HG23	2.14	0.48
1:A:75:ASN:HB3	1:A:100:TRP:CD1	2.50	0.46
1:B:86:MSE:CG	1:B:90:SER:HB2	2.46	0.44
1:B:219:GLU:HG3	3:B:476:HOH:O	2.18	0.43
1:A:86:MSE:HE3	1:A:86:MSE:HB3	1.94	0.43
2:B:301:MES:H32	2:B:301:MES:H82	1.46	0.43
1:B:50:ARG:O	1:B:54:ARG:HG2	2.19	0.42
1:B:133:ARG:HD3	1:B:158:ASP:C	2.39	0.42
1:A:101:ARG:NH1	3:A:405:HOH:O	2.43	0.42
1:B:206:LYS:HA	1:B:206:LYS:HD2	1.78	0.42
2:A:301:MES:H82	2:A:301:MES:H31	1.75	0.42
1:B:228:LYS:HG3	3:B:430:HOH:O	2.20	0.42
1:B:223:GLN:HB3	1:B:228:LYS:HE3	2.03	0.40
1:B:175:LYS:HB3	1:B:175:LYS:HE3	1.78	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:465:HOH:O	3:B:497:HOH:O[1_655]	2.17	0.03

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/269 (72%)	192 (100%)	1 (0%)	0	100	100
1	B	194/269 (72%)	192 (99%)	2 (1%)	0	100	100
All	All	387/538 (72%)	384 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/233 (71%)	165 (99%)	1 (1%)	86	79
1	B	164/233 (70%)	162 (99%)	2 (1%)	71	56
All	All	330/466 (71%)	327 (99%)	3 (1%)	78	67

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	206	LYS
1	B	88	ASP
1	B	133	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MES	A	301	-	12,12,12	1.92	2 (16%)	14,16,16	2.46	7 (50%)
2	MES	B	301	-	12,12,12	1.85	2 (16%)	14,16,16	3.33	7 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MES	A	301	-	-	4/6/14/14	0/1/1/1
2	MES	B	301	-	-	5/6/14/14	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	MES	C8-S	-5.87	1.69	1.77
2	B	301	MES	C8-S	-5.52	1.69	1.77
2	B	301	MES	O2S-S	2.45	1.52	1.45
2	A	301	MES	O2S-S	2.37	1.52	1.45

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	MES	O1S-S-C8	8.96	117.71	106.92
2	B	301	MES	O2S-S-C8	-4.58	101.39	106.92
2	A	301	MES	C5-N4-C3	3.70	117.16	108.83
2	B	301	MES	C2-C3-N4	-3.68	104.52	110.10
2	A	301	MES	C2-C3-N4	-3.53	104.75	110.10
2	A	301	MES	O1S-S-C8	3.38	110.98	106.92
2	B	301	MES	C5-N4-C3	3.36	116.38	108.83
2	A	301	MES	C6-C5-N4	-3.35	105.02	110.10
2	A	301	MES	C7-N4-C5	3.13	119.24	111.23
2	B	301	MES	O1-C6-C5	2.84	118.04	111.80
2	A	301	MES	O2S-S-C8	2.70	110.17	106.92
2	A	301	MES	O3S-S-C8	2.57	109.93	105.77
2	B	301	MES	C6-C5-N4	-2.57	106.21	110.10
2	B	301	MES	C7-N4-C3	2.56	117.79	111.23

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	MES	C7-C8-S-O1S
2	A	301	MES	C7-C8-S-O2S
2	A	301	MES	C7-C8-S-O3S
2	B	301	MES	C8-C7-N4-C3
2	B	301	MES	C7-C8-S-O1S
2	B	301	MES	C7-C8-S-O2S
2	B	301	MES	C7-C8-S-O3S
2	A	301	MES	C8-C7-N4-C5
2	B	301	MES	C8-C7-N4-C5

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	MES	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	MES	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	192/269 (71%)	0.38	9 (4%) 31 37	20, 29, 50, 56	0
1	B	193/269 (71%)	0.50	16 (8%) 11 14	21, 30, 51, 65	0
All	All	385/538 (71%)	0.44	25 (6%) 18 24	20, 30, 50, 65	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	42	SER	6.0
1	B	174	GLY	4.4
1	A	237	ALA	4.1
1	B	43	THR	3.7
1	B	173	ARG	3.6
1	A	174	GLY	3.5
1	B	54	ARG	3.4
1	B	140	ASN	3.2
1	A	192	TRP	3.0
1	A	172	ASP	2.8
1	B	176	ARG	2.7
1	B	88	ASP	2.5
1	B	147	ASP	2.4
1	A	190	ASN	2.4
1	A	121	LEU	2.3
1	B	123	CYS	2.3
1	B	89	GLY	2.3
1	A	119	LYS	2.2
1	A	193	ASN	2.2
1	B	157	GLY	2.2
1	B	50	ARG	2.2
1	B	146	LYS	2.2
1	B	87	VAL	2.1
1	B	145	PRO	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	107	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MES	B	301	12/12	0.83	0.21	29,36,38,42	0
2	MES	A	301	12/12	0.91	0.12	31,36,38,38	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.