

Full wwPDB EM Validation Report (i)

Jun 25, 2025 – 12:48 PM JST

PDB ID : 6LR4 / pdb 00006lr4

EMDB ID : EMD-0957

Title: Molecular basis for inhibition of human gamma-secretase by small molecule

Authors: Yang, G.; Zhou, R.; Guo, X.; Lei, J.; Shi, Y.

Deposited on : 2020-01-15

Resolution : 3.00 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev118

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4-5-2 with Phenix2.0rc1

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

 $MapQ \quad : \quad 1.9.13$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.44

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain							
1	A	709	84%		10% 6%					
2	В	467	59%	7%	34%					
3	С	265	87%		5% 8%					
4	D	143	61%	6%	33%					
5	Е	2		100%						
5	G	2	50%		50%					
5	Н	2		100%						
5	I	2	50%	name of the second	50%					

Continued on next page...



 $Continued\ from\ previous\ page...$

Г			T		-:					
	\mathbf{Mol}	Chain	Length	Quality of chain						
	5	T	9	4.000/						
	9	J		100%						
	6	E								
	O	Г)	60%	40%					



2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 10838 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Nicastrin.

Mol	Chain	Residues	Atoms				AltConf	Trace	
1	A	667	Total 5235	C 3321	N 890	O 1003	S 21	0	0

• Molecule 2 is a protein called Presenilin-1.

Mol	Chain	Residues	Atoms				AltConf	Trace	
2	В	308	Total 2444	C 1655	N 371	O 404	S 14	0	0

• Molecule 3 is a protein called Gamma-secretase subunit APH-1A.

Mol	Chain	Residues	Atoms				AltConf	Trace	
3	С	243	Total 1872	C 1254	N 299	O 315	S 4	0	0

• Molecule 4 is a protein called Gamma-secretase subunit PEN-2.

Mo	l Chain	Residues	Atoms					AltConf	Trace
4	D	96	Total 814	_	N 126	O 128	S 1	0	0

There are 43 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled			Reference
D	-41	MET	-	expression tag	UNP Q9NZ42
D	-40	ALA	-	expression tag	UNP Q9NZ42
D	-39	SER	-	expression tag	UNP Q9NZ42
D	-38	TRP	-	expression tag	UNP Q9NZ42
D	-37	SER	-	expression tag	UNP Q9NZ42
D	-36	HIS	-	expression tag	UNP Q9NZ42
D	-35	PRO	-	expression tag	UNP Q9NZ42
D	-34	GLN	-	expression tag	UNP Q9NZ42
D	-33	PHE	-	expression tag	UNP Q9NZ42

Continued on next page...



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
D	-32	GLU	-	expression tag	UNP Q9NZ42
D	-31	LYS	-	expression tag	UNP Q9NZ42
D	-30	GLY	-	expression tag	UNP Q9NZ42
D	-29	GLY	-	expression tag	UNP Q9NZ42
D	-28	GLY	-	expression tag	UNP Q9NZ42
D	-27	ALA	-	expression tag	UNP Q9NZ42
D	-26	ARG	-	expression tag	UNP Q9NZ42
D	-25	GLY	-	expression tag	UNP Q9NZ42
D	-24	GLY	-	expression tag	UNP Q9NZ42
D	-23	SER	-	expression tag	UNP Q9NZ42
D	-22	GLY	-	expression tag	UNP Q9NZ42
D	-21	GLY	-	expression tag	UNP Q9NZ42
D	-20	GLY	-	expression tag	UNP Q9NZ42
D	-19	SER	-	expression tag	UNP Q9NZ42
D	-18	TRP	-	expression tag	UNP Q9NZ42
D	-17	SER	-	expression tag	UNP Q9NZ42
D	-16	HIS	-	expression tag	UNP Q9NZ42
D	-15	PRO	-	expression tag	UNP Q9NZ42
D	-14	GLN	-	expression tag	UNP Q9NZ42
D	-13	PHE	-	expression tag	UNP Q9NZ42
D	-12	GLU	-	expression tag	UNP Q9NZ42
D	-11	LYS	-	expression tag	UNP Q9NZ42
D	-10	GLY	-	expression tag	UNP Q9NZ42
D	-9	PHE	-	expression tag	UNP Q9NZ42
D	-8	ASP	-	expression tag	UNP Q9NZ42
D	-7	TYR	-	expression tag	UNP Q9NZ42
D	-6	LYS	-	expression tag	UNP Q9NZ42
D	-5	ASP	-	expression tag	UNP Q9NZ42
D	-4	ASP	-	expression tag	UNP Q9NZ42
D	-3	ASP	-	expression tag	UNP Q9NZ42
D	-2	ASP	-	expression tag	UNP Q9NZ42
D	-1	LYS	-	expression tag	UNP Q9NZ42
D	0	GLY	-	expression tag	UNP Q9NZ42
D	1	THR	-	expression tag	UNP Q9NZ42


 \bullet Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.






Mol	Chain	Residues	Atoms	AltConf	Trace
5	E	2	Total C N O	0	0
9	<u> </u>	2	28 16 2 10	0	
5	G	2	Total C N O	0	0
5	G	2	28 16 2 10	U	
5	Н	2	Total C N O	0	0
	11	2	28 16 2 10	U	
5	Ţ	2	Total C N O	0	0
	1	2	28 16 2 10	U	U
5	Ţ	2	Total C N O	0	0
	J	2	28 16 2 10		

• Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-3)-[beta-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
6	F	5	Total 61	C 34	_	O 25	0	0


 \bullet Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $\rm C_8H_{15}NO_6).$





Mol	Chain	Residues	A		AltConf		
7	Δ	1	Total	С	N	О	0
'	Λ	1	14	8	1	5	U
7	A	1	Total	С	N	Ο	0
'	11	1	14	8	1	5	U
7	A	1	Total	С	N	Ο	0
'	71	1	14	8	1	5	0
7	A	1	Total	С	N	Ο	0
'	71	1	14	8	1	5	0
7	A	1	Total	С	N	Ο	0
'	11	1	14	8	1	5	
7	A	1	Total	С	N	O	0
'	11	1	14	8	1	5	


• Molecule 8 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: PC1) (formula: $C_{44}H_{88}NO_8P$).



Mol	Chain	Residues	Atoms		AltConf
Q	B	1	Total C N (ОР	0
	Б	1	37 27 1 8	8 1	U
Q	С	1	Total C N (ЭΡ	0
8	C	1	41 31 1 8	8 1	0


• Molecule 9 is (2S)-2-hydroxy-3-methyl-N-[(2S)-1-[[(5S)-3-methyl-4-oxo-2,5-dihydro-1H-3-be nzazepin-5-yl]amino]-1-oxopropan-2-yl]butanamide (CCD ID: ESF) (formula: $C_{19}H_{27}N_3O_4$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms			AltConf	
Q	B	1	Total	С	N	О	0
	ם ו		26	19	3	4	

 \bullet Molecule 10 is CHOLESTEROL (CCD ID: CLR) (formula: $\mathrm{C_{27}H_{46}O}).$



Mol	Chain	Residues	Atoms	AltConf
10	С	1	Total C O 28 27 1	0
10	С	1	Total C O 28 27 1	0
10	С	1	Total C O 28 27 1	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nicastrin Chain A: 84% PRO ARG GLU PRO GLY ALA VAL SER TYR • Molecule 2: Presenilin-1 Chain B: 59% 34% MINITER MANUAL M







• Molecule 5:	2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)	-2-acetamido-2-	-deoxy-beta-D-	gluc
opyranose				

Chain J: 100%



 $\bullet \ \, \text{Molecule 6: beta-D-mannopyranose-(1-3)-[beta-D-mannopyranose-(1-6)]} \\ \text{beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-acetamido-2$

Chain F: 60% 40%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	1044389	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	1.5625	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.170	Depositor
Minimum map value	-0.088	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.004	Depositor
Recommended contour level	0.028	Depositor
Map size (Å)	349.12, 349.12, 349.12	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.091, 1.091, 1.091	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CLR, BMA, NAG, ESF, PC1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Clasia	Bond lengths		Bond angles	
Mol Chain	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.29	0/5358	0.56	$4/7302 \ (0.1\%)$
2	В	0.28	0/2507	0.53	0/3421
3	С	0.27	0/1924	0.55	$2/2624 \ (0.1\%)$
4	D	0.25	0/847	0.51	0/1157
All	All	0.28	0/10636	0.55	6/14504 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers	
1	A	0	2	

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	${f Z}$	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
1	A	209	THR	CA-C-N	5.83	126.99	120.06
1	A	209	THR	C-N-CA	5.83	126.99	120.06
3	С	108	GLY	CA-C-N	5.33	131.73	121.54
3	С	108	GLY	C-N-CA	5.33	131.73	121.54
1	A	92	PRO	CA-C-N	-5.08	113.49	119.84
1	A	92	PRO	C-N-CA	-5.08	113.49	119.84

There are no chirality outliers.

All (2) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	224	VAL	Peptide
1	A	91	ASN	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	5235	0	5127	36	0
2	В	2444	0	2569	21	0
3	С	1872	0	1911	9	0
4	D	814	0	804	6	0
5	Е	28	0	25	0	0
5	G	28	0	25	0	0
5	Н	28	0	25	0	0
5	I	28	0	25	0	0
5	J	28	0	25	0	0
6	F	61	0	52	0	0
7	A	84	0	78	0	0
8	В	37	0	48	1	0
8	С	41	0	56	1	0
9	В	26	0	0	0	0
10	С	84	0	138	1	0
All	All	10838	0	10908	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:350:GLY:HA2	1:A:355:GLN:HE22	1.64	0.61
2:B:106:TYR:HA	2:B:239:LYS:HD3	1.83	0.61
1:A:530:ASN:OD1	1:A:535:GLN:NE2	2.35	0.59
1:A:218:PHE:O	1:A:656:ILE:HA	2.03	0.57
8:B:501:PC1:H351	3:C:135:ILE:HD11	1.86	0.57
1:A:256:VAL:HG22	1:A:329:PHE:HB2	1.88	0.56
1:A:54:LEU:HD11	1:A:223:ALA:HB1	1.88	0.55

Continued on next page...



Continued from previous page...

Continued from preva		Interatomic	Clash
Atom-1	Atom-2	${ m distance} \; ({ m \AA})$	overlap (Å)
2:B:173:LEU:HD23	2:B:229:ILE:HB	1.89	0.55
1:A:282:LEU:HD11	1:A:567:VAL:HG21	1.89	0.55
1:A:310:GLU:HB3	1:A:574:LEU:HB3	1.90	0.54
1:A:261:LYS:NZ	1:A:321:LEU:O	2.42	0.53
1:A:559:SER:HB2	1:A:629:ARG:HH12	1.74	0.53
1:A:210:PHE:HB3	1:A:665:SER:HB2	1.91	0.52
2:B:85:LEU:HD12	2:B:422:LEU:HD21	1.92	0.52
1:A:304:THR:HG23	1:A:365:LEU:HD21	1.91	0.52
2:B:140:ILE:HA	2:B:143:ILE:HG22	1.91	0.52
1:A:252:SER:HB3	1:A:631:LEU:HD13	1.92	0.52
2:B:192:ALA:O	4:D:79:GLN:NE2	2.37	0.51
1:A:665:SER:OG	1:A:667:GLU:OE1	2.29	0.51
4:D:41:ALA:HB2	4:D:53:ILE:HG22	1.93	0.51
1:A:120:GLY:H	1:A:178:PHE:HB2	1.75	0.50
2:B:452:LEU:O	3:C:50:SER:OG	2.29	0.50
8:C:304:PC1:H362	8:C:304:PC1:H2A1	1.93	0.50
1:A:537:ILE:HG22	1:A:569:TYR:HB3	1.92	0.50
1:A:543:ARG:HD3	1:A:606:GLN:HE22	1.76	0.50
1:A:401:LEU:HD11	1:A:439:VAL:HG11	1.94	0.50
2:B:96:VAL:HG21	2:B:390:SER:HB3	1.93	0.50
3:C:17:ALA:HB2	3:C:168:ILE:HG21	1.93	0.49
1:A:253:ASP:HB3	1:A:331:GLN:HG2	1.94	0.49
1:A:691:ASN:HD22	3:C:23:ILE:HG22	1.76	0.49
2:B:171:LEU:HD23	2:B:210:MET:HE3	1.94	0.49
1:A:137:SER:OG	1:A:165:ASN:ND2	2.46	0.49
1:A:364:GLU:OE2	1:A:427:LEU:N	2.47	0.48
1:A:296:GLU:HG2	1:A:369:ALA:HB3	1.95	0.48
4:D:34:ILE:HA	4:D:57:VAL:HG23	1.95	0.48
1:A:275:VAL:HG12	1:A:324:ASN:HB3	1.96	0.47
2:B:416:ILE:HG13	3:C:40:PHE:HB2	1.96	0.47
1:A:52:ARG:NH2	1:A:63:GLN:OE1	2.47	0.47
10:C:302:CLR:H222	10:C:302:CLR:H162	1.57	0.47
1:A:75:VAL:HG12	1:A:97:LEU:HD23	1.97	0.46
1:A:286:SER:OG	1:A:287:PHE:N	2.48	0.46
4:D:27:PRO:O	4:D:30:TRP:HB2	2.16	0.46
2:B:208:VAL:HG12	2:B:225:TYR:HE2	1.82	0.45
1:A:355:GLN:H	1:A:358:ASN:HD22	1.64	0.45
3:C:82:LEU:HA	3:C:85:VAL:HG12	1.98	0.45
1:A:64:SER:HB2	1:A:179:PRO:HG3	1.99	0.43
1:A:610:HIS:ND1	1:A:611:SER:O	2.50	0.43
2:B:211:ILE:HG23	2:B:215:TRP:HE3	1.83	0.43

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance} ({ m \AA})$	overlap (Å)
3:C:87:ARG:HH21	3:C:125:PHE:HD2	1.67	0.43
2:B:194:ASP:OD1	2:B:194:ASP:N	2.46	0.43
2:B:151:VAL:HG22	2:B:271:LEU:HD12	2.01	0.42
2:B:464:GLN:O	3:C:207:ASN:ND2	2.35	0.42
1:A:51:VAL:HA	1:A:221:MET:HE3	2.00	0.42
2:B:392:LEU:HD22	2:B:414:ILE:HD11	2.02	0.42
1:A:540:GLN:HA	1:A:606:GLN:HE21	1.84	0.42
1:A:589:PRO:HD2	1:A:597:LYS:HD2	2.02	0.42
1:A:34:ASN:ND2	3:C:140:ASP:OD1	2.53	0.41
2:B:194:ASP:OD2	4:D:74:TRP:NE1	2.50	0.41
2:B:192:ALA:HB2	4:D:97:PRO:HD3	2.03	0.41
1:A:112:LYS:NZ	1:A:175:ASP:O	2.45	0.41
1:A:317:ASP:OD1	1:A:320:THR:OG1	2.39	0.41
2:B:145:VAL:HA	2:B:148:ILE:HG22	2.03	0.41
2:B:226:LEU:HD22	2:B:383:LEU:HD11	2.03	0.41
1:A:517:THR:HG22	1:A:520:ARG:HH21	1.86	0.40
2:B:109:LYS:O	2:B:239:LYS:NZ	2.54	0.40
2:B:421:THR:HG21	2:B:434:ALA:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles		
1	A	665/709 (94%)	616 (93%)	49 (7%)	0	100	100	
2	В	304/467 (65%)	295 (97%)	9 (3%)	0	100	100	
3	C	241/265 (91%)	231 (96%)	10 (4%)	0	100	100	
4	D	94/143 (66%)	87 (93%)	7 (7%)	0	100	100	
All	All	1304/1584 (82%)	1229 (94%)	75 (6%)	0	100	100	

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles			
1	A	584/612 (95%)	583 (100%)	1 (0%)	92	97		
2	В	266/408~(65%)	266 (100%)	0	100	100		
3	С	193/214 (90%)	193 (100%)	0	100	100		
4	D	84/119 (71%)	84 (100%)	0	100	100		
All	All	1127/1353 (83%)	1126 (100%)	1 (0%)	92	98		

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	147	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	ASN
1	A	58	HIS
1	A	139	GLN
1	A	163	GLN
1	A	165	ASN
1	A	194	GLN
1	A	355	GLN
1	A	358	ASN
1	A	516	GLN
1	A	531	ASN
1	A	606	GLN
1	A	691	ASN
2	В	81	HIS
2	В	135	ASN
2	В	214	HIS
2	В	464	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

15 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N / - 1	Т	Clasica	Dag	Link	Во	ond leng	ths	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	Е	1	1,5	14,14,15	0.26	0	17,19,21	0.42	0
5	NAG	Е	2	5	14,14,15	0.25	0	17,19,21	0.62	0
6	NAG	F	1	1,6	14,14,15	0.24	0	17,19,21	0.55	0
6	NAG	F	2	6	14,14,15	0.25	0	17,19,21	0.56	0
6	BMA	F	3	6	11,11,12	0.66	0	15,15,17	0.85	1 (6%)
6	BMA	F	4	6	11,11,12	0.83	0	15,15,17	0.81	0
6	BMA	F	5	6	11,11,12	1.58	3 (27%)	15,15,17	1.86	4 (26%)
5	NAG	G	1	1,5	14,14,15	0.29	0	17,19,21	1.08	2 (11%)
5	NAG	G	2	5	14,14,15	0.38	0	17,19,21	0.52	0
5	NAG	Н	1	1,5	14,14,15	0.32	0	17,19,21	0.49	0
5	NAG	Н	2	5	14,14,15	0.38	0	17,19,21	0.62	0
5	NAG	I	1	1,5	14,14,15	0.26	0	17,19,21	0.50	0
5	NAG	I	2	5	14,14,15	0.82	1 (7%)	17,19,21	2.22	3 (17%)
5	NAG	J	1	1,5	14,14,15	0.36	0	17,19,21	0.49	0
5	NAG	J	2	5	14,14,15	0.27	0	17,19,21	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	Ε	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	E	2	5	-	2/6/23/26	0/1/1/1
6	NAG	F	1	1,6	-	4/6/23/26	0/1/1/1
6	NAG	F	2	6	-	2/6/23/26	0/1/1/1
6	BMA	F	3	6	-	2/2/19/22	0/1/1/1
6	BMA	F	4	6	-	1/2/19/22	0/1/1/1
6	BMA	F	5	6	-	0/2/19/22	0/1/1/1
5	NAG	G	1	1,5	-	1/6/23/26	0/1/1/1
5	NAG	G	2	5	-	2/6/23/26	0/1/1/1
5	NAG	Н	1	1,5	-	1/6/23/26	0/1/1/1
5	NAG	Н	2	5	-	4/6/23/26	0/1/1/1
5	NAG	I	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	I	2	5	-	5/6/23/26	0/1/1/1
5	NAG	J	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	J	2	5	-	0/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$Ideal(\AA)$
6	F	5	BMA	C2-C3	3.07	1.57	1.52
6	F	5	BMA	C1-C2	2.89	1.58	1.52
5	I	2	NAG	C1-C2	2.40	1.55	1.52
6	F	5	BMA	O5-C1	2.12	1.47	1.43

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
5	I	2	NAG	C2-N2-C7	7.79	134.00	122.90
6	F	5	BMA	C1-O5-C5	5.16	119.18	112.19
5	I	2	NAG	C1-C2-N2	3.52	116.50	110.49
5	G	1	NAG	C2-N2-C7	3.17	127.42	122.90
6	F	5	BMA	C1-C2-C3	2.88	113.21	109.67
6	F	5	BMA	O5-C1-C2	2.53	114.67	110.77
6	F	5	BMA	C2-C3-C4	2.16	114.62	110.89
5	I	2	NAG	C8-C7-N2	2.11	119.67	116.10
5	G	1	NAG	C1-O5-C5	2.01	114.92	112.19
6	F	3	BMA	C1-O5-C5	2.00	114.91	112.19

There are no chirality outliers.

All (28) torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
5	I	2	NAG	O5-C5-C6-O6
6	F	2	NAG	O5-C5-C6-O6
6	F	3	BMA	C4-C5-C6-O6
5	Н	2	NAG	O5-C5-C6-O6
5	I	2	NAG	C4-C5-C6-O6
6	F	3	BMA	O5-C5-C6-O6
6	F	2	NAG	C4-C5-C6-O6
5	J	1	NAG	O5-C5-C6-O6
5	Е	2	NAG	C8-C7-N2-C2
5	Е	2	NAG	O7-C7-N2-C2
5	G	2	NAG	C8-C7-N2-C2
5	G	2	NAG	O7-C7-N2-C2
5	Н	2	NAG	C8-C7-N2-C2
5	Н	2	NAG	O7-C7-N2-C2
5	I	2	NAG	C8-C7-N2-C2
5	I	2	NAG	O7-C7-N2-C2
6	F	1	NAG	C8-C7-N2-C2
6	F	1	NAG	O7-C7-N2-C2
5	J	1	NAG	C4-C5-C6-O6
5	Е	1	NAG	O5-C5-C6-O6
5	Н	2	NAG	C4-C5-C6-O6
5	Е	1	NAG	C4-C5-C6-O6
6	F	1	NAG	C4-C5-C6-O6
6	F	4	BMA	O5-C5-C6-O6
5	Н	1	NAG	O5-C5-C6-O6
6	F	1	NAG	O5-C5-C6-O6
5	G	1	NAG	C3-C2-N2-C7
5	I	2	NAG	C3-C2-N2-C7

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





























5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain Res		Link	Bond lengths			Bond angles			
MIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	NAG	A	818	1	14,14,15	0.42	0	17,19,21	0.44	0
7	NAG	A	819	1	14,14,15	0.98	1 (7%)	17,19,21	2.23	3 (17%)
10	CLR	С	302	-	31,31,31	0.33	0	48,48,48	0.65	0
10	CLR	С	303	-	31,31,31	0.31	0	48,48,48	0.51	0
8	PC1	С	304	-	40,40,53	0.30	0	46,48,61	0.38	0
7	NAG	A	821	1	14,14,15	0.36	0	17,19,21	0.56	0



Mol	Mol Type Chain R		Dog	Res Link	Bond lengths			Bond angles			
MIOI	Туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
9	ESF	В	502	-	26,27,27	0.76	1 (3%)	32,38,38	1.14	1 (3%)	
10	CLR	С	301	-	31,31,31	0.27	0	48,48,48	0.48	0	
8	PC1	В	501	-	36,36,53	0.33	0	42,44,61	0.49	0	
7	NAG	A	820	1	14,14,15	0.38	0	17,19,21	0.63	1 (5%)	
7	NAG	A	817	-	14,14,15	0.44	0	17,19,21	1.07	2 (11%)	
7	NAG	A	810	1	14,14,15	0.34	0	17,19,21	0.59	1 (5%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	A	818	1	-	2/6/23/26	0/1/1/1
7	NAG	A	819	1	-	5/6/23/26	0/1/1/1
10	CLR	С	302	-	-	5/10/68/68	0/4/4/4
10	CLR	С	303	-	-	4/10/68/68	0/4/4/4
8	PC1	С	304	-	-	6/44/44/57	-
7	NAG	A	821	1	-	1/6/23/26	0/1/1/1
9	ESF	В	502	-	-	4/20/37/37	0/2/2/2
10	CLR	С	301	-	-	4/10/68/68	0/4/4/4
8	PC1	В	501	-	-	5/40/40/57	-
7	NAG	A	820	1	-	0/6/23/26	0/1/1/1
7	NAG	A	817	-	-	3/6/23/26	0/1/1/1
7	NAG	A	810	1	-	1/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
7	A	819	NAG	C1-C2	3.11	1.57	1.52
9	В	502	ESF	C13-N05	-2.73	1.33	1.35

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
7	A	819	NAG	C2-N2-C7	7.70	133.86	122.90
9	В	502	ESF	C20-N05-C13	5.57	123.46	118.54
7	A	819	NAG	C1-C2-N2	3.80	116.98	110.49
7	A	817	NAG	C2-N2-C7	3.08	127.29	122.90

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^o)$
7	A	820	NAG	C1-O5-C5	2.28	115.29	112.19
7	A	817	NAG	C1-O5-C5	2.10	115.04	112.19
7	A	819	NAG	C8-C7-N2	2.04	119.56	116.10
7	A	810	NAG	C1-O5-C5	2.04	114.96	112.19

There are no chirality outliers.

All (40) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	В	501	PC1	C11-O13-P-O14
8	В	501	PC1	C11-O13-P-O11
8	В	501	PC1	O32-C31-O31-C3
8	В	501	PC1	C32-C31-O31-C3
8	С	304	PC1	C11-O13-P-O14
8	С	304	PC1	O22-C21-O21-C2
9	В	502	ESF	C18-C19-C23-N07
9	В	502	ESF	C18-C19-C23-O04
9	В	502	ESF	O02-C19-C23-N07
9	В	502	ESF	O02-C19-C23-O04
8	С	304	PC1	C22-C21-O21-C2
10	С	301	CLR	C16-C17-C20-C21
10	С	301	CLR	C13-C17-C20-C21
10	С	301	CLR	C13-C17-C20-C22
10	С	301	CLR	C16-C17-C20-C22
7	A	819	NAG	O5-C5-C6-O6
10	С	303	CLR	C13-C17-C20-C22
10	С	302	CLR	C13-C17-C20-C22
7	A	819	NAG	C4-C5-C6-O6
7	A	819	NAG	C8-C7-N2-C2
7	A	819	NAG	O7-C7-N2-C2
7	A	817	NAG	C4-C5-C6-O6
10	С	303	CLR	C16-C17-C20-C22
7	A	818	NAG	O5-C5-C6-O6
10	С	302	CLR	C16-C17-C20-C22
7	A	818	NAG	C4-C5-C6-O6
10	С	303	CLR	C16-C17-C20-C21
7	A	817	NAG	O5-C5-C6-O6
10	С	302	CLR	C16-C17-C20-C21
10	С	303	CLR	C13-C17-C20-C21
7	A	821	NAG	O5-C5-C6-O6
10	С	302	CLR	C13-C17-C20-C21
8	В	501	PC1	C12-C11-O13-P

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms
8	С	304	PC1	C12-C11-O13-P
7	A	810	NAG	C4-C5-C6-O6
8	С	304	PC1	O13-C11-C12-N
8	С	304	PC1	C11-O13-P-O11
10	С	302	CLR	C17-C20-C22-C23
7	A	817	NAG	C3-C2-N2-C7
7	A	819	NAG	C3-C2-N2-C7


There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	С	302	CLR	1	0
8	С	304	PC1	1	0
8	В	501	PC1	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-0957. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map






The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.028. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $66~\mathrm{nm^3}$; this corresponds to an approximate mass of $60~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



^{*}Reported resolution corresponds to spatial frequency of 0.333 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)


This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-0957 and PDB model 6LR4. Per-residue inclusion information can be found in section 3 on page 9.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.028 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.028).



9.4 Atom inclusion (i)



At the recommended contour level, 87% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.028) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8130	0.5510
A	0.8440	0.5680
В	0.7620	0.5350
С	0.8050	0.5420
D	0.8090	0.5240
Е	0.6430	0.5340
F	0.7540	0.5550
G	0.8210	0.4950
Н	0.7140	0.4550
I	0.5710	0.5070
J	0.8210	0.5510



