



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 30, 2023 – 04:08 PM JST

PDB ID : 6LR7
Title : Crystal structure of GFPuv complexed with the nanobody LaG16 at 1.67 Angstrom resolution
Authors : Zhang, Z.Y.; Ding, Y.; Hattori, M.
Deposited on : 2020-01-15
Resolution : 1.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.32.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.32.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	239	
2	B	131	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	CRO	A	301	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3068 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	232	1919	1225	322	366	6	0	10	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P42212
A	-1	PRO	-	expression tag	UNP P42212
A	0	HIS	-	expression tag	UNP P42212
A	301	CRO	SER	chromophore	UNP P42212
A	301	CRO	TYR	chromophore	UNP P42212
A	301	CRO	GLY	chromophore	UNP P42212
A	80	ARG	GLN	conflict	UNP P42212
A	99	SER	PHE	conflict	UNP P42212
A	153	THR	MET	conflict	UNP P42212
A	163	ALA	VAL	conflict	UNP P42212
A	206	LYS	ALA	conflict	UNP P42212
A	208	LEU	SER	conflict	UNP P42212

- Molecule 2 is a protein called Nanobody LaG16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	125	979	609	175	191	4	0	3	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	132	Total 132	O 132	0	0
3	B	38	Total 38	O 38	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein



- Molecule 2: Nanobody LaG16



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.37Å 41.78Å 81.71Å 90.00° 91.89° 90.00°	Depositor
Resolution (Å)	42.22 – 1.67 48.34 – 1.67	Depositor EDS
% Data completeness (in resolution range)	99.3 (42.22-1.67) 99.3 (48.34-1.67)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 1.67Å)	Xtrriage
Refinement program	PHENIX 1.12-2829	Depositor
R, R_{free}	0.185 , 0.226 0.185 , 0.225	Depositor DCC
R_{free} test set	1992 reflections (5.25%)	wwPDB-VP
Wilson B-factor (Å ²)	17.9	Xtrriage
Anisotropy	0.197	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.024 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3068	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/1971	0.59	0/2661
2	B	0.38	0/1004	0.62	0/1359
All	All	0.37	0/2975	0.60	0/4020

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1919	0	1881	18	0
2	B	979	0	964	7	0
3	A	132	0	0	1	0
3	B	38	0	0	1	0
All	All	3068	0	2845	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:CRO:HG11	1:A:220:LEU:HD21	1.60	0.82
1:A:301:CRO:HE1	1:A:148[B]:HIS:CE1	2.26	0.70
1:A:162:LYS:NZ	1:A:182[B]:TYR:OH	2.35	0.58
1:A:61:VAL:O	1:A:301:CRO:HG13	2.06	0.55
1:A:301:CRO:N2	1:A:301:CRO:HG12	2.24	0.52
2:B:11:ARG:HD3	2:B:19:LEU:CD1	2.39	0.51
1:A:301:CRO:OH	1:A:148[B]:HIS:HE1	1.95	0.49
1:A:301:CRO:OH	1:A:148[B]:HIS:CE1	2.67	0.48
1:A:203:THR:HG23	1:A:224:VAL:HG22	1.95	0.47
2:B:11:ARG:HD3	2:B:19:LEU:HD11	1.97	0.46
2:B:84:MET:HB3	2:B:87:LEU:HD21	1.97	0.45
1:A:143:TYR:CZ	1:A:209:LYS:HE2	2.52	0.45
1:A:301:CRO:N2	1:A:301:CRO:HD2	2.31	0.45
1:A:122:ARG:HG2	2:B:30:PHE:CZ	2.52	0.43
1:A:230:THR:HG22	1:A:231:HIS:ND1	2.32	0.43
2:B:19:LEU:HD12	2:B:19:LEU:HA	1.50	0.43
1:A:301:CRO:HE1	1:A:148[B]:HIS:HE1	1.80	0.43
2:B:58:ASN:ND2	3:B:201:HOH:O	2.41	0.42
1:A:126[A]:LYS:HE3	1:A:128:ILE:CG2	2.50	0.42
1:A:301:CRO:CE1	1:A:148[B]:HIS:CE1	3.01	0.42
1:A:148[B]:HIS:CE1	1:A:167:ILE:CD1	3.02	0.42
1:A:236:LEU:HD21	3:A:531:HOH:O	2.19	0.41
1:A:301:CRO:CE1	1:A:148[B]:HIS:HE1	2.34	0.40
2:B:11:ARG:HD3	2:B:19:LEU:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	237/239 (99%)	234 (99%)	3 (1%)	0	100 100
2	B	126/131 (96%)	126 (100%)	0	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	363/370 (98%)	360 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/208 (102%)	212 (100%)	0	100	100
2	B	106/107 (99%)	104 (98%)	2 (2%)	57	38
All	All	318/315 (101%)	316 (99%)	2 (1%)	84	79

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	11	ARG
2	B	73	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRO	A	301	1	23,23,24	3.59	5 (21%)	30,32,34	3.99	12 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	A	301	1	-	1/12/31/32	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	301	CRO	CB2-CA2	15.63	1.48	1.35
1	A	301	CRO	CA2-C2	-4.03	1.44	1.48
1	A	301	CRO	C1-N2	3.70	1.37	1.32
1	A	301	CRO	C2-N3	-3.11	1.32	1.39
1	A	301	CRO	O2-C2	2.55	1.28	1.23

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	301	CRO	O2-C2-CA2	-13.61	123.32	130.96
1	A	301	CRO	CA2-C2-N3	13.07	109.55	103.37
1	A	301	CRO	C2-N3-C1	-4.14	105.87	107.97
1	A	301	CRO	CA1-C1-N3	-4.12	119.80	124.75
1	A	301	CRO	CB2-CA2-C2	3.57	126.54	122.28
1	A	301	CRO	O3-C3-CA3	-3.42	116.08	126.39
1	A	301	CRO	C2-CA2-N2	-3.40	106.55	108.93
1	A	301	CRO	CA1-C1-N2	2.99	128.07	123.89
1	A	301	CRO	CG2-CB2-CA2	-2.96	126.31	129.94
1	A	301	CRO	CA3-N3-C2	2.58	129.73	123.80
1	A	301	CRO	CA3-N3-C1	-2.34	124.36	127.16
1	A	301	CRO	C1-CA1-N1	-2.27	106.28	109.96

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	301	CRO	C3-CA3-N3-C2

There are no ring outliers.

1 monomer is involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	301	CRO	10	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/239 (96%)	0.05	4 (1%) 70 74	8, 16, 30, 40	0
2	B	125/131 (95%)	0.48	10 (8%) 12 13	10, 22, 35, 52	0
All	All	356/370 (96%)	0.20	14 (3%) 39 42	8, 18, 33, 52	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	127	SER	6.2
2	B	28	ARG	3.7
1	A	191	GLY	3.4
1	A	232	GLY	3.3
2	B	15	ALA	3.0
2	B	126	VAL	2.8
2	B	17	ASP	2.8
2	B	14	GLN	2.6
2	B	88	GLU	2.5
1	A	148[A]	HIS	2.3
2	B	13	VAL	2.2
2	B	112	VAL	2.2
2	B	86	ASN	2.2
1	A	4	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	CRO	A	301	22/23	0.95	0.11	9,12,16,35	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.