

Full wwPDB X-ray Structure Validation Report (i)

Jun 11, 2024 – 02:52 PM EDT

PDB ID : 6MIV

Title: Crystal structure of the mCD1d/xxq (JJ300)/iNKTCR ternary complex

Authors : Zajonc, D.M.; Bitra, A.; Janssens, J.

Deposited on : 2018-09-20

Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36.2

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	Similar resolution $(\# \text{Entries, resolution range}(\text{\AA}))$
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	С	209	90%	6% •
2	D	241	95%	
3	A	285		% 5%
4	В	99	97%	
5	Е	2	100%	



 $Continued\ from\ previous\ page...$

Mol	Chain	Length		Quality of chain	
6	F	3	33%	67%	_



2 Entry composition (i)

There are 11 unique types of molecules in this entry. The entry contains 6914 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called T cell receptor alpha variable 11,T cell receptor alpha variable 11,T cell receptor alpha joining 18,Human nkt tcr alpha chain, CHIMERIC PROTEIN,Human nkt tcr alpha chain.

\mathbf{Mol}	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	С	201	Total 1518	C 944	N 261	O 305	S 8	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	0	MET	-	initiating methionine	UNP A0A0B4J1J9
С	113	ILE	-	linker	UNP A0A0B4J1J9

• Molecule 2 is a protein called Beta-chain, Tcell receptor chain, T cell receptor beta constant 2.

Mo	Chain	Residues		Ato	oms		ZeroOcc	AltConf	Trace	
2	D	239	Total	C 1174	N 333	O 357	S	0	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	95	ASP	-	linker	UNP A2NTY6
D	96	GLU	-	linker	UNP A2NTY6
D	97	GLY	-	linker	UNP A2NTY6
D	98	TYR	-	linker	UNP A2NTY6
D	130	ALA	ALA	linker	UNP A0N8J3
D	168	CYS	SER	variant	UNP A0A5B9
D	186	SER	CYS	variant	UNP A0A5B9

• Molecule 3 is a protein called Antigen-presenting glycoprotein CD1d1.



Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
3	A	271	Total	C	N 271	0	S	0	0	0
	11	211	2160	1379	371	397	13			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	280	HIS	-	expression tag	UNP A0A0R4J090
A	281	HIS	-	expression tag	UNP A0A0R4J090
A	282	HIS	-	expression tag	UNP A0A0R4J090
A	283	HIS	-	expression tag	UNP A0A0R4J090
A	284	HIS	-	expression tag	UNP A0A0R4J090
A	285	HIS	-	expression tag	UNP A0A0R4J090

• Molecule 4 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	D	98	Total	С	N	О	S	0	0	0
4	Б	90	789	503	133	146	7	0	0	U

• Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	E	2	Total	С	N	O	0	0	0
	2	_	28	16	2	10			O

• Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[al pha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	F	3	Total 38	C 22	N 2	O 14	0	0	0

• Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	С	1	Total Na 1 1	0	0
7	D	2	Total Na 2 2	0	0


 \bullet Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: $\mathrm{C_3H_8O_3}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	С	1	Total C O 6 3 3	0	0
8	D	1	Total C O 6 3 3	0	0
8	A	1	Total C O 6 3 3	0	0
8	A	1	Total C O 6 3 3	0	0


• Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total 14	C 8	N 1	O 5	0	0

• Molecule 10 is N-[(2S,3S,4R)-1-({4-O-[(4-tert-butylphenyl)methyl]-alpha-D-galactopyran osyl}oxy)-3,4-dihydroxyoctadecan-2-yl]hexacosanamide (three-letter code: JU1) (formula: $C_{61}H_{113}NO_9$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	Δ	1	Total	С	N	О	0	0
10	Λ	1	71	61	1	9	U	0

• Molecule 11 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	С	86	Total O 86 86	0	0
11	D	147	Total O 147 147	0	0
11	A	123	Total O 123 123	0	0
11	В	43	Total O 43 43	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: T cell receptor alpha variable 11,T cell receptor alpha variable 11,T cell receptor alpha joining 18,Human nkt tcr alpha chain, CHIMERIC PROTEIN,Human nkt tcr alpha chain



• Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose







 \bullet Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 33% 67%





4 Data and refinement statistics (i)

Property	Value	Source	
Space group	C 2 2 21	Depositor	
Cell constants	78.98Å 191.44Å 151.37Å	Donogitor	
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	41.54 - 2.05	Depositor	
resolution (A)	41.51 - 2.05	EDS	
% Data completeness	98.1 (41.54-2.05)	Depositor	
(in resolution range)	98.1 (41.51-2.05)	EDS	
R_{merge}	0.11	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	2.17 (at 2.05Å)	Xtriage	
Refinement program	REFMAC 5.8.0222	Depositor	
P.P.	0.185 , 0.219	Depositor	
R, R_{free}	0.193 , 0.226	DCC	
R_{free} test set	3613 reflections (5.10%)	wwPDB-VP	
Wilson B-factor (Å ²)	34.5	Xtriage	
Anisotropy	0.055	Xtriage	
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 39.3	EDS	
L-test for twinning ²	$ < L > = 0.50, < L^2> = 0.33$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
F_o, F_c correlation	0.96	EDS	
Total number of atoms	6914	wwPDB-VP	
Average B, all atoms $(Å^2)$	40.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, NA, NAG, JU1, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	\mathbf{angles}
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	С	0.39	0/1545	0.59	0/2103
2	D	0.40	0/1921	0.61	0/2618
3	A	0.38	0/2223	0.54	0/3024
4	В	0.35	0/815	0.58	0/1112
All	All	0.39	0/6504	0.58	0/8857

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	С	1518	0	1432	8	0
2	D	1870	0	1772	6	0
3	A	2160	0	2056	13	0
4	В	789	0	735	1	0
5	Е	28	0	25	0	0
6	F	38	0	34	0	0
7	С	1	0	0	0	0
7	D	2	0	0	0	0
8	A	12	0	16	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	С	6	0	8	0	0
8	D	6	0	8	0	0
9	A	14	0	13	0	0
10	A	71	0	0	0	0
11	A	123	0	0	0	0
11	В	43	0	0	1	0
11	С	86	0	0	0	0
11	D	147	0	0	0	0
All	All	6914	0	6099	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A.11	A	Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ ({\rm \AA})$	overlap (Å)
3:A:59:SER:H	3:A:62:GLN:HE21	1.43	0.67
1:C:22:GLN:OE1	1:C:74:THR:HG22	1.95	0.67
4:B:67:HIS:HD2	11:B:101:HOH:O	1.78	0.66
3:A:267:HIS:HD2	3:A:269:SER:OG	1.82	0.63
3:A:215:PRO:O	3:A:267:HIS:HE1	1.82	0.61
3:A:191:ALA:HA	3:A:209:HIS:O	2.06	0.56
3:A:59:SER:H	3:A:62:GLN:NE2	2.08	0.52
3:A:145:LEU:HB3	3:A:146:PRO:HD3	1.94	0.50
3:A:88:MET:HE3	3:A:146:PRO:HD3	1.96	0.48
3:A:168:CYS:HB3	3:A:169:PRO:HD3	1.97	0.47
3:A:219:TRP:CZ2	3:A:221:MET:HG3	2.50	0.46
3:A:267:HIS:CD2	3:A:269:SER:H	2.34	0.46
3:A:188:LYS:HA	3:A:269:SER:OG	2.17	0.45
1:C:126:LEU:HD12	1:C:126:LEU:N	2.31	0.45
2:D:78:LEU:HD12	2:D:78:LEU:N	2.32	0.45
1:C:37:LYS:HE2	1:C:39:ASP:OD1	2.18	0.44
2:D:208:GLN:HG3	2:D:231:ILE:HG23	2.00	0.44
1:C:203:PRO:O	1:C:204:SER:CB	2.67	0.43
1:C:157:TYR:O	1:C:178:ALA:HA	2.19	0.42
3:A:219:TRP:HB3	3:A:266:LYS:HB2	2.02	0.42
3:A:258:GLU:HB3	3:A:279:TRP:CD1	2.54	0.42
2:D:174:LEU:C	2:D:174:LEU:HD12	2.41	0.41
2:D:199:GLN:HA	2:D:239:ARG:O	2.19	0.41
1:C:32:HIS:CE1	2:D:98:TYR:CD1	3.08	0.41
1:C:125:GLN:C	1:C:126:LEU:HD12	2.41	0.41
		Continu	ed on nert nage



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	$egin{aligned} \operatorname{Clash} \ \operatorname{overlap}\ (\mbox{\normalfont\AA}) \end{aligned}$
2:D:190:ARG:CD	2:D:190:ARG:N	2.84	0.41
1:C:30:ASP:N	1:C:30:ASP:OD1	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	\mathbf{C}	197/209~(94%)	187 (95%)	10 (5%)	0	100	100
2	D	237/241 (98%)	233 (98%)	4 (2%)	0	100	100
3	A	$267/285 \ (94\%)$	263 (98%)	4 (2%)	0	100	100
4	В	96/99~(97%)	96 (100%)	0	0	100	100
All	All	797/834 (96%)	779 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	C	169/188 (90%)	169 (100%)	0	100	100
2	D	201/208 (97%)	199 (99%)	2 (1%)	76	75
3	A	232/249 (93%)	229 (99%)	3 (1%)	69	67



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
4	В	87/93 (94%)	86 (99%)	1 (1%)	73 73		
All	All	689/738 (93%)	683 (99%)	6 (1%)	78 79		

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	128	SER
2	D	168	CYS
3	A	170	LEU
3	A	203	HIS
3	A	204	ARG
4	В	70	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	117	GLN
2	D	230	GLN
3	A	7	ASN
3	A	62	GLN
3	A	227	GLN
3	A	267	HIS
4	В	2	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain		Res	Ros	Link	Bo	ond leng	ths	В	ond ang	les
Wor Type	Chain	Lilik		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2		
5	NAG	Е	1	3,5	14,14,15	0.56	0	17,19,21	0.83	1 (5%)	
5	NAG	Е	2	5	14,14,15	0.39	0	17,19,21	0.86	1 (5%)	
6	NAG	F	1	3,6	14,14,15	0.45	0	17,19,21	0.95	1 (5%)	
6	NAG	F	2	6	14,14,15	0.47	0	17,19,21	0.92	0	
6	FUC	F	3	6	10,10,11	0.59	0	14,14,16	0.97	1 (7%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	Е	1	3,5	-	0/6/23/26	0/1/1/1
5	NAG	Е	2	5	-	1/6/23/26	0/1/1/1
6	NAG	F	1	3,6	-	0/6/23/26	0/1/1/1
6	NAG	F	2	6	-	2/6/23/26	0/1/1/1
6	FUC	F	3	6	-	-	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
6	F	3	FUC	O5-C1-C2	-2.52	106.89	110.77
5	Е	1	NAG	O5-C5-C6	2.35	110.89	107.20
5	Е	2	NAG	C1-C2-N2	-2.32	106.52	110.49
6	F	1	NAG	O5-C5-C6	2.31	110.83	107.20

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	F	2	NAG	O5-C5-C6-O6
6	F	2	NAG	C4-C5-C6-O6
5	Е	2	NAG	O5-C5-C6-O6




There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain Res	Chain	Ros	Link	Во	Bond lengths			Bond angles		
MIOI		LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2			
8	GOL	A	308	-	5,5,5	0.43	0	5,5,5	0.26	0	
10	JU1	A	307	-	72,72,72	0.89	2 (2%)	83,87,87	0.92	3 (3%)	
8	GOL	С	302	-	5,5,5	0.24	0	5,5,5	0.29	0	



Mol Type	Chain	Res	Link	Bond lengths			В	ond ang	les	
MIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	GOL	A	309	-	5,5,5	0.28	0	5,5,5	0.24	0
8	GOL	D	302	-	5,5,5	0.36	0	5,5,5	0.66	0
9	NAG	A	301	3	14,14,15	0.57	0	17,19,21	1.31	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	A	308	-	-	4/4/4/4	-
10	JU1	A	307	-	-	12/69/89/89	0/2/2/2
8	GOL	С	302	-	-	2/4/4/4	-
8	GOL	A	309	-	-	3/4/4/4	-
8	GOL	D	302	-	-	2/4/4/4	-
9	NAG	A	301	3	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	${f Atoms}$	${f Z}$	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
10	A	307	JU1	CBA-CAX	-4.25	1.45	1.53
10	A	307	JU1	CAT-CAU	-3.25	1.43	1.50

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
9	A	301	NAG	C1-O5-C5	3.87	117.44	112.19
10	A	307	JU1	O4-CAT-CAU	3.33	117.57	109.91
10	A	307	JU1	CAN-NAO-CBE	-2.78	118.80	123.48
9	A	301	NAG	O5-C5-C6	2.58	111.25	107.20
10	A	307	JU1	CBN-CBO-CBP	-2.08	103.88	114.42

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	С	302	GOL	C1-C2-C3-O3
8	D	302	GOL	C1-C2-C3-O3
8	D	302	GOL	O2-C2-C3-O3
8	A	308	GOL	C1-C2-C3-O3



Continued from previous page...


Mol	Chain	Res	Type	Atoms
10	A	307	JU1	CCE-CCF-CCG-CCH
10	A	307	JU1	CCK-CCL-CCM-CCN
8	A	308	GOL	O1-C1-C2-C3
8	A	309	GOL	O1-C1-C2-C3
10	A	307	JU1	CBP-CBQ-CBR-CBS
10	A	307	JU1	CCG-CCH-CCI-CCJ
8	С	302	GOL	O2-C2-C3-O3
8	A	308	GOL	O1-C1-C2-O2
10	A	307	JU1	CCA-CCB-CCC-CCD
10	A	307	JU1	O6-C2-C3-O2
10	A	307	JU1	CBZ-CCA-CCB-CCC
10	A	307	JU1	CCB-CCC-CCD-CCE
8	A	308	GOL	O2-C2-C3-O3
10	A	307	JU1	CBK-CBL-CBM-CBN
10	A	307	JU1	CBX-CBY-CBZ-CCA
8	A	309	GOL	C1-C2-C3-O3
10	A	307	JU1	CCL-CCM-CCN-CCO
8	A	309	GOL	O1-C1-C2-O2
10	A	307	JU1	CAU-CAT-O4-C4

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$		$OWAB(A^2)$	Q<0.9
1	С	201/209 (96%)	0.14	4 (1%) 65	69	25, 37, 70, 92	0
2	D	239/241 (99%)	-0.12	2 (0%) 86	88	24, 34, 54, 71	0
3	A	271/285 (95%)	0.30	16 (5%) 22	24	25, 36, 75, 104	0
4	В	98/99 (98%)	0.06	3 (3%) 49	53	28, 42, 65, 90	0
All	All	809/834 (97%)	0.11	25 (3%) 49	53	24, 36, 68, 104	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	В	2	GLN	4.1
4	В	99	MET	3.9
3	A	200	ALA	3.8
1	С	132	SER	3.8
1	С	133	ASP	3.6
3	A	203	HIS	3.5
3	A	279	TRP	3.4
3	A	255	ALA	3.3
3	A	256	GLY	3.1
2	D	179	ALA	3.0
3	A	198	SER	2.8
3	A	227	GLN	2.6
1	С	117	GLN	2.6
3	A	199	SER	2.6
3	A	109	GLY	2.6
3	A	254	GLU	2.5
3	A	273	GLN	2.5
3	A	206	LEU	2.4
3	A	197	PRO	2.3
3	A	196	VAL	2.3
3	A	259	ALA	2.3



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	С	187	CYS	2.1
4	В	4	THR	2.1
2	D	181	ASN	2.1
3	A	276	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
5	NAG	Ε	2	14/15	0.83	0.27	61,68,74,78	0
6	FUC	F	3	10/11	0.85	0.16	54,56,59,61	0
6	NAG	F	2	14/15	0.94	0.16	44,53,57,59	0
5	NAG	E	1	14/15	0.96	0.10	38,42,46,54	0
6	NAG	F	1	14/15	0.96	0.11	36,39,45,51	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.









6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
9	NAG	A	301	14/15	0.73	0.25	55,68,73,74	0
8	GOL	A	308	6/6	0.78	0.19	63,65,67,68	0
8	GOL	D	302	6/6	0.84	0.23	42,57,61,62	0
8	GOL	С	302	6/6	0.90	0.22	46,55,59,63	0
7	NA	С	301	1/1	0.92	0.31	51,51,51,51	0
7	NA	D	303	1/1	0.92	0.30	70,70,70,70	0
8	GOL	A	309	6/6	0.94	0.18	50,60,61,66	0



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
10	JU1	A	307	71/71	0.95	0.16	25,38,45,47	0
7	NA	D	301	1/1	0.98	0.12	41,41,41,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers (i)

There are no such residues in this entry.

