



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 4, 2023 – 06:40 AM EDT

PDB ID : 6MM8
Title : Catalytic subunit of cAMP-dependent protein kinase A in complex with RyR2
K2879A, S2813D phosphomimetic (2699-2904) crystal form 2
Authors : van Petegem, F.; Haji-Ghassemi, O.
Deposited on : 2018-09-29
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	339	
2	D	209	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACY	C	402	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 5031 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-dependent protein kinase catalytic subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
1	C	326	2729	1775	457	484	2	11	0	8	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	12	SER	-	expression tag	UNP P05132
C	13	ASN	-	expression tag	UNP P05132
C	14	ALA	-	expression tag	UNP P05132

- Molecule 2 is a protein called Ryanodine receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	206	1701	1081	295	317	8	0	2	0

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	2696	SER	-	expression tag	UNP E9Q401
D	2697	ASN	-	expression tag	UNP E9Q401
D	2698	ALA	-	expression tag	UNP E9Q401
D	2813	ASP	SER	engineered mutation	UNP E9Q401
D	2879	ALA	LYS	engineered mutation	UNP E9Q401

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			7	4	3		


- Molecule 4 is ACETIC ACID (three-letter code: ACY) (formula: $C_2H_4O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter

code: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
5	C	1	62	20	12	24	6	0	1

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	C	1	4	2	2	0	0
6	C	1	4	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	D	1	Total Cl 1 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	C	342	Total O 342 342	0	0
8	D	161	Total O 161 161	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP-dependent protein kinase catalytic subunit alpha



- Molecule 2: Ryanodine receptor 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	179.64Å 67.86Å 58.53Å 90.00° 92.22° 90.00°	Depositor
Resolution (Å)	33.95 – 1.85 33.93 – 1.85	Depositor EDS
% Data completeness (in resolution range)	92.7 (33.95-1.85) 92.7 (33.93-1.85)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 1.85Å)	Xtrriage
Refinement program	REFMAC 5.8.0232	Depositor
R, R_{free}	0.181 , 0.221 0.190 , 0.229	Depositor DCC
R_{free} test set	2798 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtrriage
Anisotropy	0.044	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.023 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5031	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, SEP, ACY, ANP, CL, TPO, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.69	0/2794	0.77	0/3760
2	D	0.70	0/1749	0.73	0/2358
All	All	0.69	0/4543	0.76	0/6118

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2729	0	2743	26	0
2	D	1701	0	1681	8	0
3	C	7	0	10	2	0
4	C	4	0	3	3	0
4	D	4	0	3	1	0
5	C	62	0	26	2	0
6	C	16	0	24	2	0
6	D	4	0	6	0	0
7	D	1	0	0	0	0
8	C	342	0	0	7	0
8	D	161	0	0	0	0
All	All	5031	0	4496	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:403[C]:ANP:H3'	5:C:403[C]:ANP:O1A	1.71	0.91
1:C:135:ILE:HG21	4:C:402:ACY:CH3	2.07	0.84
1:C:101:PRO:HA	3:C:401:PEG:H32	1.74	0.70
1:C:53:SER:HA	8:C:661:HOH:O	1.91	0.70
1:C:25:ASP:HB3	8:C:663:HOH:O	1.92	0.69
1:C:135:ILE:CG2	4:C:402:ACY:CH3	2.77	0.61
1:C:194:ARG:HD2	8:C:525:HOH:O	2.00	0.61
1:C:252:SER:HA	8:C:760:HOH:O	2.03	0.58
2:D:2753:GLN:HB3	2:D:2756:MET:HG3	1.87	0.56
1:C:135:ILE:CG2	4:C:402:ACY:H1	2.38	0.53
2:D:2755:LEU:HD21	2:D:2766:GLU:OE1	2.09	0.53
1:C:94:ILE:CD1	1:C:188:ALA:HB3	2.41	0.51
1:C:253:GLY:HA2	1:C:275:VAL:HG11	1.92	0.51
1:C:105[B]:LYS:HE2	1:C:121:GLU:CD	2.31	0.50
1:C:94:ILE:HD13	1:C:188:ALA:HB3	1.93	0.50
2:D:2743:GLY:HA3	2:D:2752:ILE:HG23	1.93	0.50
2:D:2826:ASP:OD1	2:D:2828:SER:HB2	2.12	0.50
1:C:53:SER:CA	8:C:661:HOH:O	2.56	0.49
1:C:72:LYS:HD2	6:C:405:EDO:H12	1.95	0.49
1:C:336[B]:ARG:HH12	1:C:338:SEP:P	2.36	0.48
1:C:173:LEU:C	1:C:173:LEU:HD12	2.34	0.48
1:C:248:GLU:O	1:C:252:SER:HB3	2.14	0.48
1:C:42:GLN:HE22	6:C:406:EDO:C1	2.27	0.47
1:C:99:ASN:OD1	3:C:401:PEG:C1	2.62	0.47
1:C:105[B]:LYS:HE2	1:C:121:GLU:OE2	2.16	0.46
2:D:2859:LEU:HD22	2:D:2865:GLY:HA3	1.98	0.46
2:D:2755:LEU:CD2	2:D:2766:GLU:OE1	2.65	0.44
1:C:75:ASP:HA	1:C:115:ASN:HD22	1.83	0.44
1:C:53:SER:HB3	8:C:661:HOH:O	2.18	0.43
5:C:403[C]:ANP:O1A	5:C:403[C]:ANP:C3'	2.54	0.43
2:D:2712:ILE:HB	4:D:3001:ACY:H2	2.00	0.43
1:C:226:VAL:HG13	1:C:237:PRO:HD2	2.01	0.42
2:D:2707:THR:HB	2:D:2779:LYS:HB3	2.01	0.42
1:C:286[B]:ASN:OD1	8:C:501:HOH:O	2.22	0.41
1:C:58:MET:HE3	1:C:58:MET:HB3	1.97	0.41
1:C:253:GLY:CA	1:C:275:VAL:HG11	2.51	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	328/339 (97%)	319 (97%)	9 (3%)	0	100	100
2	D	206/209 (99%)	198 (96%)	8 (4%)	0	100	100
All	All	534/548 (97%)	517 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	290/297 (98%)	288 (99%)	2 (1%)	84	79
2	D	183/184 (100%)	179 (98%)	4 (2%)	52	36
All	All	473/481 (98%)	467 (99%)	6 (1%)	69	58

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	44	ASP
1	C	53	SER
2	D	2715	LYS
2	D	2753	GLN
2	D	2790	ARG
2	D	2791	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such

sidechains are listed below:

Mol	Chain	Res	Type
1	C	90	ASN
1	C	113	ASN
1	C	115	ASN
2	D	2709	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	C	197	1	8,10,11	0.92	0	10,14,16	0.89	0
1	SEP	C	338	1	8,9,10	0.71	0	8,12,14	1.73	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	C	197	1	-	1/9/11/13	-
1	SEP	C	338	1	-	4/5/8/10	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	338	SEP	O3P-P-OG	-3.34	97.84	106.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	338	SEP	O3P-P-O2P	2.37	116.70	107.64

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	338	SEP	N-CA-CB-OG
1	C	338	SEP	CA-CB-OG-P
1	C	338	SEP	CB-OG-P-O2P
1	C	338	SEP	CB-OG-P-O3P
1	C	197	TPO	O-C-CA-CB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	338	SEP	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	EDO	C	405	-	3,3,3	0.22	0	2,2,2	0.29	0
6	EDO	C	406	-	3,3,3	0.18	0	2,2,2	0.12	0
5	ANP	C	403[C]	-	29,33,33	1.27	4 (13%)	31,52,52	1.24	4 (12%)
4	ACY	C	402	-	3,3,3	0.66	0	3,3,3	1.32	0
4	ACY	D	3001	-	3,3,3	0.94	0	3,3,3	0.81	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ANP	C	403[B]	-	29,33,33	1.33	5 (17%)	31,52,52	1.25	3 (9%)
6	EDO	C	407	-	3,3,3	0.09	0	2,2,2	0.08	0
6	EDO	D	3002	-	3,3,3	0.32	0	2,2,2	0.13	0
6	EDO	C	404	-	3,3,3	0.25	0	2,2,2	0.20	0
3	PEG	C	401	-	6,6,6	0.27	0	5,5,5	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	C	405	-	-	0/1/1/1	-
6	EDO	C	406	-	-	1/1/1/1	-
5	ANP	C	403[C]	-	-	3/14/38/38	0/3/3/3
5	ANP	C	403[B]	-	-	6/14/38/38	0/3/3/3
6	EDO	C	407	-	-	0/1/1/1	-
6	EDO	D	3002	-	-	0/1/1/1	-
6	EDO	C	404	-	-	0/1/1/1	-
3	PEG	C	401	-	-	3/4/4/4	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	403[B]	ANP	PG-O1G	3.57	1.51	1.46
5	C	403[C]	ANP	PG-O1G	3.31	1.51	1.46
5	C	403[C]	ANP	PB-O1B	3.19	1.51	1.46
5	C	403[B]	ANP	PB-O1B	3.05	1.51	1.46
5	C	403[B]	ANP	PB-O3A	2.76	1.62	1.59
5	C	403[C]	ANP	PB-O3A	2.30	1.62	1.59
5	C	403[B]	ANP	PB-O2B	-2.13	1.51	1.56
5	C	403[C]	ANP	PB-O2B	-2.08	1.51	1.56
5	C	403[B]	ANP	PG-O3G	-2.00	1.51	1.56

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	403[C]	ANP	O2B-PB-O1B	4.29	118.91	109.92
5	C	403[B]	ANP	O2B-PB-O1B	4.05	118.41	109.92
5	C	403[C]	ANP	O2G-PG-O1G	-2.82	106.36	113.45
5	C	403[B]	ANP	O3G-PG-O1G	-2.79	106.45	113.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	403[C]	ANP	C5-C6-N6	2.58	124.28	120.35
5	C	403[B]	ANP	C5-C6-N6	2.55	124.23	120.35
5	C	403[C]	ANP	O3G-PG-O1G	-2.22	107.87	113.45

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	403[B]	ANP	PB-N3B-PG-O1G
5	C	403[B]	ANP	PA-O3A-PB-O1B
5	C	403[B]	ANP	PA-O3A-PB-O2B
5	C	403[C]	ANP	PB-N3B-PG-O1G
5	C	403[C]	ANP	PA-O3A-PB-O2B
5	C	403[B]	ANP	O4'-C4'-C5'-O5'
6	C	406	EDO	O1-C1-C2-O2
3	C	401	PEG	O2-C3-C4-O4
5	C	403[B]	ANP	C3'-C4'-C5'-O5'
5	C	403[C]	ANP	C4'-C5'-O5'-PA
3	C	401	PEG	O1-C1-C2-O2
5	C	403[B]	ANP	C4'-C5'-O5'-PA
3	C	401	PEG	C4-C3-O2-C2


There are no ring outliers.


6 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	405	EDO	1	0
6	C	406	EDO	1	0
5	C	403[C]	ANP	2	0
4	C	402	ACY	3	0
4	D	3001	ACY	1	0
3	C	401	PEG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	324/339 (95%)	-0.08	8 (2%) 57 56	11, 18, 45, 97	0
2	D	206/209 (98%)	0.48	29 (14%) 2 3	14, 33, 71, 94	0
All	All	530/548 (96%)	0.14	37 (6%) 16 15	11, 23, 62, 97	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	2745	ILE	7.3
2	D	2749	SER	7.1
1	C	318	PHE	7.1
1	C	15	VAL	6.4
2	D	2748	ASP	5.4
2	D	2796	SER	5.4
2	D	2746	TYR	5.3
1	C	18	PHE	4.8
2	D	2799	LEU	4.5
2	D	2750	SER	4.5
2	D	2744	GLU	4.3
2	D	2741	ILE	3.9
2	D	2797	MET	3.8
2	D	2747	SER	3.8
2	D	2829	ASN	3.7
2	D	2699	ASN	3.5
1	C	19	LEU	3.4
1	C	330	TYR	3.3
2	D	2737	ALA	3.3
2	D	2765	LYS	3.2
2	D	2752	ILE	3.2
1	C	253	GLY	3.1
1	C	14	ALA	2.9
2	D	2828	SER	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	2755	LEU	2.7
2	D	2709	ASN	2.7
2	D	2739	GLY	2.6
2	D	2795	ASP	2.5
2	D	2793[A]	GLU	2.5
2	D	2738	ASN	2.3
2	D	2740	TRP	2.3
2	D	2791	THR	2.2
2	D	2761	LEU	2.2
1	C	331	GLU	2.1
2	D	2764	GLU	2.1
2	D	2826	ASP	2.0
2	D	2794	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	C	338	10/11	0.94	0.10	27,32,39,41	0
1	TPO	C	197	11/12	0.99	0.06	17,18,19,20	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	C	407	4/4	0.66	0.23	45,50,50,53	0
5	ANP	C	403[C]	31/31	0.78	0.27	31,52,67,69	31
5	ANP	C	403[B]	31/31	0.78	0.27	37,60,67,70	31


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PEG	C	401	7/7	0.80	0.19	40,42,46,49	0
4	ACY	D	3001	4/4	0.87	0.29	48,53,54,57	0
6	EDO	C	406	4/4	0.91	0.16	50,51,52,52	0
4	ACY	C	402	4/4	0.93	0.22	26,41,41,50	0
6	EDO	D	3002	4/4	0.93	0.15	26,31,33,34	0
6	EDO	C	405	4/4	0.94	0.10	35,36,37,39	0
6	EDO	C	404	4/4	0.97	0.07	20,22,24,24	0
7	CL	D	3003	1/1	1.00	0.09	15,15,15,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.