

Full wwPDB X-ray Structure Validation Report (i)

Oct 26, 2024 – 01:08 PM EDT

PDB ID	:	6P8Z
Title	:	Crystal structure of human KRAS G12C covalently bound to an acryloylaze-
		tidine acetamide inhibitor
Authors	:	Mohr, C.
Deposited on	:	2019-06-08
Resolution	:	1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


Xtriage (Phenix) EDS buster-report Percentile statistics CCP4 Density-Fitness	:::::::::::::::::::::::::::::::::::::::	3.0 1.1.7 (2018) 20231227.v01 (using entries in the PDB archive December 27th 2023) 9.0.003 (Gargrove) 1.0.11
Density-Fitness Ideal geometry (proteins) Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)	:	Engh & Huber (2001)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	164625	2328 (1.66-1.66)
Clashscore	180529	2515 (1.66-1.66)
Ramachandran outliers	177936	2475(1.66-1.66)
Sidechain outliers	177891	2475 (1.66-1.66)
RSRZ outliers	164620	2328 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	183	6% 84% 5% •	10%
1	В	183	85% 5%	9%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 3036 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Λ	165	Total	С	Ν	0	S	0	0	0
	A	105	1313	823	227	258	5	0		
1	В	166	Total	С	Ν	Ο	S	0) 0	0
	D	100	1328	833	228	263	4			U

• Molecule 1 is a protein called GTPase KRas.

Chain	Residue	Modelled	Actual	Comment	Reference
А	-13	MET	-	expression tag	UNP P01116
А	-12	LYS	-	expression tag	UNP P01116
А	-11	HIS	-	expression tag	UNP P01116
А	-10	HIS	-	expression tag	UNP P01116
А	-9	HIS	-	expression tag	UNP P01116
А	-8	HIS	-	expression tag	UNP P01116
А	-7	HIS	-	expression tag	UNP P01116
А	-6	HIS	-	expression tag	UNP P01116
А	-5	HIS	-	expression tag	UNP P01116
А	-4	ASP	-	expression tag	UNP P01116
А	-3	GLU	-	expression tag	UNP P01116
А	-2	VAL	-	expression tag	UNP P01116
А	-1	ASP	-	expression tag	UNP P01116
А	0	GLY	-	expression tag	UNP P01116
А	12	CYS	GLY	variant	UNP P01116
А	51	SER	CYS	engineered mutation	UNP P01116
А	80	LEU	CYS	engineered mutation	UNP P01116
А	118	SER	CYS	engineered mutation	UNP P01116
В	-13	MET	-	expression tag	UNP P01116
В	-12	LYS	-	expression tag	UNP P01116
В	-11	HIS	-	expression tag	UNP P01116
В	-10	HIS	-	- expression tag	
В	-9	HIS	-	- expression tag	
В	-8	HIS	-	expression tag	UNP P01116
В	-7	HIS	_	expression tag	UNP P01116

There are 36 discrepancies between the modelled and reference sequences:



Chain

В

В

В

В

В

В

В

В

В

В

В

-	expression tag
-	expression tag
-	expression tag
-	expression tag

Comment

expression tag

expression tag

expression tag

variant

engineered mutation

engineered mutation

engineered mutation

Reference

UNP P01116

Continued from previous page...

Modelled

HIS

HIS

ASP

GLU

VAL

ASP

GLY

CYS

SER

LEU

SER

Residue

-6

-5

-4

-3

-2

-1

0

12

51

80

118

• Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Actual

-

-

-

GLY


CYS

CYS

CYS

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	2	Total Ca 2 2	0	0
2	В	2	Total Ca 2 2	0	0

• Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).




Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf
2	۸	1	Total	С	Ν	Ο	Р	0	0
Э	A	1	28	10	5	11	2	0	0



Continued from previous page...

Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf
3	В	1	Total		-	0	Р	0	0
			28	10	\mathbf{b}	11	2		

• Molecule 4 is 2-[5-chloro-2-cyclopropyl-3-(5-methoxy-3,4-dihydroisoquinoline-2(1H)-carbony l)-7-methyl-1H-indol-1-yl]-N-(1-propanoylazetidin-3-yl)acetamide (three-letter code: O5S) (formula: C₃₁H₃₅ClN₄O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
4	Λ	1	Total	С	Cl	Ν	Ο	0	0	
4	A	1	40	31	1	4	4	0	0	
4	В	1	Total	С	Cl	Ν	Ο	0	0	
4	D		40	31	1	4	4	0	0	


• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	147	Total O 147 147	0	0
5	В	108	Total O 108 108	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: GTPase KRas



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	39.76Å 65.39Å 62.60Å	Depositor
a, b, c, α , β , γ	90.00° 105.09° 90.00°	Depositor
Resolution (Å)	30.00 - 1.65	Depositor
Resolution (A)	30.00 - 1.65	EDS
% Data completeness	98.1 (30.00-1.65)	Depositor
(in resolution range)	98.1 (30.00-1.65)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	0.11	Depositor
$< I/\sigma(I) > 1$	1.55 (at 1.65 Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
D D.	0.229 , 0.269	Depositor
R, R_{free}	0.238 , 0.278	DCC
R_{free} test set	1814 reflections (4.84%)	wwPDB-VP
Wilson B-factor $(Å^2)$	23.1	Xtriage
Anisotropy	0.222	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33, 28.6	EDS
L-test for twinning ²	$ < L >=0.47, < L^2>=0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3036	wwPDB-VP
Average B, all atoms $(Å^2)$	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.65% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GDP, $\rm O5S$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain		lengths	Bond	angles
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.33	0/1332	0.57	0/1795
1	В	0.32	0/1349	0.58	0/1821
All	All	0.33	0/2681	0.58	0/3616

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1313	0	1303	14	0
1	В	1328	0	1309	4	0
2	А	2	0	0	0	0
2	В	2	0	0	0	0
3	А	28	0	12	0	0
3	В	28	0	12	0	0
4	А	40	0	0	0	0
4	В	40	0	0	0	0
5	А	147	0	0	2	0
5	В	108	0	0	0	0
All	All	3036	0	2636	18	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:A:97:ARG:HH21	1:A:111:MET:CE	1.96	0.79
1:A:41:ARG:HH21	1:A:41:ARG:HG2	1.53	0.73
1:A:97:ARG:HH21	1:A:111:MET:HE3	1.53	0.73
1:A:68:ARG:HA	1:A:71:TYR:CE2	2.24	0.72
1:B:161:ARG:O	1:B:165:LYS:HG3	1.94	0.67
1:B:49:GLU:OE2	1:B:164:ARG:NH2	2.35	0.59
1:A:111:MET:HE2	5:A:479:HOH:O	2.02	0.58
1:A:41:ARG:HH21	1:A:41:ARG:CG	2.18	0.54
1:A:150:GLN:NE2	5:A:403:HOH:O	2.41	0.53
1:A:97:ARG:HH21	1:A:111:MET:HE1	1.74	0.52
1:A:97:ARG:NH2	1:A:111:MET:CE	2.72	0.48
1:B:22:GLN:HG3	1:B:146:ALA:O	2.14	0.47
1:A:41:ARG:CG	1:A:41:ARG:NH2	2.77	0.45
1:B:127:THR:HG22	1:B:143:GLU:OE2	2.17	0.45
1:A:68:ARG:HA	1:A:71:TYR:CD2	2.54	0.43
1:A:97:ARG:NH2	1:A:111:MET:HE1	2.34	0.42
1:A:3:GLU:HG2	1:A:52:LEU:HB3	2.02	0.42
1:A:24:ILE:HD13	1:A:40:TYR:HB3	2.02	0.42

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percenti	les
1	А	161/183~(88%)	158 (98%)	3~(2%)	0	100 10	00
1	В	164/183~(90%)	160 (98%)	4 (2%)	0	100 10	00
All	All	325/366~(89%)	318 (98%)	7(2%)	0	100 10	00



There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	145/163~(89%)	143~(99%)	2(1%)	62 43		
1	В	147/163~(90%)	145~(99%)	2(1%)	62 43		
All	All	292/326~(90%)	288~(99%)	4 (1%)	62 43		

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	41	ARG
1	А	67	MET
1	В	51	SER
1	В	71	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	99	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.



5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	ol Type Chain		Res	Link	В	ond leng	gths	Bond angles							
10101	Type	Unam	nes	nes	nes	nes	nes	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
3	GDP	А	303	2	$25,\!30,\!30$	0.97	2 (8%)	30,47,47	1.08	3 (10%)					
4	O5S	В	304	1	39,45,45	2.84	17 (43%)	53,67,67	2.83	13 (24%)					
3	GDP	В	303	2	25,30,30	1.08	1 (4%)	30,47,47	1.10	3 (10%)					
4	O5S	А	304	1	$39,\!45,\!45$	2.79	16 (41%)	53,67,67	2.83	9 (16%)					

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	А	303	2	-	0/12/32/32	0/3/3/3
4	O5S	В	304	1	-	3/21/47/47	0/6/6/6
3	GDP	В	303	2	-	1/12/32/32	0/3/3/3
4	O5S	А	304	1	-	2/21/47/47	0/6/6/6

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
4	В	304	O5S	C18-C19	7.79	1.57	1.51
4	А	304	O5S	C18-C19	7.39	1.57	1.51
4	А	304	O5S	C19-N21	6.61	1.48	1.34
4	В	304	O5S	C19-N21	6.58	1.48	1.34
4	В	304	O5S	C10-N09	6.31	1.47	1.34
4	В	304	O5S	C03-N05	6.06	1.46	1.35
4	А	304	O5S	C10-N09	5.98	1.46	1.34
4	А	304	O5S	C03-N05	5.79	1.46	1.35
4	А	304	O5S	C08-N05	4.07	1.50	1.47
4	В	304	O5S	C06-N05	4.07	1.50	1.47
4	В	304	O5S	C08-N05	3.87	1.50	1.47



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	В	304	O5S	C23-C24	3.65	1.57	1.51
4	А	304	O5S	C23-C24	3.55	1.57	1.51
4	В	304	O5S	O26-C25	3.47	1.42	1.37
3	В	303	GDP	PA-O3A	3.46	1.63	1.59
4	А	304	O5S	O26-C25	3.40	1.42	1.37
4	А	304	O5S	C06-N05	3.31	1.49	1.47
4	А	304	O5S	C30-C31	2.80	1.44	1.39
4	А	304	O5S	C25-C24	2.68	1.44	1.40
4	В	304	O5S	C25-C24	2.58	1.44	1.40
4	В	304	O5S	C30-C31	2.52	1.43	1.39
4	А	304	O5S	C14-C15	2.50	1.55	1.51
4	А	304	O5S	C34-C35	2.49	1.41	1.36
4	В	304	O5S	C02-C03	2.45	1.55	1.51
3	А	303	GDP	PA-O3A	2.44	1.62	1.59
4	А	304	O5S	C02-C03	2.43	1.55	1.51
4	В	304	O5S	C34-C35	2.36	1.40	1.36
4	В	304	O5S	C14-C15	2.28	1.55	1.51
4	А	304	O5S	C22-C23	2.24	1.56	1.51
4	В	304	O5S	C22-C23	2.13	1.55	1.51
4	А	304	O5S	C37-C38	2.08	1.40	1.37
4	В	304	O5S	C37-C38	2.07	1.40	1.37
4	В	304	O5S	C38-C40	2.05	1.45	1.43
4	А	304	O5S	C33-C40	-2.01	1.37	1.41
4	В	304	O5S	C37-C35	2.01	1.41	1.38
3	А	303	GDP	C6-N1	-2.01	1.34	1.37

Continued from previous page...

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
4	А	304	O5S	C17-C15-C14	16.99	138.89	119.66
4	В	304	O5S	C17-C15-C14	15.10	136.75	119.66
4	В	304	O5S	C08-N05-C06	-6.64	90.04	94.69
4	А	304	O5S	C08-N05-C06	-5.89	90.57	94.69
4	В	304	O5S	C23-C22-N21	-5.39	104.47	110.03
4	В	304	O5S	C18-C14-C15	-4.39	122.70	129.87
4	А	304	O5S	C16-C15-C14	-4.36	114.72	119.66
4	В	304	O5S	O26-C25-C24	3.62	119.03	115.30
4	А	304	O5S	C06-C07-N09	-3.59	107.12	116.59
4	А	304	O5S	O26-C25-C24	3.47	118.89	115.30
4	В	304	O5S	C31-C32-N21	-3.22	106.69	111.56
3	В	303	GDP	C8-N7-C5	2.94	107.55	102.55
3	А	303	GDP	C8-N7-C5	2.86	107.42	102.55



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	А	304	O5S	C18-C14-C15	-2.80	125.30	129.87
4	В	304	O5S	C16-C15-C14	-2.67	116.64	119.66
4	А	304	O5S	C23-C22-N21	-2.64	107.30	110.03
4	В	304	O5S	C07-N09-C10	-2.48	119.22	122.88
4	В	304	O5S	O11-C10-N09	-2.26	119.13	122.95
3	В	303	GDP	C5-C6-N1	2.22	118.30	114.07
3	А	303	GDP	C5-C6-N1	2.20	118.27	114.07
3	В	303	GDP	O6-C6-C5	-2.14	120.08	124.32
4	А	304	O5S	C34-C33-C40	2.13	121.70	117.07
4	В	304	O5S	C08-C07-N09	-2.13	110.99	116.59
3	А	303	GDP	O6-C6-C5	-2.10	120.15	124.32
4	В	304	O5S	C34-C33-C40	2.06	121.54	117.07
4	В	304	O5S	O20-C19-N21	-2.04	119.14	122.35
4	В	304	O5S	C25-C24-C31	2.02	120.59	118.35
4	А	304	O5S	O11-C10-N09	-2.00	119.56	122.95

Continued from previous page...

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	А	304	O5S	N13-C14-C15-C16
4	В	304	O5S	C01-C02-C03-N05
4	В	304	O5S	N13-C14-C15-C16
3	В	303	GDP	PA-O3A-PB-O2B
4	В	304	O5S	C01-C02-C03-O04
4	А	304	O5S	C28-C25-O26-C27

There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz> $#$RSRZ>2</rsrz>		$OWAB(Å^2)$	Q<0.9
1	А	165/183~(90%)	0.67	11 (6%) 25 28	17, 26, 46, 60	0
1	В	166/183~(90%)	1.15	30 (18%) 4 5	22, 34, 51, 58	0
All	All	331/366~(90%)	0.91	41 (12%) 9 10	17, 30, 51, 60	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	69	ASP	4.3
1	В	29	VAL	4.1
1	В	120	LEU	3.9
1	А	0	GLY	3.6
1	А	71	TYR	3.5
1	А	66	ALA	3.4
1	В	24	ILE	3.4
1	А	103	VAL	3.2
1	В	28	PHE	3.1
1	В	50	THR	3.1
1	В	127	THR	3.0
1	В	64	TYR	2.9
1	В	46	ILE	2.8
1	В	148	THR	2.7
1	В	124	THR	2.7
1	В	36	ILE	2.7
1	А	60	GLY	2.7
1	А	1	MET	2.5
1	В	48	GLY	2.5
1	В	71	TYR	2.5
1	В	149	ARG	2.5
1	В	26	ASN	2.4
1	В	31	GLU	2.4
1	B	122	SER	2.4



Mol	Chain	Res	Type	RSRZ
1	А	68	ARG	2.4
1	А	67	MET	2.4
1	А	165	LYS	2.4
1	В	30	ASP	2.3
1	В	69	ASP	2.3
1	В	49	GLU	2.3
1	В	72	MET	2.3
1	В	60	GLY	2.3
1	В	91	GLU	2.3
1	В	125	VAL	2.2
1	В	121	PRO	2.2
1	В	4	TYR	2.1
1	В	39	SER	2.1
1	В	157	TYR	2.1
1	В	45	VAL	2.0
1	В	134	ALA	2.0
1	А	105	ASP	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{\AA}^2)$	Q<0.9
2	CA	А	302	1/1	0.88	0.20	39,39,39,39	0
4	O5S	А	304	40/40	0.91	0.10	$26,\!29,\!31,\!32$	0
4	O5S	В	304	40/40	0.92	0.10	26,28,30,30	0
2	CA	В	302	1/1	0.94	0.10	44,44,44,44	0
3	GDP	В	303	28/28	0.94	0.09	27,32,34,36	0
3	GDP	А	303	28/28	0.97	0.06	20,21,24,26	0




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
2	CA	В	301	1/1	0.98	0.04	30,30,30,30	0
2	CA	А	301	1/1	0.98	0.05	21,21,21,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















6.5 Other polymers (i)

There are no such residues in this entry.

