



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 25, 2024 – 11:27 PM EDT

PDB ID : 6R7T  
Title : Crystal Structure of human Melanoma-associated antigen B1 (MAGEB1) in complex with nanobody  
Authors : Ye, M.; Newman, J.; Pike, A.C.W.; Burgess-Brown, N.; Cooper, C.D.O.; Bountra, C.; Arrowsmith, C.; Edwards, A.; Gileadi, O.; von Delft, F.  
Deposited on : 2019-03-29  
Resolution : 2.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3863 (2.70-2.66)
Clashscore	141614	4210 (2.70-2.66)
Ramachandran outliers	138981	4141 (2.70-2.66)
Sidechain outliers	138945	4141 (2.70-2.66)
RSRZ outliers	127900	3780 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	123	<p>4% (poor fit), 91% (0-1 outliers), 8% (2-3 outliers)</p>
2	B	251	<p>9% (poor fit), 60% (0-1 outliers), 11% (2 outliers), 28% (3+ outliers or not modelled)</p>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2442 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called anti MAGEB1 nanobody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	123	926	576	163	183	4	0	1	0

- Molecule 2 is a protein called Melanoma-associated antigen B1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	180	1473	959	243	262	9	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	99	SER	-	expression tag	UNP P43366
B	100	MET	-	expression tag	UNP P43366
B	146	GLU	ASP	conflict	UNP P43366
B	147	TYR	HIS	conflict	UNP P43366


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	16	Total	O	0	0
			16	16		
3	B	27	Total	O	0	0
			27	27		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: anti MAGEB1 nanobody



- Molecule 2: Melanoma-associated antigen B1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.98Å 61.98Å 361.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	53.67 – 2.68 53.67 – 2.68	Depositor EDS
% Data completeness (in resolution range)	99.5 (53.67-2.68) 99.6 (53.67-2.68)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.67 (at 2.69Å)	Xtrriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, $R_{free}$	0.258 , 0.288 0.259 , 0.288	Depositor DCC
$R_{free}$ test set	625 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	66.6	Xtrriage
Anisotropy	0.014	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 58.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	2442	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/948	0.47	0/1283
2	B	0.25	0/1507	0.40	0/2028
All	All	0.25	0/2455	0.43	0/3311

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	926	0	883	8	0
2	B	1473	0	1447	20	0
3	A	16	0	0	1	1
3	B	27	0	0	3	1
All	All	2442	0	2330	26	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:199:ARG:NH1	3:B:402:HOH:O	2.11	0.70

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:LYS:O	3:A:201:HOH:O	2.10	0.69
2:B:252:VAL:O	3:B:403:HOH:O	2.12	0.67
2:B:161:LEU:HD23	2:B:294:MET:HB2	1.82	0.61
2:B:218:THR:HG22	2:B:220:GLU:H	1.65	0.60
2:B:165:LEU:HD23	2:B:179:LEU:HD23	1.87	0.56
1:A:84:MET:HB3	1:A:87:LEU:HD21	1.89	0.54
1:A:17:SER:HA	1:A:84:MET:O	2.09	0.52
2:B:195:TRP:N	3:B:408:HOH:O	2.44	0.50
2:B:274:TRP:CZ2	2:B:287:VAL:HG11	2.47	0.50
1:A:84:MET:HE2	1:A:87:LEU:HD21	1.97	0.47
2:B:110:VAL:HG23	2:B:151:ILE:HG12	1.96	0.47
1:A:68:ARG:HB3	1:A:85:ASN:O	2.15	0.46
2:B:274:TRP:HZ2	2:B:287:VAL:HG11	1.81	0.45
2:B:132:MET:HA	2:B:176:THR:HA	1.98	0.44
1:A:110:ARG:HD3	2:B:195:TRP:CE2	2.53	0.43
2:B:256:TYR:CE1	2:B:277:ARG:HD2	2.54	0.43
2:B:298:THR:OG1	2:B:301:ASP:OD1	2.24	0.43
1:A:95:TYR:O	1:A:113:GLY:HA2	2.19	0.42
2:B:132:MET:HG2	2:B:135:ASP:HB2	2.00	0.42
2:B:255:LYS:HB3	2:B:276:PRO:HG2	2.02	0.41
1:A:103:ARG:NH2	2:B:201:GLY:O	2.39	0.41
2:B:212:LEU:HD13	2:B:291:LEU:HD21	2.02	0.41
2:B:256:TYR:CD1	2:B:277:ARG:HB3	2.55	0.41
2:B:122:LEU:HD23	2:B:122:LEU:HA	1.92	0.40
2:B:222:ILE:O	2:B:226:MET:HG2	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:216:HOH:O	3:B:426:HOH:O[8_555]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	122/123 (99%)	121 (99%)	1 (1%)	0	100	100
2	B	170/251 (68%)	162 (95%)	7 (4%)	1 (1%)	25	47
All	All	292/374 (78%)	283 (97%)	8 (3%)	1 (0%)	41	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	166	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	95/95 (100%)	93 (98%)	2 (2%)	53	78
2	B	151/222 (68%)	147 (97%)	4 (3%)	46	73
All	All	246/317 (78%)	240 (98%)	6 (2%)	49	75

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	ARG
1	A	122	LYS
2	B	110	VAL
2	B	132	MET
2	B	209	VAL
2	B	277	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.



### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	123/123 (100%)	0.26	5 (4%) 37 35	33, 50, 79, 100	0
2	B	180/251 (71%)	0.75	22 (12%) 4 3	38, 69, 107, 113	0
All	All	303/374 (81%)	0.55	27 (8%) 9 8	33, 58, 102, 113	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	120	SER	4.4
2	B	168	LYS	3.8
2	B	131	ILE	3.7
2	B	153	ASN	3.6
2	B	166	ASP	3.5
2	B	214	GLY	3.3
2	B	269	ARG	3.3
2	B	180	VAL	3.2
2	B	157	ARG	3.1
2	B	291	LEU	3.1
1	A	121	SER	3.1
2	B	116	MET	2.8
2	B	295	ASN	2.7
2	B	137	LEU	2.6
2	B	130	PRO	2.4
2	B	165	LEU	2.4
2	B	134	ALA	2.4
1	A	5	GLN	2.4
1	A	54	SER	2.4
2	B	114	ALA	2.3
2	B	139	VAL	2.3
1	A	1	GLN	2.3
2	B	160	GLU	2.2
2	B	149	THR	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	155	ALA	2.1
2	B	318	GLN	2.0
2	B	133	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.