



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2025 – 06:24 pm BST

PDB ID : 6SAK / pdb_00006sak
Title : Structure of the OTULINcat C129A - SNX27 PDZ domain complex.
Authors : Elliott, P.R.; Komander, D.
Deposited on : 2019-07-17
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	8% 88% 12%
1	B	275	7% 87% 9% .
2	C	96	36% 84% 15% .
2	D	96	7% 89% 11%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11628 atoms, of which 5602 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin thioesterase otulin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	274	Total	C	H	N	O	S	0	4	0
			4295	1401	2107	363	410	14			
1	B	264	Total	C	H	N	O	S	0	3	0
			4153	1355	2046	347	392	13			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	78	GLY	-	expression tag	UNP Q96BN8
A	79	PRO	-	expression tag	UNP Q96BN8
A	129	ALA	CYS	engineered mutation	UNP Q96BN8
B	78	GLY	-	expression tag	UNP Q96BN8
B	79	PRO	-	expression tag	UNP Q96BN8
B	129	ALA	CYS	engineered mutation	UNP Q96BN8

- Molecule 2 is a protein called Sorting nexin-27.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	95	Total	C	H	N	O		0	0	0
			1305	414	642	120	129				
2	D	96	Total	C	H	N	O		0	2	0
			1454	448	738	137	131				

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			13	3	7	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			12	3	6	3		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	106	Total	O	0	0
			106	106		
5	B	114	Total	O	0	0
			114	114		
5	C	16	Total	O	0	0
			16	16		
5	D	32	Total	O	0	0
			32	32		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ubiquitin thioesterase otulin




- Molecule 1: Ubiquitin thioesterase otulin



- Molecule 2: Sorting nexin-27



- Molecule 2: Sorting nexin-27





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.65Å 62.91Å 72.02Å 65.13° 85.98° 85.38°	Depositor
Resolution (Å)	42.65 – 2.00 42.65 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.5 (42.65-2.00) 95.6 (42.65-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.87 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.16_3549, PHENIX 1.16_3549	Depositor
R, R_{free}	0.215 , 0.253 0.214 , 0.250	Depositor DCC
R_{free} test set	2790 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtriage
Anisotropy	0.568	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 56.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.003 for -h,-k,-k+l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11628	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.13	0/2247	0.33	0/3054
1	B	0.12	0/2163	0.30	0/2942
2	C	0.24	0/670	0.53	2/913 (0.2%)
2	D	0.16	0/730	0.39	0/987
All	All	0.15	0/5810	0.36	2/7896 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	126	LYS	CA-C-N	5.59	132.92	122.74
2	C	126	LYS	C-N-CA	5.59	132.92	122.74

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2188	2107	2113	26	0
1	B	2107	2046	2050	23	0
2	C	663	642	642	12	0
2	D	716	738	740	8	0
3	A	30	40	40	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	24	29	32	1	0
4	A	15	0	0	1	0
4	B	15	0	0	0	0
5	A	106	0	0	6	1
5	B	114	0	0	6	1
5	C	16	0	0	1	0
5	D	32	0	0	3	0
All	All	6026	5602	5617	66	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:115:GLN:OE1	5:B:601:HOH:O	1.85	0.93
2:D:63:GLU:OE1	5:D:201:HOH:O	1.90	0.90
1:A:172:LYS:NZ	5:A:502:HOH:O	2.06	0.88
1:B:205:MET:HE1	1:B:214:ALA:HB2	1.64	0.80
1:A:352:LEU:HD11	2:C:55:PHE:CZ	2.17	0.79
1:B:348:GLU:OE1	5:B:602:HOH:O	2.00	0.79
2:D:111:GLY:O	5:D:201:HOH:O	2.01	0.77
4:A:407:SO4:O3	5:A:501:HOH:O	2.03	0.74
1:B:124:ARG:NH1	5:B:604:HOH:O	2.22	0.72
1:A:101[B]:GLN:NE2	5:A:506:HOH:O	2.22	0.72
3:B:504:GOL:O2	5:B:603:HOH:O	2.07	0.72
1:A:224:GLU:OE2	5:A:503:HOH:O	2.09	0.69
3:A:405:GOL:O1	5:D:201:HOH:O	2.13	0.66
1:A:100:THR:HG23	1:A:103:ALA:H	1.61	0.65
2:C:116:GLN:NE2	5:C:201:HOH:O	2.29	0.65
1:B:222:GLU:N	1:B:222:GLU:OE2	2.25	0.63
1:B:139:MET:HE1	1:B:228:LEU:HD13	1.80	0.62
1:A:308[A]:SER:OG	5:A:504:HOH:O	2.15	0.61
1:B:133:ALA:HA	1:B:342[A]:ILE:HD11	1.82	0.60
5:B:601:HOH:O	2:D:58:ARG:HD3	2.01	0.60
1:B:288:GLN:NE2	5:B:610:HOH:O	2.38	0.56
1:A:155:MET:HE2	1:A:193:THR:HA	1.89	0.55
1:A:157:LEU:HD12	1:A:161:LEU:HD13	1.91	0.52
2:C:55:PHE:HB2	2:C:83:ALA:O	2.10	0.52
1:B:139:MET:CE	1:B:228:LEU:HD13	2.40	0.52
1:A:352:LEU:N	1:A:352:LEU:HD23	2.25	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:LEU:HD11	2:C:55:PHE:CE2	2.46	0.51
1:A:205:MET:HE1	1:A:214:ALA:HB2	1.92	0.51
1:B:121:ARG:HB2	1:B:342[B]:ILE:HG13	1.92	0.51
2:C:79:GLN:HG3	2:C:117:VAL:HG21	1.94	0.50
1:A:94:LYS:O	1:A:97:ARG:NH1	2.45	0.49
1:A:87:ASP:OD1	2:C:68:ARG:NH2	2.44	0.49
1:B:120:ILE:HG12	1:B:341:ASN:HB3	1.95	0.49
1:B:169:LYS:HB3	1:B:169:LYS:NZ	2.28	0.49
1:A:205:MET:HE1	1:A:210:ALA:O	2.15	0.47
1:A:117:PHE:CG	1:A:343:PRO:HB2	2.49	0.47
1:B:120:ILE:HD11	1:B:341:ASN:HB2	1.97	0.46
1:A:352:LEU:HD23	2:C:55:PHE:O	2.16	0.46
1:B:348:GLU:HB2	2:D:61:VAL:HG22	1.97	0.46
1:B:120:ILE:HD11	1:B:341:ASN:CB	2.46	0.45
1:B:117:PHE:CG	1:B:343:PRO:HB2	2.52	0.45
1:B:266:SER:O	1:B:266:SER:OG	2.32	0.45
2:C:46:ILE:HD11	2:C:95:VAL:CG2	2.47	0.45
1:A:280:VAL:HA	1:A:284:GLY:O	2.17	0.45
2:D:85:LEU:HD12	2:D:86:PRO:HD2	1.99	0.44
1:A:267:ASN:HB2	1:A:271:GLN:OE1	2.18	0.44
2:C:117:VAL:O	2:C:121:ILE:HG12	2.18	0.44
1:B:352:LEU:HD21	2:D:57[A]:VAL:HG12	1.99	0.44
1:A:127:ASN:HB3	1:A:281:GLY:H	1.82	0.43
1:A:194:LEU:HD23	1:A:227:SER:HB3	2.00	0.43
1:A:202:LEU:CD2	1:A:214:ALA:HB3	2.48	0.43
1:A:352:LEU:HD23	1:A:352:LEU:H	1.82	0.43
1:A:336:ASP:OD2	1:A:339:HIS:CD2	2.72	0.43
1:B:303:GLN:O	1:B:331:THR:HA	2.18	0.43
2:C:133:LEU:C	2:C:133:LEU:HD23	2.44	0.42
1:A:157:LEU:CD1	1:A:161:LEU:HD13	2.50	0.42
1:B:139:MET:HE1	1:B:228:LEU:CD1	2.48	0.41
1:A:97:ARG:CZ	1:A:97:ARG:HB2	2.50	0.41
1:B:128:TYR:CZ	1:B:281:GLY:HA3	2.55	0.41
2:D:79:GLN:HG3	2:D:117:VAL:HG21	2.02	0.41
1:A:212:GLN:OE1	5:A:505:HOH:O	2.21	0.41
2:C:76:ALA:HB3	2:C:110:GLU:HG2	2.02	0.41
2:D:46:ILE:HD11	2:D:95:VAL:HG21	2.03	0.41
1:B:137:GLN:CD	1:B:342[A]:ILE:CG2	2.94	0.41
1:B:283:THR:O	1:B:283:THR:HG22	2.20	0.41
2:C:104:VAL:O	2:C:104:VAL:HG13	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:599:HOH:O	5:B:610:HOH:O[1_464]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	276/275 (100%)	270 (98%)	6 (2%)	0	100	100
1	B	263/275 (96%)	257 (98%)	6 (2%)	0	100	100
2	C	93/96 (97%)	89 (96%)	4 (4%)	0	100	100
2	D	96/96 (100%)	95 (99%)	1 (1%)	0	100	100
All	All	728/742 (98%)	711 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/243 (95%)	228 (99%)	2 (1%)	75	81
1	B	222/243 (91%)	222 (100%)	0	100	100
2	C	65/76 (86%)	65 (100%)	0	100	100
2	D	75/76 (99%)	75 (100%)	0	100	100
All	All	592/638 (93%)	590 (100%)	2 (0%)	92	94

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	215[A]	CYS
1	A	215[B]	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	127	ASN
1	A	275	ASN
1	A	303	GLN
1	A	339	HIS
1	B	288	GLN
2	C	105	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	403	-	5,5,5	0.70	0	5,5,5	1.06	0
3	GOL	A	405	-	5,5,5	0.55	0	5,5,5	0.72	0
4	SO4	B	506	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	A	406	-	4,4,4	0.14	0	6,6,6	0.06	0
3	GOL	B	503	-	5,5,5	0.42	0	5,5,5	0.67	0
3	GOL	A	402	-	5,5,5	0.76	0	5,5,5	0.97	0
3	GOL	B	504	-	5,5,5	0.72	0	5,5,5	1.05	0
4	SO4	B	507	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	A	408	-	4,4,4	0.14	0	6,6,6	0.05	0
3	GOL	A	404	-	5,5,5	0.85	0	5,5,5	1.06	0
4	SO4	B	505	-	4,4,4	0.14	0	6,6,6	0.06	0
3	GOL	B	502	-	5,5,5	0.70	0	5,5,5	0.91	0
3	GOL	B	501	-	5,5,5	0.75	0	5,5,5	1.11	1 (20%)
4	SO4	A	407	-	4,4,4	0.14	0	6,6,6	0.04	0
3	GOL	A	401	-	5,5,5	0.52	0	5,5,5	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	403	-	-	4/4/4/4	-
3	GOL	A	405	-	-	2/4/4/4	-
3	GOL	B	503	-	-	4/4/4/4	-
3	GOL	A	402	-	-	0/4/4/4	-
3	GOL	B	504	-	-	4/4/4/4	-
3	GOL	A	404	-	-	2/4/4/4	-
3	GOL	B	502	-	-	4/4/4/4	-
3	GOL	B	501	-	-	2/4/4/4	-
3	GOL	A	401	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	501	GOL	C3-C2-C1	-2.00	103.92	111.70

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	GOL	C1-C2-C3-O3
3	A	405	GOL	O1-C1-C2-O2
3	A	405	GOL	O1-C1-C2-C3
3	B	501	GOL	C1-C2-C3-O3
3	B	501	GOL	O2-C2-C3-O3
3	B	502	GOL	C1-C2-C3-O3
3	B	503	GOL	O1-C1-C2-O2
3	B	503	GOL	O1-C1-C2-C3
3	B	503	GOL	C1-C2-C3-O3
3	B	504	GOL	O1-C1-C2-C3
3	B	504	GOL	O1-C1-C2-O2
3	A	403	GOL	O1-C1-C2-C3
3	B	502	GOL	O1-C1-C2-C3
3	B	504	GOL	C1-C2-C3-O3
3	A	403	GOL	O2-C2-C3-O3
3	B	503	GOL	O2-C2-C3-O3
3	B	502	GOL	O2-C2-C3-O3
3	A	404	GOL	O2-C2-C3-O3
3	B	504	GOL	O2-C2-C3-O3
3	B	502	GOL	O1-C1-C2-O2
3	A	403	GOL	O1-C1-C2-O2
3	A	404	GOL	C1-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	405	GOL	1	0
3	B	504	GOL	1	0
4	A	407	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/275 (99%)	0.75	23 (8%) 18 17	22, 54, 82, 116	4 (1%)
1	B	264/275 (96%)	0.60	20 (7%) 21 20	19, 50, 75, 99	3 (1%)
2	C	95/96 (98%)	1.67	35 (36%) 1 1	39, 81, 129, 159	0
2	D	96/96 (100%)	0.81	7 (7%) 22 21	25, 60, 100, 132	2 (2%)
All	All	729/742 (98%)	0.82	85 (11%) 10 9	19, 55, 98, 159	9 (1%)

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	55	PHE	4.8
2	D	126	LYS	4.5
1	A	181	ASN	4.2
1	B	89	MET	3.9
2	C	123	ALA	3.9
1	A	338	ARG	3.8
2	C	92	ARG	3.8
2	C	121	ILE	3.8
2	C	94	GLY	3.8
2	C	133	LEU	3.6
2	C	51	SER	3.5
2	C	50	GLU	3.5
2	C	43	VAL	3.4
2	D	50	GLU	3.4
1	B	216	ASP	3.3
1	A	352	LEU	3.3
2	C	93	ALA	3.3
1	A	80	LEU	3.1
1	A	93	LYS	3.1
1	A	311	ASN	3.0
1	A	180	LYS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	79	PRO	3.0
1	B	311	ASN	2.9
1	B	125	GLY	2.9
2	C	120	LEU	2.9
2	D	135	VAL	2.9
1	B	80	LEU	2.8
1	B	350	THR	2.8
1	B	338	ARG	2.8
2	D	125	GLU	2.8
1	A	166	ASN	2.7
1	B	106	MET	2.7
1	A	249	LYS	2.7
2	C	42	ARG	2.7
1	B	173	LEU	2.7
1	A	255	PHE	2.6
1	A	89	MET	2.6
1	A	155	MET	2.6
2	C	130	LEU	2.6
1	A	351	SER	2.6
1	B	120	ILE	2.6
2	C	53	TYR	2.5
1	B	204	GLU	2.5
1	A	236	MET	2.5
1	B	205	MET	2.5
2	C	40	GLY	2.4
1	A	101[A]	GLN	2.4
1	A	175	LEU	2.4
2	C	128	LEU	2.4
2	C	108	ASN	2.4
2	D	40	GLY	2.4
1	B	264	ASP	2.3
2	C	48	LYS	2.3
2	C	71	ASN	2.3
2	C	102	LEU	2.3
2	C	87	GLY	2.3
2	C	126	LYS	2.3
1	A	276	HIS	2.3
1	A	161	LEU	2.3
1	B	122	ARG	2.3
2	C	89	ALA	2.3
2	C	104	VAL	2.3
1	A	152	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	188	ILE	2.2
2	C	46	ILE	2.2
2	C	90	ALA	2.2
2	C	125	GLU	2.2
2	C	86	PRO	2.2
2	D	41	PRO	2.2
2	C	88	GLY	2.2
2	D	45	ARG	2.2
2	C	66	GLN	2.2
1	A	182	GLU	2.1
1	A	100	THR	2.1
1	A	247	LYS	2.1
1	B	183	ASP	2.1
1	B	97	ARG	2.1
1	B	282	HIS	2.1
2	C	129	ILE	2.1
1	B	189	LYS	2.1
1	B	152	PRO	2.1
2	C	65	GLY	2.1
2	C	73	GLU	2.1
2	C	106	HIS	2.0
2	C	116	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	404	6/6	0.67	0.21	77,93,94,94	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	SO4	B	507	5/5	0.68	0.15	111,111,111,111	0
3	GOL	B	501	6/6	0.73	0.20	75,91,92,92	0
3	GOL	A	401	6/6	0.75	0.20	74,89,91,93	0
3	GOL	B	503	6/6	0.76	0.22	75,90,91,91	0
3	GOL	A	402	6/6	0.76	0.16	64,77,81,83	0
4	SO4	A	407	5/5	0.78	0.10	128,128,128,128	0
3	GOL	B	504	6/6	0.78	0.19	91,92,110,110	0
3	GOL	A	403	6/6	0.79	0.18	67,81,83,84	0
4	SO4	B	505	5/5	0.81	0.24	121,121,122,122	0
4	SO4	A	406	5/5	0.82	0.17	98,98,98,98	0
3	GOL	B	502	6/6	0.82	0.17	67,71,85,85	0
4	SO4	A	408	5/5	0.83	0.13	105,106,106,106	0
4	SO4	B	506	5/5	0.86	0.16	103,103,103,103	0
3	GOL	A	405	6/6	0.92	0.12	68,81,82,96	0

6.5 Other polymers [i](#)

There are no such residues in this entry.