



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2025 – 02:25 am BST

PDB ID : 6T5U / pdb\_00006t5u  
Title : KRasG12C ligand complex  
Authors : Phillips, C.  
Deposited on : 2019-10-17  
Resolution : 1.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.46


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7106 (1.74-1.70)
Clashscore	180529	7746 (1.74-1.70)
Ramachandran outliers	177936	7654 (1.74-1.70)
Sidechain outliers	177891	7654 (1.74-1.70)
RSRZ outliers	164620	7104 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	168	<div> <div>11%</div> <div>74%</div> <div>24%</div> <div>.</div> </div>
1	B	168	<div> <div>23%</div> <div>89%</div> <div>10%</div> <div>..</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2984 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called V-Ki-ras2 Kirsten rat sarcoma viral oncogene homolog, isoform CRA\_b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	168	Total	C	N	O	S	0	1	0
			1349	848	230	266	5			
1	B	167	Total	C	N	O	S	0	1	0
			1343	843	229	266	5			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP A0A024RAV5
A	12	CYS	GLY	conflict	UNP A0A024RAV5
A	51	SER	CYS	conflict	UNP A0A024RAV5
A	80	LEU	CYS	conflict	UNP A0A024RAV5
A	118	SER	CYS	conflict	UNP A0A024RAV5
A	167	LYS	-	expression tag	UNP A0A024RAV5
B	0	GLY	-	expression tag	UNP A0A024RAV5
B	12	CYS	GLY	conflict	UNP A0A024RAV5
B	51	SER	CYS	conflict	UNP A0A024RAV5
B	80	LEU	CYS	conflict	UNP A0A024RAV5
B	118	SER	CYS	conflict	UNP A0A024RAV5
B	167	LYS	-	expression tag	UNP A0A024RAV5

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>11</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
3	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 4 is 1-[(7R)-16-chloro-15-(5-methyl-1H-indazol-4-yl)-9-oxa-2,5,12-triazatetracyclo[8.8.0.02,7.013,18]octadeca-1(10),11,13,15,17-pentaen-5-yl]prop-2-en-1-one (CCD ID: MKW) (formula: C<sub>25</sub>H<sub>24</sub>ClN<sub>5</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	0	0
			33	25	1	5	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	Cl	N	O	0	0
			33	25	1	5	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	106	Total	O	0	0
			106	106		
5	B	62	Total	O	0	0
			62	62		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: V-Ki-ras2 Kirsten rat sarcoma viral oncogene homolog, isoform CRA\_b



- Molecule 1: V-Ki-ras2 Kirsten rat sarcoma viral oncogene homolog, isoform CRA\_b



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	33.60Å 41.08Å 65.19Å 78.47° 85.59° 68.14°	Depositor
Resolution (Å)	31.94 – 1.72 31.94 – 1.72	Depositor EDS
% Data completeness (in resolution range)	90.0 (31.94-1.72) 89.8 (31.94-1.72)	Depositor EDS
$R_{merge}$	0.01	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	50.57 (at 1.72Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.237 , 0.291 0.244 , 0.298	Depositor DCC
$R_{free}$ test set	1460 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.8	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 44.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.59$ , $\langle L^2 \rangle = 0.45$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2984	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MKW, GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.48	17/1374 (1.2%)	1.36	7/1854 (0.4%)
1	B	1.07	1/1365 (0.1%)	1.11	4/1843 (0.2%)
All	All	1.29	18/2739 (0.7%)	1.24	11/3697 (0.3%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	78	PHE	N-CA	7.60	1.55	1.46
1	A	114	VAL	N-CA	-6.86	1.37	1.46
1	A	157	TYR	N-CA	6.56	1.54	1.46
1	A	77	GLY	N-CA	-6.56	1.39	1.45
1	A	164	ARG	N-CA	6.37	1.54	1.46
1	A	114	VAL	C-O	-6.12	1.17	1.24
1	A	139	ILE	C-O	-5.79	1.19	1.24
1	A	161	ARG	N-CA	5.50	1.53	1.46
1	B	99	GLN	C-O	-5.49	1.17	1.24
1	A	115	GLY	N-CA	5.41	1.53	1.45
1	A	85	ASN	N-CA	5.40	1.52	1.46
1	A	143	GLU	N-CA	5.29	1.53	1.46
1	A	78	PHE	CA-C	-5.24	1.46	1.52
1	A	116	ASN	N-CA	-5.22	1.39	1.45
1	A	32	TYR	C-O	-5.13	1.18	1.24
1	A	145	SER	C-O	5.12	1.30	1.24
1	A	4	TYR	C-O	-5.11	1.18	1.24
1	A	135	ARG	N-CA	5.10	1.52	1.46

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	114	VAL	CA-C-O	-6.14	114.06	120.27

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	116	ASN	CA-C-O	-5.72	114.97	121.72
1	B	109	VAL	CA-C-N	-5.69	113.93	119.90
1	B	109	VAL	C-N-CA	-5.69	113.93	119.90
1	A	116	ASN	O-C-N	5.62	129.24	122.94
1	A	29	VAL	CB-CA-C	5.54	118.28	111.25
1	A	77	GLY	O-C-N	5.45	127.41	123.27
1	A	152	VAL	N-CA-CB	5.39	117.47	110.57
1	B	74	THR	CA-C-N	5.28	124.99	119.92
1	B	74	THR	C-N-CA	5.28	124.99	119.92
1	A	13	GLY	N-CA-C	5.27	122.62	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1349	0	1333	18	0
1	B	1343	0	1319	6	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	28	0	12	0	0
3	B	28	0	12	0	0
4	A	33	0	0	2	0
4	B	33	0	0	0	0
5	A	106	0	0	2	0
5	B	62	0	0	0	0
All	All	2984	0	2676	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:24:ILE:HD12	1:A:42:LYS:HB2	1.62	0.81
1:A:62:GLU:HG3	4:A:203:MKW:C12	2.12	0.78
1:B:46:ILE:HD13	1:B:160:VAL:HG11	1.66	0.78
1:A:63:GLU:OE2	5:A:301:HOH:O	2.02	0.76
1:A:22:GLN:HG3	1:A:146:ALA:O	2.01	0.59
1:A:112:VAL:HG22	1:A:140:PRO:HG2	1.83	0.59
1:A:24:ILE:HD12	1:A:42:LYS:CB	2.34	0.57
1:A:67:MET:HE2	1:A:71:TYR:CE1	2.39	0.57
1:A:72:MET:HB2	1:A:103:VAL:HG21	1.88	0.55
1:A:62:GLU:HG3	4:A:203:MKW:C13	2.35	0.55
1:A:32:TYR:O	1:A:33:ASP:C	2.53	0.52
1:B:64[A]:TYR:CD1	1:B:64[A]:TYR:N	2.77	0.51
1:B:39:SER:HA	1:B:55:ILE:O	2.10	0.51
1:B:97:ARG:O	1:B:101:LYS:HG3	2.10	0.50
1:B:33:ASP:CG	1:B:35:THR:HG1	2.25	0.45
1:A:106:SER:OG	1:A:107:GLU:N	2.48	0.45
1:B:64[B]:TYR:CD1	1:B:64[B]:TYR:C	2.91	0.45
1:A:26:ASN:HD22	1:A:149:ARG:HH22	1.65	0.43
1:A:79:LEU:HD23	1:A:159:LEU:HD22	1.99	0.43
1:A:41:ARG:HH21	1:A:52:LEU:HD21	1.84	0.42
1:A:166:HIS:HB3	5:A:374:HOH:O	2.19	0.41
1:A:33:ASP:CG	1:A:36:ILE:HG13	2.46	0.41
1:A:26:ASN:HD22	1:A:149:ARG:NH2	2.18	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	167/168 (99%)	164 (98%)	1 (1%)	2 (1%)	11	2
1	B	166/168 (99%)	157 (95%)	8 (5%)	1 (1%)	22	9
All	All	333/336 (99%)	321 (96%)	9 (3%)	3 (1%)	14	4

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	59	ALA
1	B	31	GLU
1	A	33	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	149/148 (101%)	146 (98%)	3 (2%)	50	33
1	B	148/148 (100%)	143 (97%)	5 (3%)	32	14
All	All	297/296 (100%)	289 (97%)	8 (3%)	40	21

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	84	ILE
1	A	107	GLU
1	A	149	ARG
1	B	31	GLU
1	B	49	GLU
1	B	62	GLU
1	B	106	SER
1	B	129	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	GLN
1	A	26	ASN
1	A	27	HIS
1	A	43	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	GDP	B	202	2	24,30,30	1.10	1 (4%)	30,47,47	1.50	7 (23%)
4	MKW	B	203	-	35,38,38	0.81	1 (2%)	44,57,57	1.25	4 (9%)
4	MKW	A	203	-	35,38,38	0.92	3 (8%)	44,57,57	1.22	6 (13%)
3	GDP	A	202	2	24,30,30	1.28	4 (16%)	30,47,47	1.86	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	B	202	2	-	1/12/32/32	0/3/3/3
4	MKW	B	203	-	-	2/10/32/32	0/5/6/6
4	MKW	A	203	-	-	0/10/32/32	0/5/6/6
3	GDP	A	202	2	-	3/12/32/32	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	203	MKW	C1-C2	2.54	1.55	1.51
3	A	202	GDP	O3'-C3'	2.49	1.48	1.43
4	A	203	MKW	C20-C21	-2.41	1.37	1.41
3	B	202	GDP	O4'-C4'	-2.29	1.39	1.45
4	A	203	MKW	C12-C13	2.22	1.42	1.38
3	A	202	GDP	PA-O2A	-2.17	1.45	1.55
4	A	203	MKW	C1-C2	2.13	1.54	1.51
3	A	202	GDP	C2'-C1'	-2.12	1.50	1.53
3	A	202	GDP	O4'-C1'	-2.03	1.38	1.41

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	202	GDP	C2-N1-C6	-4.22	117.33	125.10
4	B	203	MKW	C15-C16-N	4.07	115.05	109.57
3	B	202	GDP	O2A-PA-O5'	3.77	125.28	107.75
3	A	202	GDP	C5-C6-N1	3.77	120.61	113.95
3	A	202	GDP	O6-C6-N1	-3.73	116.24	120.65
3	A	202	GDP	O3B-PB-O2B	3.45	120.83	107.64
4	A	203	MKW	O1-C13-C12	2.97	121.63	117.06
4	B	203	MKW	C4-N1-C5	-2.79	113.29	118.71
3	B	202	GDP	O6-C6-N1	-2.76	117.39	120.65
3	A	202	GDP	O2'-C2'-C3'	2.67	120.46	111.82
3	A	202	GDP	PA-O3A-PB	-2.64	123.75	132.83
3	B	202	GDP	C8-N7-C5	2.64	108.02	102.99
4	A	203	MKW	C16-N-C2	2.57	130.31	122.03
4	B	203	MKW	O1-C13-C12	2.56	120.98	117.06
3	B	202	GDP	O4'-C1'-C2'	-2.54	103.21	106.93
3	B	202	GDP	C3'-C2'-C1'	2.50	104.75	100.98
4	A	203	MKW	C3-N-C2	-2.33	115.22	122.81
4	A	203	MKW	C4-N1-C5	-2.31	114.22	118.71
4	A	203	MKW	C15-C16-N	2.14	112.45	109.57
3	B	202	GDP	O2B-PB-O1B	-2.13	102.35	110.68
3	B	202	GDP	C5-C6-N1	2.12	117.69	113.95
4	B	203	MKW	C16-N-C2	2.08	128.72	122.03
4	A	203	MKW	C1-C2-N	2.03	121.23	117.95

There are no chirality outliers.

All (6) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	202	GDP	PA-O3A-PB-O2B
3	A	202	GDP	PA-O3A-PB-O3B
4	B	203	MKW	C-C1-C2-N
3	B	202	GDP	PA-O3A-PB-O3B
4	B	203	MKW	C-C1-C2-O
3	A	202	GDP	PA-O3A-PB-O1B


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	203	MKW	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers ⓘ

There are no such residues in this entry.



## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	168/168 (100%)	0.82	18 (10%)	12 13	12, 26, 48, 85	1 (0%)
1	B	167/168 (99%)	1.51	39 (23%)	2 2	17, 37, 59, 91	1 (0%)
All	All	335/336 (99%)	1.17	57 (17%)	5 5	12, 31, 57, 91	2 (0%)

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	64[A]	TYR	5.2
1	B	66	ALA	4.6
1	B	157	TYR	4.2
1	B	36	ILE	4.2
1	B	0	GLY	4.2
1	B	107	GLU	4.0
1	B	120	LEU	4.0
1	B	40	TYR	3.9
1	A	1	MET	3.9
1	B	153	ASP	3.9
1	A	64[A]	TYR	3.8
1	B	121	PRO	3.6
1	A	32	TYR	3.4
1	B	124	THR	3.3
1	B	71	TYR	3.3
1	B	57	ASP	3.3
1	A	48	GLY	3.2
1	A	103	VAL	3.0
1	A	166	HIS	3.0
1	B	106	SER	2.9
1	B	105	ASP	2.8
1	A	157	TYR	2.8
1	A	0	GLY	2.8
1	B	125	VAL	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	34	PRO	2.6
1	A	29	VAL	2.6
1	B	33	ASP	2.6
1	B	166	HIS	2.6
1	B	12	CYS	2.6
1	A	30	ASP	2.6
1	A	49	GLU	2.5
1	B	39	SER	2.4
1	A	107	GLU	2.4
1	B	45	VAL	2.4
1	B	103	VAL	2.4
1	B	152	VAL	2.3
1	B	74	THR	2.3
1	B	6	LEU	2.3
1	A	71	TYR	2.3
1	B	32	TYR	2.3
1	B	44	VAL	2.3
1	B	136	SER	2.3
1	B	38	ASP	2.2
1	B	126	ASP	2.2
1	B	146	ALA	2.2
1	B	127	THR	2.2
1	B	53	LEU	2.2
1	A	24	ILE	2.1
1	B	52	LEU	2.1
1	B	26	ASN	2.1
1	B	142	ILE	2.1
1	A	31	GLU	2.1
1	B	84	ILE	2.0
1	A	40	TYR	2.0
1	B	91	GLU	2.0
1	A	167	LYS	2.0
1	A	105	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MKW	B	203	33/33	0.78	0.14	32,37,41,43	0
4	MKW	A	203	33/33	0.85	0.11	22,24,28,31	0
2	MG	B	201	1/1	0.87	0.08	37,37,37,37	0
3	GDP	B	202	28/28	0.90	0.10	29,34,39,41	0
2	MG	A	201	1/1	0.96	0.04	21,21,21,21	0
3	GDP	A	202	28/28	0.97	0.06	18,24,27,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around MKW A 203:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around GDP B 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around GDP A 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.