



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 4, 2023 – 06:39 PM EDT

PDB ID : 6UJE
Title : Crystal structure of the Clostridial cellulose synthase subunit Z (CcsZ) from Clostridioides difficile
Authors : Scott, W.; Lowrance, B.; Anderson, A.C.; Weadge, J.T.
Deposited on : 2019-10-03
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2763 atoms, of which 86 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endoglucanase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	301	2642	1656	86	422	474	4	0	1	0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	120	Total	O	0	0
			120	120		

SEQUENCE-PLOTS INFOmissingINFO

3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.95Å 45.63Å 59.94Å 90.00° 94.99° 90.00°	Depositor
Resolution (Å)	43.04 – 1.75	Depositor
% Data completeness (in resolution range)	96.9 (43.04-1.75)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 1.75Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.227 , 0.260	Depositor
Wilson B-factor (Å ²)	36.4	Xtrriage
Anisotropy	0.339	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.016 for l,-k,h	Xtrriage
Total number of atoms	2763	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality (i)

4.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	5/2626 (0.2%)	0.57	6/3544 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	226	GLU	C-N	7.74	1.49	1.34
1	A	127	GLU	CD-OE2	-6.95	1.18	1.25
1	A	226	GLU	CD-OE2	-6.37	1.18	1.25
1	A	226	GLU	CD-OE1	-5.85	1.19	1.25
1	A	315	GLU	CD-OE1	-5.54	1.19	1.25

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	315	GLU	CB-CA-C	11.03	132.46	110.40
1	A	315	GLU	CB-CG-CD	-9.01	89.88	114.20
1	A	314	TRP	O-C-N	6.78	133.54	122.70
1	A	315	GLU	N-CA-CB	-6.13	99.56	110.60
1	A	314	TRP	N-CA-CB	6.08	121.53	110.60
1	A	314	TRP	CA-C-N	-5.19	105.78	117.20

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2556	86	2483	40	1
2	A	1	0	0	0	0
3	A	120	0	0	2	0
All	All	2677	86	2483	40	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215:ASP:HB3	1:A:218:ILE:HD12	1.53	0.91
1:A:239:LYS:NZ	1:A:239:LYS:HB3	1.94	0.81
1:A:236:ILE:HD12	1:A:237:TYR:N	2.05	0.70
1:A:239:LYS:HB3	1:A:239:LYS:HZ2	1.57	0.70
1:A:212:ILE:HD12	1:A:212:ILE:H	1.59	0.68
1:A:101:PHE:CZ	1:A:149:ARG:HD3	2.37	0.60
1:A:213:PRO:HG2	1:A:218:ILE:HD13	1.85	0.59
1:A:268:ASN:O	1:A:272:LYS:HG3	2.03	0.59
1:A:215:ASP:HB3	1:A:218:ILE:CD1	2.30	0.58
1:A:200:ASN:O	1:A:203:GLN:HB2	2.04	0.57
1:A:101:PHE:CE1	1:A:149:ARG:HD3	2.41	0.56
1:A:43:GLY:HA2	1:A:77:THR:O	2.09	0.52
1:A:197:GLY:HA3	1:A:202:TYR:CE1	2.44	0.52
1:A:164:GLU:HB3	1:A:201:PHE:HA	1.92	0.51
1:A:195:ILE:HG12	1:A:219:VAL:HB	1.93	0.51
1:A:223:HIS:CG	1:A:281:GLU:HB2	2.46	0.51
1:A:236:ILE:HD11	1:A:237:TYR:CE2	2.46	0.51
1:A:151:GLN:HG3	1:A:152:LYS:HD3	1.93	0.51
1:A:314:TRP:CD2	1:A:315:GLU:HB2	2.47	0.50
1:A:143:TRP:CZ2	1:A:161:LEU:HA	2.47	0.49
1:A:92:ASP:OD1	1:A:92:ASP:N	2.45	0.49
1:A:212:ILE:HD12	1:A:212:ILE:N	2.28	0.49
1:A:236:ILE:HD12	1:A:237:TYR:H	1.76	0.49
1:A:261:LYS:HD2	1:A:261:LYS:O	2.13	0.49
1:A:195:ILE:HA	1:A:219:VAL:O	2.14	0.48
1:A:94:PHE:O	1:A:145:GLN:NE2	2.40	0.46
1:A:54:LYS:HE2	1:A:86:ASP:O	2.15	0.46
1:A:81:PRO:HB3	1:A:123:HIS:CG	2.52	0.45
1:A:89:SER:OG	1:A:94:PHE:N	2.50	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:THR:OG1	1:A:138:LYS:HE3	2.18	0.44
1:A:51:GLU:HB3	1:A:81:PRO:HB2	2.00	0.43
1:A:213:PRO:CG	1:A:218:ILE:HD13	2.48	0.43
1:A:89:SER:O	1:A:94:PHE:HA	2.18	0.43
1:A:284:VAL:HB	1:A:293:ARG:HG3	2.01	0.43
1:A:148:ASN:O	1:A:151:GLN:HG2	2.20	0.42
1:A:280:GLY:HA3	3:A:519:HOH:O	2.19	0.42
1:A:256:MET:O	1:A:260:LYS:HG3	2.20	0.42
1:A:141:LYS:O	1:A:145:GLN:HG3	2.20	0.41
1:A:315:GLU:CD	1:A:318:SER:HB3	2.41	0.41
1:A:319:GLY:HA3	3:A:560:HOH:O	2.19	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55:ASP:O	1:A:93:ASN:ND2[2_546]	2.18	0.02

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	298/320 (93%)	284 (95%)	13 (4%)	1 (0%)	41 22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	315	GLU

4.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	276/294 (94%)	264 (96%)	12 (4%)	29 9

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	ASP
1	A	56	PHE
1	A	86	ASP
1	A	90	ASP
1	A	92	ASP
1	A	101	PHE
1	A	125	PHE
1	A	127	GLU
1	A	191	LYS
1	A	205	ASP
1	A	216	SER
1	A	239	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers

There are no such residues in this entry.

4.8 Polymer linkage issues

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

5.4 Ligands

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers

EDS failed to run properly - this section is therefore empty.