



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2023 – 03:27 AM EDT

PDB ID : 6VUA
Title : X-ray structure of human CD38 catalytic domain with 2'-Cl-araNAD+
Authors : Dai, Z.; Zhang, X.N.; Nasertorabi, F.; Han, G.W.; Stevens, R.C.; Zhang, Y.
Deposited on : 2020-02-14
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 4973 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	255	2216	1405	373	419	19	0	26	0
1	B	251	2094	1332	356	388	18	0	16	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	SER	-	expression tag	UNP P28907
A	100	ASP	ASN	engineered mutation	UNP P28907
A	164	ALA	ASN	engineered mutation	UNP P28907
A	209	ASP	ASN	engineered mutation	UNP P28907
A	219	ASP	ASN	engineered mutation	UNP P28907
B	44	SER	-	expression tag	UNP P28907
B	100	ASP	ASN	engineered mutation	UNP P28907
B	164	ALA	ASN	engineered mutation	UNP P28907
B	209	ASP	ASN	engineered mutation	UNP P28907
B	219	ASP	ASN	engineered mutation	UNP P28907

- Molecule 2 is [(2 {R},3 {S},4 {R},5 {R})-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-oxidanyl-phosphoryl [(2 {R},3 {R},4 {R})-4-chloranyl-3-oxidanyl-oxolan-2-yl] methyl hydrogen phosphate (three-letter code: ROJ) (formula: C₁₅H₂₂ClN₅O₁₂P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 7 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	C O	0	0
			10	6 4		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	282	Total	O	0	4
			286	286		
8	B	220	Total	O	0	6
			226	226		

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	114.76Å 114.76Å 97.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.71 – 1.50	Depositor
% Data completeness (in resolution range)	100.0 (28.71-1.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.166 , 0.190	Depositor
Wilson B-factor (Å ²)	23.9	Xtrriage
Anisotropy	0.427	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	4973	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	MPD	B	404	-	7,7,7	0.18	0	9,10,10	0.48	0
4	PO4	A	403	-	4,4,4	0.94	0	6,6,6	0.58	0
2	ROJ	A	401	1	32,38,38	1.19	3 (9%)	33,58,58	1.47	3 (9%)
6	MPD	A	405	-	7,7,7	0.14	0	9,10,10	0.29	0
5	PEG	B	406	-	6,6,6	0.24	0	5,5,5	0.20	0
5	PEG	B	407	-	6,6,6	0.17	0	5,5,5	0.18	0
7	PGE	A	406	-	9,9,9	0.23	0	8,8,8	0.21	0
4	PO4	B	403	-	4,4,4	1.70	1 (25%)	6,6,6	0.79	0
2	ROJ	B	401	1	32,38,38	1.23	2 (6%)	33,58,58	1.17	3 (9%)
5	PEG	A	407	-	6,6,6	0.25	0	5,5,5	0.18	0
5	PEG	B	408	-	6,6,6	0.19	0	5,5,5	0.13	0
6	MPD	B	405	-	7,7,7	0.14	0	9,10,10	0.44	0
5	PEG	A	404	-	6,6,6	0.15	0	5,5,5	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MPD	B	404	-	-	2/5/5/5	-
2	ROJ	A	401	1	-	1/18/51/51	0/4/4/4
6	MPD	A	405	-	-	0/5/5/5	-
5	PEG	B	406	-	-	2/4/4/4	-
5	PEG	B	407	-	-	3/4/4/4	-
7	PGE	A	406	-	-	1/7/7/7	-
2	ROJ	B	401	1	-	1/18/51/51	0/4/4/4
5	PEG	A	407	-	-	2/4/4/4	-
5	PEG	B	408	-	-	1/4/4/4	-
6	MPD	B	405	-	-	2/5/5/5	-
5	PEG	A	404	-	-	2/4/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	ROJ	C12-N5	3.96	1.38	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	ROJ	C12-N5	3.42	1.37	1.32
2	A	401	ROJ	C1-C4	-2.48	1.46	1.53
4	B	403	PO4	P-O1	2.40	1.56	1.50
2	B	401	ROJ	O11-C14	-2.38	1.37	1.43
2	A	401	ROJ	C9-N2	-2.05	1.31	1.34

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	ROJ	O2-C4-C1	5.27	109.37	104.70
2	A	401	ROJ	C10-C11-N3	2.94	124.82	120.35
2	B	401	ROJ	C10-C11-N3	2.90	124.76	120.35
2	B	401	ROJ	C8-N1-C13	-2.14	122.89	126.64
2	B	401	ROJ	O12-C15-C7	2.02	116.90	111.05
2	A	401	ROJ	N5-C12-N4	-2.00	125.55	128.68

There are no chirality outliers.

All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	B	404	MPD	O2-C2-C3-C4
6	B	404	MPD	CM-C2-C3-C4
5	A	407	PEG	O2-C3-C4-O4
5	A	404	PEG	O1-C1-C2-O2
5	A	407	PEG	O1-C1-C2-O2
5	B	407	PEG	O2-C3-C4-O4
5	B	407	PEG	O1-C1-C2-O2
5	A	404	PEG	C1-C2-O2-C3
5	B	408	PEG	O2-C3-C4-O4
6	B	405	MPD	C2-C3-C4-C5
5	B	406	PEG	C4-C3-O2-C2
5	B	407	PEG	C4-C3-O2-C2
7	A	406	PGE	C4-C3-O2-C2
2	B	401	ROJ	O9-C6-C7-O10
2	A	401	ROJ	O9-C6-C7-O10
5	B	406	PEG	O1-C1-C2-O2
6	B	405	MPD	C2-C3-C4-O4


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.