



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 9, 2023 – 10:06 AM EDT

PDB ID : 6WP3
Title : Pyruvate Kinase M2 Mutant-K433Q
Authors : Nandi, S.; Razzaghi, M.; Srivastava, D.; Dey, M.
Deposited on : 2020-04-26
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4003 (1.86-1.82)
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)
RSRZ outliers	127900	3957 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	550	 4% 84% 9% 7%
1	B	550	 3% 71% 7% 22%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7444 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate kinase PKM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	510	3793	2389	666	713	25	0	0	0
1	B	428	3200	2008	572	598	22	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MET	-	initiating methionine	UNP P14618
A	-17	GLY	-	expression tag	UNP P14618
A	-16	SER	-	expression tag	UNP P14618
A	-15	SER	-	expression tag	UNP P14618
A	-14	HIS	-	expression tag	UNP P14618
A	-13	HIS	-	expression tag	UNP P14618
A	-12	HIS	-	expression tag	UNP P14618
A	-11	HIS	-	expression tag	UNP P14618
A	-10	HIS	-	expression tag	UNP P14618
A	-9	HIS	-	expression tag	UNP P14618
A	-8	SER	-	expression tag	UNP P14618
A	-7	SER	-	expression tag	UNP P14618
A	-6	GLY	-	expression tag	UNP P14618
A	-5	LEU	-	expression tag	UNP P14618
A	-4	VAL	-	expression tag	UNP P14618
A	-3	PRO	-	expression tag	UNP P14618
A	-2	ARG	-	expression tag	UNP P14618
A	-1	GLY	-	expression tag	UNP P14618
A	0	SER	-	expression tag	UNP P14618
A	433	GLN	LYS	engineered mutation	UNP P14618
B	-18	MET	-	initiating methionine	UNP P14618
B	-17	GLY	-	expression tag	UNP P14618
B	-16	SER	-	expression tag	UNP P14618
B	-15	SER	-	expression tag	UNP P14618
B	-14	HIS	-	expression tag	UNP P14618

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-13	HIS	-	expression tag	UNP P14618
B	-12	HIS	-	expression tag	UNP P14618
B	-11	HIS	-	expression tag	UNP P14618
B	-10	HIS	-	expression tag	UNP P14618
B	-9	HIS	-	expression tag	UNP P14618
B	-8	SER	-	expression tag	UNP P14618
B	-7	SER	-	expression tag	UNP P14618
B	-6	GLY	-	expression tag	UNP P14618
B	-5	LEU	-	expression tag	UNP P14618
B	-4	VAL	-	expression tag	UNP P14618
B	-3	PRO	-	expression tag	UNP P14618
B	-2	ARG	-	expression tag	UNP P14618
B	-1	GLY	-	expression tag	UNP P14618
B	0	SER	-	expression tag	UNP P14618
B	433	GLN	LYS	engineered mutation	UNP P14618

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is OXALATE ION (three-letter code: OXL) (formula: C_2O_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	2	4		
3	B	1	Total	C	O	0	0
			6	2	4		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is NITRATE ION (three-letter code: NO3) (formula: NO_3).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total N O 4 1 3	0	0
5	B	1	Total N O 4 1 3	0	0

- Molecule 6 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total K 1 1	0	0

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total Cl 1 1	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	213	Total O 213 213	0	0
8	B	172	Total O 172 172	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	112.38Å 94.23Å 109.64Å 90.00° 94.80° 90.00°	Depositor
Resolution (Å)	47.11 – 1.84 47.11 – 1.84	Depositor EDS
% Data completeness (in resolution range)	97.6 (47.11-1.84) 97.6 (47.11-1.84)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 1.84Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.203 , 0.230 0.203 , 0.229	Depositor DCC
R_{free} test set	4815 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	26.1	Xtrriage
Anisotropy	0.499	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 52.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7444	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, OXL, CL, NO3, MG, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/3850	0.57	0/5213
1	B	0.39	0/3250	0.61	3/4398 (0.1%)
All	All	0.40	0/7100	0.59	3/9611 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	487	ASP	CB-CG-OD2	-12.52	107.04	118.30
1	B	487	ASP	CB-CG-OD1	5.57	123.31	118.30
1	B	467	ARG	NE-CZ-NH1	-5.52	117.54	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3793	0	3747	32	1
1	B	3200	0	3186	22	1
2	A	24	0	31	4	0
2	B	18	0	24	2	0
3	A	6	0	0	0	0
3	B	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	4	0	0	0	0
5	B	4	0	0	0	0
6	B	1	0	0	0	0
7	B	1	0	0	0	0
8	A	213	0	0	4	0
8	B	172	0	0	1	0
All	All	7444	0	6988	53	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:410:GLU:HG3	1:B:440:GLN:HE21	1.30	0.94
1:A:83:TYR:HE1	2:A:604:GOL:H2	1.50	0.75
1:B:407:ASP:OD2	1:B:436:ARG:NH2	2.21	0.74
1:A:494:MET:CE	1:A:531:PRO:HD2	2.18	0.73
1:A:494:MET:HE1	1:A:531:PRO:HD2	1.71	0.73
1:B:410:GLU:HG3	1:B:440:GLN:NE2	2.04	0.73
1:A:83:TYR:CE1	2:A:604:GOL:H2	2.25	0.71
1:B:16:GLN:HB2	1:B:18:LEU:HD13	1.75	0.67
1:A:87:THR:HG21	2:A:604:GOL:H11	1.81	0.62
1:B:436:ARG:HB3	2:B:603:GOL:H11	1.80	0.62
1:A:494:MET:HE1	1:A:530:VAL:HA	1.82	0.62
1:A:479:GLN:C	1:A:481:ALA:HB3	2.21	0.61
1:B:244:PHE:HD1	1:B:273:ASN:HD22	1.50	0.60
1:A:39:PRO:O	8:A:701:HOH:O	2.16	0.59
1:B:472:VAL:HG11	1:B:496:VAL:HG11	1.85	0.59
1:A:339:ARG:NH2	8:A:707:HOH:O	2.34	0.59
1:A:127:SER:O	8:A:702:HOH:O	2.17	0.57
1:A:395:PHE:CZ	1:A:399:ARG:HD2	2.39	0.57
1:B:272:GLU:HB3	1:B:293:ALA:HB3	1.86	0.56
1:A:395:PHE:CE2	1:A:399:ARG:HD2	2.40	0.56
1:A:132:VAL:HG21	1:A:153:ASP:HA	1.89	0.55
1:A:272:GLU:HB3	1:A:293:ALA:HB3	1.90	0.54
1:A:472:VAL:HG11	1:A:496:VAL:HG11	1.89	0.53
1:A:16:GLN:NE2	1:A:447:ARG:HD3	2.24	0.52
1:A:331:LEU:O	1:A:334:MET:HB2	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149:MET:HG3	1:A:150:GLU:HG3	1.92	0.52
1:B:39:PRO:O	8:B:701:HOH:O	2.18	0.51
1:A:40:ILE:O	8:A:703:HOH:O	2.20	0.49
1:A:497:GLY:HA3	1:A:503:PHE:CZ	2.48	0.49
1:A:175:TYR:HB3	1:A:179:GLY:HA2	1.95	0.48
1:A:416:ALA:HB2	1:A:512:LEU:HD21	1.94	0.48
1:A:27:LEU:HD23	1:B:401:LEU:HD12	1.96	0.47
1:B:505:LYS:HE2	1:B:531:PRO:OXT	2.14	0.46
1:A:505:LYS:HA	1:A:530:VAL:HG12	1.97	0.46
1:B:224:LYS:NZ	1:B:228:ASP:OD2	2.48	0.45
1:A:437:SER:HB3	1:A:522:THR:HG21	1.99	0.45
1:A:514:GLY:HA2	1:A:522:THR:HA	1.98	0.44
1:B:216:VAL:HB	2:B:602:GOL:H31	2.00	0.44
1:A:494:MET:HE2	1:A:531:PRO:HD2	1.96	0.44
1:B:393:GLN:O	1:B:397:GLU:HG3	2.17	0.43
1:B:40:ILE:O	1:B:383:ARG:NH2	2.49	0.43
1:A:244:PHE:CZ	2:A:603:GOL:H11	2.54	0.43
1:A:163:ASN:ND2	1:A:217:ASP:OD2	2.53	0.42
1:A:164:ILE:O	1:A:168:VAL:HG22	2.20	0.42
1:B:55:SER:HA	1:B:60:THR:HG21	2.03	0.41
1:B:18:LEU:O	1:B:22:MET:HG2	2.20	0.41
1:B:288:ASP:O	1:B:323:PRO:HD2	2.21	0.40
1:B:15:THR:HG22	1:B:38:PRO:HG2	2.02	0.40
1:A:494:MET:HE3	1:A:494:MET:HB3	1.92	0.40
1:B:331:LEU:HD23	1:B:344:GLU:HB3	2.03	0.40
1:B:453:VAL:HG21	1:B:493:ALA:HB2	2.03	0.40
1:A:121:THR:HB	1:A:157:LEU:HD11	2.03	0.40
1:B:103:ILE:HD13	1:B:492:PHE:CE1	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:434:SER:OG	1:B:223:GLU:OE1[3_455]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	500/550 (91%)	487 (97%)	10 (2%)	3 (1%)	25	12
1	B	420/550 (76%)	414 (99%)	5 (1%)	1 (0%)	47	33
All	All	920/1100 (84%)	901 (98%)	15 (2%)	4 (0%)	34	20

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	41	THR
1	A	517	PRO
1	A	328	THR
1	B	328	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	387/452 (86%)	387 (100%)	0	100	100
1	B	331/452 (73%)	331 (100%)	0	100	100
All	All	718/904 (79%)	718 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	GLN
1	A	227	GLN
1	B	433	GLN
1	B	440	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 4 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	601	-	5,5,5	0.72	0	5,5,5	1.25	1 (20%)
2	GOL	B	603	-	5,5,5	1.02	0	5,5,5	1.06	0
3	OXL	B	604	4	5,5,5	1.22	0	6,6,6	1.51	2 (33%)
5	NO3	B	606	-	1,3,3	0.63	0	0,3,3	-	-
2	GOL	A	601	-	5,5,5	0.87	0	5,5,5	0.88	0
2	GOL	A	603	-	5,5,5	0.63	0	5,5,5	1.25	0
2	GOL	A	602	-	5,5,5	0.78	0	5,5,5	1.12	1 (20%)
2	GOL	B	602	-	5,5,5	0.73	0	5,5,5	1.00	0
2	GOL	A	604	-	5,5,5	1.44	1 (20%)	5,5,5	1.31	0
3	OXL	A	605	4	5,5,5	1.28	0	6,6,6	1.19	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NO3	A	607	-	1,3,3	0.44	0	0,3,3	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	601	-	-	0/4/4/4	-
2	GOL	B	603	-	-	0/4/4/4	-
3	OXL	B	604	4	-	0/4/4/4	-
2	GOL	A	601	-	-	0/4/4/4	-
2	GOL	A	603	-	-	2/4/4/4	-
2	GOL	A	602	-	-	0/4/4/4	-
2	GOL	B	602	-	-	3/4/4/4	-
2	GOL	A	604	-	-	4/4/4/4	-
3	OXL	A	605	4	-	3/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	604	GOL	O2-C2	-2.06	1.37	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	604	OXL	O4-C2-C1	2.21	119.74	113.16
2	A	602	GOL	C3-C2-C1	-2.20	103.15	111.70
3	B	604	OXL	O3-C1-C2	2.17	119.59	113.16
2	B	601	GOL	C3-C2-C1	-2.08	103.62	111.70

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	603	GOL	C1-C2-C3-O3
2	A	604	GOL	O1-C1-C2-O2
2	A	604	GOL	O1-C1-C2-C3
2	A	604	GOL	C1-C2-C3-O3
2	B	602	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	602	GOL	C1-C2-C3-O3
2	A	603	GOL	O2-C2-C3-O3
2	A	604	GOL	O2-C2-C3-O3
2	B	602	GOL	O2-C2-C3-O3
3	A	605	OXL	O1-C1-C2-O2
3	A	605	OXL	O3-C1-C2-O4
3	A	605	OXL	O3-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	603	GOL	1	0
2	A	603	GOL	1	0
2	B	602	GOL	1	0
2	A	604	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	510/550 (92%)	0.07	20 (3%) 39 36	17, 28, 49, 61	0
1	B	428/550 (77%)	0.01	17 (3%) 38 35	17, 29, 46, 56	0
All	All	938/1100 (85%)	0.04	37 (3%) 39 36	17, 29, 48, 61	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	517	PRO	4.9
1	B	215	ALA	4.5
1	A	156	ILE	3.9
1	A	190	ALA	3.9
1	A	191	ASP	3.6
1	A	139	THR	3.6
1	B	480	GLU	3.2
1	A	138	ALA	3.2
1	B	478	VAL	3.2
1	A	170	VAL	3.1
1	A	194	VAL	3.0
1	B	481	ALA	3.0
1	A	187	GLN	2.9
1	A	131	GLU	2.9
1	A	401	LEU	2.8
1	B	477	PRO	2.8
1	A	481	ALA	2.8
1	B	520	GLY	2.6
1	A	193	LEU	2.6
1	B	214	ALA	2.6
1	B	479	GLN	2.5
1	B	58	VAL	2.5
1	B	491	ASN	2.4
1	B	13	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	184	GLN	2.3
1	A	188	LYS	2.3
1	B	211	LEU	2.3
1	A	134	LEU	2.3
1	B	103	ILE	2.2
1	B	482	TRP	2.2
1	B	100	SER	2.2
1	A	482	TRP	2.1
1	B	476	ASP	2.1
1	A	203	LEU	2.1
1	A	148	TYR	2.1
1	A	197	VAL	2.1
1	B	488	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	603	6/6	0.81	0.19	27,29,32,35	0
6	K	B	607	1/1	0.83	0.12	50,50,50,50	0
2	GOL	B	602	6/6	0.88	0.20	34,39,46,52	0
2	GOL	A	604	6/6	0.89	0.22	26,29,34,38	0
2	GOL	B	603	6/6	0.90	0.26	31,40,43,54	0
7	CL	B	608	1/1	0.90	0.06	59,59,59,59	0
2	GOL	A	602	6/6	0.92	0.14	27,35,36,41	0
4	MG	B	605	1/1	0.92	0.09	26,26,26,26	0
2	GOL	A	601	6/6	0.94	0.12	23,25,28,30	0
2	GOL	B	601	6/6	0.95	0.09	25,28,30,30	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NO3	B	606	4/4	0.98	0.12	31,32,32,35	0
3	OXL	B	604	6/6	0.98	0.09	25,28,30,31	0
3	OXL	A	605	6/6	0.98	0.09	19,20,22,23	0
5	NO3	A	607	4/4	0.99	0.12	32,33,36,36	0
4	MG	A	606	1/1	0.99	0.07	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.