



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 26, 2024 – 06:53 AM EDT

PDB ID : 6YQ8
Title : Crystal structure of native Phycocyanin from *T. elongatus* in spacegroup P63 at 1.8 Angstroms
Authors : Feiler, C.G.; Falke, S.; Sarrou, I.
Deposited on : 2020-04-16
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	162	
2	B	172	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	202	-	-	-	X
4	GOL	A	203	-	-	-	X
5	EDO	A	204	-	-	-	X
5	EDO	B	204	-	-	-	X
5	EDO	B	205	-	-	-	X
5	EDO	B	208	-	-	-	X
5	EDO	B	210	-	-	-	X
6	PGE	B	209	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 5677 atoms, of which 2720 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called C-phycoerythrin alpha chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	162	2479	783	1233	209	247	7	0	5	0

- Molecule 2 is a protein called C-phycoerythrin beta chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	172	2564	794	1282	229	251	8	0	2	0

- Molecule 3 is PHYCOCYANOBILIN (three-letter code: CYC) (formula: $C_{33}H_{40}N_4O_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
3	A	1	80	33	37	4	6	0	0
3	B	1	80	33	37	4	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	H	N			O
3	B	1	80	33	37	4	6	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	14	3	8	3	0	0
4	A	1	14	3	8	3	0	0
4	B	1	14	3	8	3	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			10	2	6	2		
5	B	1	Total	C	H	O	0	0
			10	2	6	2		
5	B	1	Total	C	H	O	0	0
			10	2	6	2		
5	B	1	Total	C	H	O	0	0
			10	2	6	2		
5	B	1	Total	C	H	O	0	0
			10	2	6	2		
5	B	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C H O 24 6 14 4	0	0
6	A	1	Total C O 7 4 3	0	0
6	B	1	Total C O 7 4 3	0	0
6	B	1	Total C H O 24 6 14 4	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	78	Total O 78 78	0	0
7	B	142	Total O 142 142	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: C-phyco cyanin alpha chain



- Molecule 2: C-phyco cyanin beta chain



4 Data and refinement statistics i

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	153.51Å 153.51Å 39.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.32 – 1.82 44.32 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.32-1.82) 99.9 (44.32-1.82)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.02 (at 1.82Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.179 , 0.201 0.179 , 0.201	Depositor DCC
R_{free} test set	2100 reflections (4.36%)	wwPDB-VP
Wilson B-factor (Å ²)	41.3	Xtrriage
Anisotropy	0.027	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 52.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.024 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5677	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CYC, PGE, EDO, GOL, MEN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.50	0/1287	0.65	3/1744 (0.2%)
2	B	0.66	1/1292 (0.1%)	0.66	1/1748 (0.1%)
All	All	0.58	1/2579 (0.0%)	0.65	4/3492 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	86	MET	CG-SD	-5.36	1.67	1.81

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	38	MET	CG-SD-CE	6.95	111.32	100.20
1	A	33	ARG	NE-CZ-NH2	-5.98	117.31	120.30
2	B	86	MET	CB-CG-SD	5.93	130.19	112.40
1	A	33	ARG	NE-CZ-NH1	5.56	123.08	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1246	1233	1229	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1282	1282	1286	10	0
3	A	43	37	37	5	0
3	B	86	74	74	7	0
4	A	12	16	16	3	0
4	B	6	8	8	1	0
5	A	4	6	6	0	0
5	B	24	36	36	1	0
6	A	17	14	23	4	0
6	B	17	14	23	4	0
7	A	78	0	0	3	1
7	B	142	0	0	3	1
All	All	2957	2720	2738	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:86:MET:CE	2:B:86:MET:SD	2.04	1.44
1:A:15:GLN:NE2	7:A:301:HOH:O	1.88	1.07
6:B:207:PGE:O3	7:B:301:HOH:O	2.07	0.72
2:B:144:ASP:OD2	6:B:207:PGE:H3	1.99	0.63
2:B:141:ILE:HA	6:B:207:PGE:H32	1.83	0.59
2:B:113:LEU:HD13	3:B:201:CYC:HMB3	1.84	0.59
1:A:84:CYS:HA	3:A:201:CYC:HHD	1.84	0.59
2:B:36:LYS:HE2	3:B:202:CYC:HMD3	1.86	0.58
1:A:42[A]:ARG:NH1	2:B:25:ASP:OD1	2.31	0.56
2:B:86:MET:CE	2:B:86:MET:CG	2.83	0.56
1:A:117:GLU:OE1	4:A:202:GOL:O2	2.26	0.53
3:B:201:CYC:HMD2	3:B:201:CYC:HC	1.75	0.52
6:A:206:PGE:H62	2:B:1:MET:CA	2.39	0.51
3:B:202:CYC:HBC3	3:B:202:CYC:HHD	1.93	0.50
1:A:55:ALA:O	1:A:59:VAL:HG23	2.13	0.49
3:B:202:CYC:HMA1	3:B:202:CYC:NB	2.28	0.48
4:B:203:GOL:H2	7:B:363:HOH:O	2.14	0.48
2:B:102:SER:OG	6:B:209:PGE:H42	2.15	0.47
1:A:114:GLY:H	4:A:202:GOL:H2	1.82	0.45
3:A:201:CYC:HMD2	3:A:201:CYC:HC	1.81	0.45
3:B:201:CYC:HMA3	3:B:201:CYC:HBA1	1.99	0.45
6:A:206:PGE:H62	2:B:1:MET:N	2.33	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:201:CYC:NB	3:A:201:CYC:HMA1	2.33	0.44
1:A:93:ARG:HG2	1:A:97:TYR:CZ	2.53	0.43
1:A:129:TYR:O	1:A:133:LEU:HG	2.18	0.43
1:A:22:THR:HG21	6:A:205:PGE:H6	2.01	0.43
6:A:205:PGE:C5	7:A:314:HOH:O	2.67	0.43
1:A:90:TYR:O	1:A:94:MET:HG2	2.19	0.43
1:A:21[A]:ASN:OD1	1:A:25:GLN:OE1	2.37	0.42
1:A:1:MET:H2	4:A:203:GOL:C3	2.34	0.41
1:A:83:LYS:HE2	7:A:356:HOH:O	2.19	0.41
3:A:201:CYC:HHA	3:A:201:CYC:HBD1	2.02	0.41
3:A:201:CYC:HMD2	3:A:201:CYC:NC	2.35	0.41
3:B:201:CYC:NB	3:B:201:CYC:HMA1	2.35	0.41
5:B:204:EDO:H21	7:B:320:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:374:HOH:O	7:B:434:HOH:O[1_556]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	165/162 (102%)	161 (98%)	3 (2%)	1 (1%)	25	12
2	B	171/172 (99%)	169 (99%)	2 (1%)	0	100	100
All	All	336/334 (101%)	330 (98%)	5 (2%)	1 (0%)	41	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	76	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	128/123 (104%)	122 (95%)	6 (5%)	26	11
2	B	129/127 (102%)	129 (100%)	0	100	100
All	All	257/250 (103%)	251 (98%)	6 (2%)	55	37

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	ARG
1	A	62	LYS
1	A	76	SER
1	A	127[A]	SER
1	A	127[B]	SER
1	A	140	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MEN	B	72	2	7,8,9	1.19	1 (14%)	6,9,11	0.98	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MEN	B	72	2	-	2/7/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	72	MEN	OD1-CG	2.45	1.28	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	72	MEN	CB-CA-C	-2.18	107.39	111.47

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	72	MEN	CA-CB-CG-OD1
2	B	72	MEN	CA-CB-CG-ND2

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	B	204	-	3,3,3	0.50	0	2,2,2	0.21	0
5	EDO	B	206	-	3,3,3	0.52	0	2,2,2	0.05	0
3	CYC	B	202	2	42,46,46	5.01	17 (40%)	50,67,67	2.15	18 (36%)
3	CYC	B	201	2	42,46,46	5.24	17 (40%)	50,67,67	2.39	12 (24%)
5	EDO	B	208	-	3,3,3	0.49	0	2,2,2	0.23	0
5	EDO	B	211	-	3,3,3	0.47	0	2,2,2	0.29	0
6	PGE	B	209	-	9,9,9	0.32	0	8,8,8	0.40	0
4	GOL	A	202	-	5,5,5	0.75	0	5,5,5	0.83	0
6	PGE	B	207	-	6,6,9	0.29	0	5,5,8	0.37	0
5	EDO	B	205	-	3,3,3	0.45	0	2,2,2	0.47	0
4	GOL	A	203	-	5,5,5	0.87	0	5,5,5	0.87	0
5	EDO	B	210	-	3,3,3	0.53	0	2,2,2	0.08	0
4	GOL	B	203	-	5,5,5	1.08	0	5,5,5	0.87	0
6	PGE	A	205	-	9,9,9	0.30	0	8,8,8	0.43	0
5	EDO	A	204	-	3,3,3	0.40	0	2,2,2	0.52	0
3	CYC	A	201	1	42,46,46	5.25	18 (42%)	50,67,67	2.09	11 (22%)
6	PGE	A	206	-	6,6,9	0.25	0	5,5,8	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	204	-	-	1/1/1/1	-
5	EDO	B	206	-	-	1/1/1/1	-
3	CYC	B	202	2	-	8/25/74/74	0/4/4/4
3	CYC	B	201	2	-	12/25/74/74	0/4/4/4
5	EDO	B	208	-	-	0/1/1/1	-
5	EDO	B	211	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PGE	B	209	-	-	2/7/7/7	-
4	GOL	A	202	-	-	2/4/4/4	-
6	PGE	B	207	-	-	3/4/4/7	-
5	EDO	B	205	-	-	1/1/1/1	-
4	GOL	A	203	-	-	0/4/4/4	-
5	EDO	B	210	-	-	0/1/1/1	-
4	GOL	B	203	-	-	2/4/4/4	-
6	PGE	A	205	-	-	5/7/7/7	-
5	EDO	A	204	-	-	1/1/1/1	-
3	CYC	A	201	1	-	10/25/74/74	0/4/4/4
6	PGE	A	206	-	-	2/4/4/7	-

All (52) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	201	CYC	C2C-C1C	24.00	1.73	1.52
3	B	202	CYC	C2C-C1C	23.36	1.72	1.52
3	B	201	CYC	C2C-C1C	23.29	1.72	1.52
3	B	201	CYC	C1C-NC	-11.23	1.23	1.37
3	A	201	CYC	C1C-NC	-9.83	1.24	1.37
3	A	201	CYC	C4C-NC	9.31	1.57	1.37
3	B	201	CYC	C2C-C3C	-9.29	1.28	1.54
3	B	202	CYC	C4C-NC	8.84	1.56	1.37
3	B	202	CYC	C1C-NC	-8.05	1.27	1.37
3	B	201	CYC	C4C-NC	7.89	1.54	1.37
3	A	201	CYC	C3C-C4C	7.82	1.62	1.50
3	B	201	CYC	C3C-C4C	7.63	1.62	1.50
3	A	201	CYC	C2C-C3C	-7.52	1.33	1.54
3	B	202	CYC	C2C-C3C	-7.48	1.33	1.54
3	B	201	CYC	C1B-NB	7.41	1.50	1.37
3	A	201	CYC	C1B-NB	7.27	1.50	1.37
3	B	202	CYC	C4B-NB	6.97	1.52	1.38
3	B	202	CYC	C1B-NB	6.57	1.48	1.37
3	A	201	CYC	C4B-NB	6.54	1.51	1.38
3	B	202	CYC	C3C-C4C	5.82	1.59	1.50
3	B	201	CYC	C4B-NB	5.59	1.49	1.38
3	B	202	CYC	CHB-C4A	5.28	1.52	1.40
3	B	201	CYC	CMC-C2C	5.26	1.64	1.53
3	A	201	CYC	CHB-C4A	5.21	1.52	1.40
3	B	201	CYC	CHB-C4A	5.15	1.52	1.40
3	B	201	CYC	C1A-NA	5.08	1.49	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	202	CYC	C4A-NA	4.59	1.47	1.36
3	B	202	CYC	C1A-NA	4.55	1.48	1.38
3	B	201	CYC	CAC-C3C	4.42	1.62	1.54
3	B	202	CYC	C2A-C3A	4.42	1.46	1.36
3	A	201	CYC	CHA-C1A	4.42	1.38	1.35
3	B	201	CYC	C2A-C3A	4.35	1.46	1.36
3	A	201	CYC	C1A-NA	4.18	1.47	1.38
3	A	201	CYC	CAC-C3C	4.12	1.62	1.54
3	B	202	CYC	CAC-C3C	4.11	1.62	1.54
3	A	201	CYC	C4A-NA	4.10	1.46	1.36
3	A	201	CYC	C2A-C3A	3.86	1.44	1.36
3	A	201	CYC	C1A-C2A	3.66	1.51	1.45
3	B	202	CYC	CHA-C1A	3.65	1.38	1.35
3	A	201	CYC	C1D-CHD	3.65	1.55	1.41
3	B	201	CYC	C4A-NA	3.61	1.44	1.36
3	A	201	CYC	CMC-C2C	3.46	1.60	1.53
3	B	202	CYC	C1D-CHD	3.43	1.54	1.41
3	B	201	CYC	C1D-CHD	3.39	1.54	1.41
3	B	201	CYC	C1A-C2A	3.35	1.51	1.45
3	B	202	CYC	C1A-C2A	3.27	1.50	1.45
3	A	201	CYC	C4D-CHA	3.25	1.53	1.41
3	A	201	CYC	C3B-C2B	2.50	1.42	1.36
3	B	201	CYC	C4D-CHA	2.45	1.50	1.41
3	B	202	CYC	C1B-C2B	2.41	1.49	1.45
3	B	202	CYC	C4D-CHA	2.25	1.49	1.41
3	B	201	CYC	C1B-C2B	2.13	1.48	1.45

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	201	CYC	C4D-CHA-C1A	-9.28	117.72	128.81
3	B	201	CYC	C4D-CHA-C1A	-7.62	119.70	128.81
3	B	201	CYC	CHD-C4C-NC	7.42	134.03	125.20
3	B	202	CYC	C4D-CHA-C1A	-6.81	120.67	128.81
3	B	201	CYC	C2C-C3C-C4C	5.13	109.02	101.34
3	B	201	CYC	CMC-C2C-C1C	-4.90	101.84	112.40
3	B	202	CYC	CMA-C3A-C4A	4.27	131.63	125.06
3	B	202	CYC	OC-C1C-C2C	-4.22	122.81	126.17
3	B	201	CYC	CBC-CAC-C3C	-4.18	104.17	113.47
3	B	201	CYC	OC-C1C-C2C	-4.11	122.90	126.17
3	A	201	CYC	CHD-C4C-NC	3.87	129.81	125.20
3	B	202	CYC	CMC-C2C-C1C	-3.58	104.69	112.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	202	CYC	C1B-NB-C4B	-3.25	106.53	110.67
3	B	202	CYC	O2A-CGA-CBA	3.16	124.20	114.03
3	A	201	CYC	CAB-C3B-C4B	3.05	126.20	121.38
3	A	201	CYC	C1A-C2A-C3A	2.96	110.06	106.78
3	A	201	CYC	C2A-C1A-NA	-2.95	105.76	110.05
3	A	201	CYC	C1B-CHB-C4A	-2.95	120.89	128.08
3	B	202	CYC	CMB-C2B-C1B	-2.90	120.55	124.17
3	B	202	CYC	C1B-CHB-C4A	-2.87	121.06	128.08
3	B	202	CYC	OB-C4B-C3B	2.84	131.12	128.04
3	B	201	CYC	C1B-NB-C4B	-2.82	107.08	110.67
3	A	201	CYC	C1B-C2B-C3B	2.76	110.75	107.87
3	B	201	CYC	CAB-C3B-C4B	2.75	125.73	121.38
3	B	202	CYC	C2C-C3C-C4C	2.73	105.43	101.34
3	B	202	CYC	CHB-C4A-NA	-2.69	119.31	124.93
3	A	201	CYC	O2D-CGD-CBD	2.69	122.66	114.03
3	B	202	CYC	C1A-C2A-C3A	2.68	109.75	106.78
3	B	202	CYC	OC-C1C-NC	2.64	128.14	124.94
3	A	201	CYC	CMA-C3A-C4A	2.64	129.13	125.06
3	A	201	CYC	C1B-NB-C4B	-2.53	107.44	110.67
3	B	202	CYC	O1A-CGA-CBA	-2.43	115.29	123.08
3	B	201	CYC	CAB-C3B-C2B	-2.30	123.59	127.53
3	B	202	CYC	CAC-C3C-C4C	-2.26	106.86	112.67
3	B	201	CYC	C2A-C1A-NA	-2.26	106.76	110.05
3	B	202	CYC	C2A-C1A-NA	-2.25	106.78	110.05
3	B	201	CYC	O2A-CGA-CBA	2.24	121.23	114.03
3	B	202	CYC	CBD-CAD-C3D	2.17	116.32	112.62
3	B	202	CYC	CAB-C3B-C4B	2.14	124.75	121.38
3	B	201	CYC	C1B-CHB-C4A	-2.13	122.89	128.08
3	A	201	CYC	CHB-C4A-NA	-2.03	120.69	124.93

There are no chirality outliers.

All (50) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	201	CYC	NA-C4A-CHB-C1B
3	A	201	CYC	C3A-C4A-CHB-C1B
3	A	201	CYC	C2C-C3C-CAC-CBC
3	A	201	CYC	C4C-C3C-CAC-CBC
3	B	201	CYC	NA-C4A-CHB-C1B
3	B	201	CYC	C3A-C4A-CHB-C1B
3	B	201	CYC	C4B-C3B-CAB-CBB
3	B	201	CYC	ND-C1D-CHD-C4C

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
3	B	201	CYC	C2D-C1D-CHD-C4C
3	B	202	CYC	NA-C4A-CHB-C1B
3	B	202	CYC	C3A-C4A-CHB-C1B
4	B	203	GOL	O1-C1-C2-C3
3	B	201	CYC	C2B-C3B-CAB-CBB
4	B	203	GOL	O1-C1-C2-O2
6	B	209	PGE	O2-C3-C4-O3
4	A	202	GOL	O1-C1-C2-C3
6	A	205	PGE	O1-C1-C2-O2
6	A	205	PGE	O3-C5-C6-O4
5	A	204	EDO	O1-C1-C2-O2
5	B	204	EDO	O1-C1-C2-O2
3	A	201	CYC	C2B-C3B-CAB-CBB
6	B	209	PGE	O1-C1-C2-O2
4	A	202	GOL	O1-C1-C2-O2
3	B	202	CYC	C4C-C3C-CAC-CBC
6	B	207	PGE	O1-C1-C2-O2
6	B	207	PGE	C1-C2-O2-C3
3	B	201	CYC	C3A-C2A-CAA-CBA
6	A	205	PGE	C4-C3-O2-C2
3	B	201	CYC	C1A-C2A-CAA-CBA
3	A	201	CYC	CAA-CBA-CGA-O2A
3	A	201	CYC	CAA-CBA-CGA-O1A
3	A	201	CYC	NA-C1A-CHA-C4D
3	B	202	CYC	CAA-CBA-CGA-O1A
3	A	201	CYC	NC-C4C-CHD-C1D
3	B	202	CYC	C2C-C3C-CAC-CBC
3	B	202	CYC	CAA-CBA-CGA-O2A
6	A	205	PGE	C1-C2-O2-C3
3	B	202	CYC	CAD-CBD-CGD-O2D
6	A	205	PGE	O2-C3-C4-O3
3	B	201	CYC	CAD-CBD-CGD-O1D
3	B	201	CYC	CAA-CBA-CGA-O1A
3	B	202	CYC	CAD-CBD-CGD-O1D
5	B	205	EDO	O1-C1-C2-O2
5	B	206	EDO	O1-C1-C2-O2
6	A	206	PGE	C6-C5-O3-C4
3	B	201	CYC	CAD-CBD-CGD-O2D
3	B	201	CYC	CAA-CBA-CGA-O2A
6	A	206	PGE	C3-C4-O3-C5
6	B	207	PGE	O2-C3-C4-O3
3	A	201	CYC	CAD-CBD-CGD-O2D


There are no ring outliers.

11 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	204	EDO	1	0
3	B	202	CYC	3	0
3	B	201	CYC	4	0
6	B	209	PGE	1	0
4	A	202	GOL	2	0
6	B	207	PGE	3	0
4	A	203	GOL	1	0
4	B	203	GOL	1	0
6	A	205	PGE	2	0
3	A	201	CYC	5	0
6	A	206	PGE	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	162/162 (100%)	0.38	13 (8%) 12 9	39, 64, 100, 116	0
2	B	171/172 (99%)	-0.16	0 100 100	34, 46, 62, 79	0
All	All	333/334 (99%)	0.10	13 (3%) 39 34	34, 51, 90, 116	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	59	VAL	6.0
1	A	63	PHE	4.4
1	A	60	TYR	3.5
1	A	67	THR	3.2
1	A	64	PRO	3.1
1	A	58	ALA	2.8
1	A	55	ALA	2.7
1	A	70	GLN	2.7
1	A	61	GLN	2.5
1	A	56	ALA	2.4
1	A	68	THR	2.3
1	A	62	LYS	2.2
1	A	138	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MEN	B	72	9/10	0.96	0.10	47,53,64,64	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	B	208	4/4	-0.02	0.41	142,170,177,180	0
4	GOL	A	202	6/6	0.16	0.41	166,199,208,210	0
5	EDO	B	206	4/4	0.26	0.30	111,133,142,145	0
6	PGE	B	209	10/10	0.28	0.48	147,181,194,195	0
5	EDO	B	205	4/4	0.31	0.54	163,196,197,197	0
5	EDO	B	210	4/4	0.40	0.57	121,145,149,149	0
5	EDO	A	204	4/4	0.41	1.11	186,224,225,225	0
4	GOL	A	203	6/6	0.45	0.41	172,207,211,211	0
6	PGE	A	206	7/10	0.50	0.29	128,129,130,130	0
5	EDO	B	204	4/4	0.52	0.49	121,145,152,154	0
5	EDO	B	211	4/4	0.61	0.15	151,181,183,183	0
6	PGE	A	205	10/10	0.62	0.38	138,168,192,193	0
4	GOL	B	203	6/6	0.74	0.26	82,98,112,114	0
6	PGE	B	207	7/10	0.84	0.28	89,106,128,131	0
3	CYC	A	201	43/43	0.90	0.12	46,61,85,89	0
3	CYC	B	201	43/43	0.94	0.11	34,51,68,69	0
3	CYC	B	202	43/43	0.96	0.11	34,41,58,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around CYC A 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.