



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 19, 2023 – 07:58 PM JST

PDB ID : 7BPZ  
Title : X-ray structure of human PPARalpha ligand binding domain-bezafibrate-SR  
C1 coactivator peptide co-crystals obtained by soaking  
Authors : Kamata, S.; Ishikawa, R.; Akahane, M.; Oyama, T.; Ishii, I.  
Deposited on : 2020-03-23  
Resolution : 2.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	273	
1	C	273	
2	B	15	
2	D	15	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4422 atoms, of which 38 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisome proliferator-activated receptor alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	265	2099	1345	352	383	19	0	0	0
1	C	259	2055	1318	343	376	18	0	0	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	196	GLY	-	expression tag	UNP Q07869
A	197	SER	-	expression tag	UNP Q07869
A	198	HIS	-	expression tag	UNP Q07869
A	199	MET	-	expression tag	UNP Q07869
C	196	GLY	-	expression tag	UNP Q07869
C	197	SER	-	expression tag	UNP Q07869
C	198	HIS	-	expression tag	UNP Q07869
C	199	MET	-	expression tag	UNP Q07869

- Molecule 2 is a protein called 15-meric peptide from Nuclear receptor coactivator 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	11	101	64	23	14	0	0	0
2	D	8	71	47	15	9	0	0	0

- Molecule 3 is 2-[P-[2-P-CHLOROBENZAMIDO)ETHYL]PHENOXY]-2-METHYLPRO PIONIC ACID (three-letter code: PEM) (formula: C<sub>19</sub>H<sub>20</sub>ClNO<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	Cl	H	N			O
3	A	1	44	19	1	19	1	4	0	0
3	C	1	44	19	1	19	1	4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	3	Total 3 3	0	0
4	C	5	Total 5 5	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Peroxisome proliferator-activated receptor alpha





- Molecule 1: Peroxisome proliferator-activated receptor alpha



- Molecule 2: 15-meric peptide from Nuclear receptor coactivator 1



- Molecule 2: 15-meric peptide from Nuclear receptor coactivator 1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.13Å 101.62Å 61.10Å 90.00° 100.93° 90.00°	Depositor
Resolution (Å)	36.07 – 2.43 36.07 – 2.43	Depositor EDS
% Data completeness (in resolution range)	93.3 (36.07-2.43) 99.6 (36.07-2.43)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.61 (at 2.42Å)	Xtrriage
Refinement program	PHENIX 1.11.1-2575-000	Depositor
R, $R_{free}$	0.201 , 0.242 0.203 , 0.245	Depositor DCC
$R_{free}$ test set	1358 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.4	Xtrriage
Anisotropy	0.151	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 45.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.024 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4422	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/2137	0.49	0/2882
1	C	0.34	0/2092	0.51	0/2822
2	B	0.24	0/102	0.42	0/134
2	D	0.24	0/71	0.51	0/93
All	All	0.33	0/4402	0.50	0/5931

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2099	0	2121	45	0
1	C	2055	0	2075	53	0
2	B	101	0	110	3	0
2	D	71	0	84	5	0
3	A	25	19	19	3	0
3	C	25	19	19	1	0
4	A	3	0	0	0	0
4	C	5	0	0	0	0
All	All	4384	38	4428	103	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:203:LEU:HD23	1:C:205:SER:HB3	1.26	1.08
1:C:203:LEU:HA	1:C:205:SER:N	1.75	1.01
1:A:324:VAL:HG22	1:A:331:LEU:HB2	1.48	0.95
1:C:206:LEU:HD11	1:C:295:PRO:HG2	1.48	0.93
1:A:230:SER:OG	1:A:231:GLY:HA2	1.75	0.87
1:C:301:ASP:O	1:C:302:LEU:HB3	1.72	0.87
1:C:230:SER:OG	1:C:231:GLY:HA2	1.76	0.84
1:C:203:LEU:HB3	1:C:206:LEU:H	1.43	0.81
1:C:203:LEU:HA	1:C:205:SER:H	1.45	0.79
1:C:403:GLY:O	1:C:407:VAL:HG23	1.83	0.78
1:C:301:ASP:OD2	1:C:392:LEU:HD12	1.86	0.75
1:A:302:LEU:O	1:A:303:ASN:HB2	1.90	0.71
1:A:224:LYS:O	1:A:227:VAL:HG12	1.95	0.67
1:A:446:ILE:O	1:A:450:THR:HB	1.95	0.67
1:C:280:SER:O	1:C:284:VAL:HG23	1.96	0.64
1:C:269:GLU:HG3	1:C:347:LEU:HG	1.79	0.64
1:A:310:LYS:HD3	1:A:311:TYR:CE2	2.33	0.64
1:C:306:VAL:HG13	2:D:690:LEU:HD23	1.80	0.63
1:C:206:LEU:CD1	1:C:295:PRO:HG2	2.26	0.62
1:A:219:ASN:O	1:A:224:LYS:HE3	2.00	0.62
1:C:388:ARG:O	1:C:391:LEU:HD22	2.01	0.61
1:C:412:LEU:HD21	1:C:426:LEU:HD22	1.82	0.61
1:A:230:SER:CB	1:A:231:GLY:HA2	2.29	0.59
1:A:462:GLU:OE1	2:B:689:ILE:HG13	2.03	0.58
1:A:255:VAL:HG23	1:A:256:ALA:H	1.69	0.58
1:C:202:ASP:OD1	1:C:203:LEU:HG	2.03	0.58
1:C:464:TYR:O	1:C:466:ASP:HA	2.03	0.58
1:C:203:LEU:CD2	1:C:205:SER:HB3	2.18	0.56
1:C:254:LEU:HD12	1:C:254:LEU:H	1.70	0.56
1:C:464:TYR:OH	3:C:501:PEM:O4	2.24	0.56
1:A:203:LEU:HG	1:A:407:VAL:HG22	1.89	0.55
1:C:307:THR:HG21	1:C:391:LEU:HD13	1.87	0.55
1:A:273:PHE:CE2	1:A:454:ALA:HB1	2.43	0.54
1:A:350:PRO:HB2	1:A:443:LEU:HD11	1.89	0.54
1:C:384:CYS:O	1:C:401:GLN:HB2	2.08	0.53
1:C:302:LEU:HD22	1:C:302:LEU:O	2.08	0.53
1:C:429:LYS:O	1:C:433:LEU:HD22	2.08	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:203:LEU:HA	1:C:204:LYS:C	2.26	0.53
1:A:277:GLN:NE2	1:A:456:LEU:HD23	2.24	0.52
1:C:203:LEU:HB3	1:C:206:LEU:N	2.20	0.52
1:C:235:ASN:O	1:C:237:PRO:HD3	2.10	0.52
1:A:221:ASN:OD1	1:A:223:VAL:HG12	2.10	0.51
1:C:465:ARG:O	1:C:465:ARG:HD3	2.10	0.51
1:C:203:LEU:CB	1:C:206:LEU:H	2.17	0.51
1:A:409:ARG:HG3	1:A:423:PHE:CD2	2.47	0.50
1:C:254:LEU:HD12	1:C:254:LEU:N	2.27	0.50
1:C:203:LEU:HB3	1:C:206:LEU:CB	2.41	0.50
1:A:324:VAL:CG2	1:A:331:LEU:HB2	2.31	0.50
1:A:324:VAL:HG13	1:A:324:VAL:O	2.12	0.49
1:C:462:GLU:OE2	2:D:689:ILE:HG22	2.10	0.49
1:A:273:PHE:CD2	1:A:454:ALA:HB1	2.47	0.49
1:A:307:THR:HG21	1:A:391:LEU:HD13	1.93	0.49
1:C:423:PHE:HB3	1:C:424:PRO:HD3	1.95	0.48
2:D:689:ILE:HD13	2:D:692:ARG:NH1	2.29	0.48
1:C:401:GLN:HG2	1:C:405:VAL:HG23	1.96	0.48
1:A:408:LEU:O	1:A:412:LEU:HG	2.14	0.48
1:C:354:ILE:HG22	1:C:355:MET:HE2	1.96	0.48
1:C:301:ASP:OD2	1:C:303:ASN:HB2	2.14	0.47
1:A:211:TYR:CZ	1:A:215:LEU:HD21	2.49	0.47
1:C:302:LEU:HD13	1:C:303:ASN:N	2.30	0.46
1:A:465:ARG:O	1:A:466:ASP:HB2	2.15	0.46
1:A:409:ARG:HG3	1:A:423:PHE:CG	2.50	0.46
1:A:403:GLY:O	1:A:407:VAL:HG23	2.16	0.46
1:A:279:THR:HG21	3:A:501:PEM:O1	2.17	0.45
1:A:301:ASP:O	1:A:302:LEU:HB3	2.16	0.45
1:A:302:LEU:O	1:A:303:ASN:CB	2.61	0.45
1:A:232:LYS:O	1:A:233:ALA:HB3	2.17	0.45
1:C:302:LEU:HD13	1:C:302:LEU:C	2.37	0.45
2:B:689:ILE:O	2:B:693:LEU:HD13	2.17	0.45
1:A:279:THR:HG21	3:A:501:PEM:C7	2.48	0.44
1:A:421:PHE:O	1:A:424:PRO:HG2	2.17	0.44
1:C:391:LEU:HG	1:C:397:ILE:HD12	1.99	0.44
1:C:421:PHE:O	1:C:424:PRO:HD2	2.17	0.44
2:D:689:ILE:HD12	2:D:689:ILE:O	2.17	0.44
1:C:321:LEU:HD11	1:C:330:MET:HE1	1.99	0.44
1:C:321:LEU:HD11	1:C:330:MET:CE	2.48	0.44
1:A:368:LEU:HD23	1:A:426:LEU:CD1	2.48	0.44
1:C:377:LEU:HB3	1:C:408:LEU:HD13	2.00	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:307:THR:CG2	1:A:391:LEU:HD13	2.47	0.44
1:C:421:PHE:C	1:C:424:PRO:HD2	2.37	0.44
1:A:291:ALA:HA	1:A:294:ILE:HD12	2.00	0.43
1:A:227:VAL:HG13	1:A:228:ILE:N	2.33	0.43
1:A:388:ARG:HB2	1:A:391:LEU:HD22	2.00	0.43
1:C:203:LEU:HD22	1:C:206:LEU:N	2.34	0.43
1:A:325:MET:HG2	1:A:330:MET:HB3	2.00	0.43
1:A:426:LEU:HA	1:A:426:LEU:HD12	1.65	0.43
1:A:384:CYS:O	1:A:401:GLN:CB	2.67	0.42
1:C:465:ARG:HA	1:C:466:ASP:HA	1.74	0.42
1:A:368:LEU:HD23	1:A:426:LEU:HD11	2.01	0.42
1:C:344:LEU:HA	1:C:347:LEU:HD22	2.01	0.42
1:C:283:THR:O	1:C:287:LEU:HB2	2.20	0.42
2:D:689:ILE:HD11	2:D:693:LEU:HD21	2.02	0.42
1:A:443:LEU:O	1:A:447:ILE:HG13	2.20	0.42
1:C:324:VAL:O	1:C:324:VAL:HG13	2.20	0.42
1:A:276:CYS:HB3	3:A:501:PEM:H183	2.00	0.41
1:A:384:CYS:O	1:A:401:GLN:HB2	2.20	0.41
1:C:230:SER:OG	1:C:231:GLY:CA	2.58	0.41
1:A:349:LYS:HE3	1:A:349:LYS:HB2	1.75	0.41
2:B:690:LEU:HD12	2:B:690:LEU:HA	1.92	0.41
1:C:385:CYS:C	1:C:387:ASP:H	2.24	0.40
1:A:223:VAL:HG13	1:A:224:LYS:N	2.36	0.40
1:C:221:ASN:OD1	1:C:223:VAL:HG12	2.21	0.40
1:C:467:MET:HB3	1:C:467:MET:HE2	1.93	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	261/273 (96%)	244 (94%)	12 (5%)	5 (2%)	<b>8</b> <b>6</b>

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	255/273 (93%)	243 (95%)	10 (4%)	2 (1%)	19	22
2	B	9/15 (60%)	7 (78%)	2 (22%)	0	100	100
2	D	6/15 (40%)	6 (100%)	0	0	100	100
All	All	531/576 (92%)	500 (94%)	24 (4%)	7 (1%)	12	11

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	303	ASN
1	A	450	THR
1	C	302	LEU
1	A	301	ASP
1	A	232	LYS
1	C	204	LYS
1	A	200	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/238 (97%)	204 (88%)	27 (12%)	5	4
1	C	227/238 (95%)	204 (90%)	23 (10%)	7	7
2	B	11/14 (79%)	8 (73%)	3 (27%)	0	0
2	D	8/14 (57%)	5 (62%)	3 (38%)	0	0
All	All	477/504 (95%)	421 (88%)	56 (12%)	5	4

All (56) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	198	HIS
1	A	202	ASP
1	A	215	LEU
1	A	219	ASN

Continued on next page...

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	229	LEU
1	A	230	SER
1	A	245	GLU
1	A	255	VAL
1	A	287	LEU
1	A	302	LEU
1	A	303	ASN
1	A	309	LEU
1	A	324	VAL
1	A	344	LEU
1	A	347	LEU
1	A	364	LYS
1	A	368	LEU
1	A	377	LEU
1	A	391	LEU
1	A	392	LEU
1	A	420	ILE
1	A	426	LEU
1	A	427	LEU
1	A	433	LEU
1	A	434	ARG
1	A	435	GLN
1	A	453	ASP
2	B	688	LYS
2	B	695	GLN
2	B	696	GLU
1	C	203	LEU
1	C	208	LYS
1	C	209	ARG
1	C	215	LEU
1	C	254	LEU
1	C	287	LEU
1	C	302	LEU
1	C	331	LEU
1	C	344	LEU
1	C	347	LEU
1	C	377	LEU
1	C	385	CYS
1	C	391	LEU
1	C	410	LEU
1	C	412	LEU
1	C	422	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	427	LEU
1	C	433	LEU
1	C	452	SER
1	C	453	ASP
1	C	460	LEU
1	C	465	ARG
1	C	466	ASP
2	D	689	ILE
2	D	693	LEU
2	D	694	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PEM	A	501	-	26,26,26	1.35	2 (7%)	36,36,36	1.04	2 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PEM	C	501	-	26,26,26	1.28	2 (7%)	36,36,36	1.22	3 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEM	A	501	-	-	7/21/21/21	0/2/2/2
3	PEM	C	501	-	-	10/21/21/21	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	PEM	C7-N1	5.31	1.45	1.33
3	C	501	PEM	C7-N1	4.83	1.44	1.33
3	C	501	PEM	O1-C7	-2.38	1.18	1.23
3	A	501	PEM	C4-CL	2.00	1.78	1.74

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	501	PEM	C16-O2-C10	-3.24	115.76	120.95
3	A	501	PEM	C16-O2-C10	-2.64	116.72	120.95
3	A	501	PEM	O2-C16-C19	2.25	113.05	109.99
3	C	501	PEM	C8-C9-C13	-2.17	107.84	112.87
3	C	501	PEM	O3-C19-C16	2.10	120.34	113.99

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	501	PEM	C2-C1-C7-O1
3	C	501	PEM	C6-C1-C7-O1
3	C	501	PEM	C2-C1-C7-N1
3	C	501	PEM	C6-C1-C7-N1
3	A	501	PEM	C2-C1-C7-O1
3	A	501	PEM	C2-C1-C7-N1
3	A	501	PEM	C6-C1-C7-O1
3	A	501	PEM	C6-C1-C7-N1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	C	501	PEM	C18-C16-C19-O3
3	A	501	PEM	C15-C10-O2-C16
3	C	501	PEM	C15-C10-O2-C16
3	C	501	PEM	O2-C16-C19-O3
3	C	501	PEM	O2-C16-C19-O4
3	C	501	PEM	C17-C16-C19-O3
3	A	501	PEM	C11-C10-O2-C16
3	C	501	PEM	C11-C10-O2-C16
3	A	501	PEM	C18-C16-C19-O4


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	PEM	3	0
3	C	501	PEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	265/273 (97%)	0.17	18 (6%) 17 13	38, 59, 89, 100	1 (0%)
1	C	259/273 (94%)	0.09	14 (5%) 25 22	42, 58, 91, 107	1 (0%)
2	B	11/15 (73%)	1.05	3 (27%) 0 0	77, 82, 92, 95	0
2	D	8/15 (53%)	3.14	6 (75%) 0 0	90, 97, 108, 113	0
All	All	543/576 (94%)	0.19	41 (7%) 13 11	38, 58, 91, 113	2 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	692	ARG	4.9
2	D	691	HIS	4.4
2	D	693	LEU	4.3
1	C	255	VAL	3.8
1	A	317	ILE	3.8
1	C	468	TYR	3.7
1	A	232	LYS	3.6
1	A	256	ALA	3.3
1	A	321	LEU	3.2
2	D	688	LYS	3.2
1	A	255	VAL	3.1
1	C	317	ILE	3.0
2	D	694	LEU	3.0
1	C	205	SER	2.9
2	D	695	GLN	2.9
1	C	204	LYS	2.9
1	C	233	ALA	2.9
1	C	302	LEU	2.9
1	A	301	ASP	2.8
1	A	316	ALA	2.8
1	A	313	VAL	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	316	ALA	2.6
2	B	691	HIS	2.6
1	A	468	TYR	2.5
1	A	231	GLY	2.5
1	A	233	ALA	2.3
1	A	318	PHE	2.3
1	A	198	HIS	2.3
1	C	396	HIS	2.3
1	A	320	MET	2.3
1	A	319	ALA	2.3
1	C	321	LEU	2.2
1	C	203	LEU	2.2
1	C	319	ALA	2.1
2	B	696	GLU	2.1
1	A	276	CYS	2.1
2	B	695	GLN	2.1
1	C	301	ASP	2.0
1	A	314	TYR	2.0
1	C	465	ARG	2.0
1	A	202	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PEM	A	501	25/25	0.87	0.24	63,83,97,109	0
3	PEM	C	501	25/25	0.93	0.18	56,66,81,83	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.