



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 21, 2023 – 05:52 AM JST

PDB ID : 7EV6
Title : Crystal structure of the Lon-like protease MtaLonC with D581A mutation in complex with F-b20-Q
Authors : Hsieh, K.Y.; Kuo, C.I.; Su, S.C.; Huang, K.F.; Chang, C.I.
Deposited on : 2021-05-20
Resolution : 2.10 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	732	
2	S	5	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5039 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endopeptidase La.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	594	4565	2902	807	848	8	0	1	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	581	ALA	ASP	engineered mutation	UNP C9DRU9
A	720	LYS	-	expression tag	UNP C9DRU9
A	721	LEU	-	expression tag	UNP C9DRU9
A	722	ALA	-	expression tag	UNP C9DRU9
A	723	ALA	-	expression tag	UNP C9DRU9
A	724	ALA	-	expression tag	UNP C9DRU9
A	725	LEU	-	expression tag	UNP C9DRU9
A	726	GLU	-	expression tag	UNP C9DRU9
A	727	HIS	-	expression tag	UNP C9DRU9
A	728	HIS	-	expression tag	UNP C9DRU9
A	729	HIS	-	expression tag	UNP C9DRU9
A	730	HIS	-	expression tag	UNP C9DRU9
A	731	HIS	-	expression tag	UNP C9DRU9
A	732	HIS	-	expression tag	UNP C9DRU9

- Molecule 2 is a protein called F-b20-Q peptide {ortho-aminobenzoic acid (Abz)- QLRSLNG EWRFAWFPAPEAV[Tyr(3-NO₂)]A}.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	S	5	42	25	6	11	0	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO₄) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	418	Total	O	0	0
			418	418		
4	S	9	Total	O	0	0
			9	9		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endopeptidase La



- Molecule 2: F-b20-Q peptide {ortho-aminobenzoic acid (Abz)- QLRSLNGEWRFAWFPAPEA V[Tyr(3-NO₂)]A}



4 Data and refinement statistics

Property	Value	Source
Space group	P 6	Depositor
Cell constants a, b, c, α , β , γ	115.83Å 115.83Å 136.26Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.85 – 2.10 19.85 – 2.10	Depositor EDS
% Data completeness (in resolution range)	85.4 (19.85-2.10) 90.1 (19.85-2.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 2.11Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.154 , 0.178 0.170 , 0.197	Depositor DCC
R_{free} test set	2757 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	23.4	Xtrriage
Anisotropy	0.068	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.027 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5039	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NIY, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/4660	0.71	4/6330 (0.1%)
2	S	0.59	0/25	1.28	0/30
All	All	0.51	0/4685	0.71	4/6360 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	432	ARG	NE-CZ-NH2	-6.12	117.24	120.30
1	A	579	GLU	OE1-CD-OE2	-5.52	116.68	123.30
1	A	631	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	A	554	ARG	NE-CZ-NH1	5.22	122.91	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	692	PHE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4565	0	4589	42	0
2	S	42	0	31	1	0
3	A	5	0	0	0	0
4	A	418	0	0	3	0
4	S	9	0	0	0	0
All	All	5039	0	4620	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:549:LEU:CD2	1:A:610:VAL:HG21	2.24	0.67
1:A:498:GLU:HG2	1:A:518:THR:HG22	1.78	0.65
1:A:573:GLN:HA	1:A:573:GLN:HE21	1.65	0.61
1:A:692:PHE:HA	1:A:694:GLY:N	2.14	0.61
1:A:411:LEU:HD13	1:A:455:LEU:HD23	1.84	0.59
1:A:693:ARG:HA	1:A:697:GLU:OE1	2.03	0.59
1:A:692:PHE:HA	1:A:694:GLY:H	1.70	0.56
1:A:573:GLN:HA	1:A:573:GLN:NE2	2.20	0.55
1:A:249:GLN:HB2	1:A:293:ASN:HB2	1.87	0.55
1:A:105:VAL:HG21	1:A:227:TYR:OH	2.08	0.53
1:A:582:SER:OG	2:S:20:ALA:C	2.47	0.53
1:A:549:LEU:CD2	1:A:610:VAL:CG2	2.86	0.53
1:A:79:LEU:HD21	1:A:93:LEU:HD22	1.91	0.52
1:A:516:ARG:CZ	1:A:572:GLU:OE1	2.61	0.49
1:A:567:ILE:HD11	1:A:595:ILE:HD11	1.93	0.49
1:A:80:TYR:CE2	1:A:297:ILE:HG21	2.48	0.48
1:A:549:LEU:HD22	1:A:588:LEU:HB2	1.95	0.48
1:A:527:HIS:HB2	1:A:566:THR:HG23	1.96	0.47
1:A:98:GLU:HG2	1:A:254:LEU:HD12	1.96	0.47
1:A:243:GLU:O	1:A:245:LEU:HD12	2.15	0.47
1:A:692:PHE:CD1	1:A:692:PHE:O	2.69	0.46
1:A:549:LEU:HD21	1:A:610:VAL:HG21	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:VAL:HG12	1:A:294:VAL:HG13	1.99	0.45
1:A:42:ALA:O	1:A:357:GLN:NE2	2.51	0.44
1:A:96:GLY:HA2	4:A:1209:HOH:O	2.17	0.44
1:A:262:THR:O	1:A:262:THR:OG1	2.35	0.44
1:A:516:ARG:NE	1:A:572:GLU:OE1	2.51	0.44
1:A:48:HIS:CD2	1:A:358:VAL:H	2.35	0.44
1:A:84:SER:OG	1:A:85:GLU:N	2.51	0.44
1:A:58:GLY:HA2	4:A:1052:HOH:O	2.19	0.43
1:A:88:VAL:CG1	1:A:294:VAL:HG13	2.49	0.43
1:A:549:LEU:HD21	1:A:610:VAL:CG2	2.48	0.43
1:A:227:TYR:OH	1:A:252:PRO:HD3	2.19	0.42
1:A:44:ARG:NH2	4:A:911:HOH:O	2.52	0.42
1:A:613:THR:OG1	1:A:615:LYS:HG2	2.19	0.42
1:A:237:ARG:HG3	1:A:240:GLU:HB3	2.01	0.42
1:A:517:LEU:HD11	1:A:587:GLU:HG3	2.01	0.41
1:A:39:LEU:O	1:A:43:ILE:HG12	2.19	0.41
1:A:593:SER:HB2	1:A:600:LEU:HD11	2.01	0.41
1:A:423:ALA:HA	1:A:426[A]:MET:SD	2.61	0.41
1:A:49:ALA:O	1:A:359:ILE:HA	2.21	0.41
1:A:107:GLY:O	1:A:108:LEU:C	2.60	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	585/732 (80%)	567 (97%)	17 (3%)	1 (0%)	47 49
2	S	2/5 (40%)	2 (100%)	0	0	100 100
All	All	587/737 (80%)	569 (97%)	17 (3%)	1 (0%)	47 49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	84	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	464/577 (80%)	437 (94%)	27 (6%)	20	17
2	S	2/2 (100%)	2 (100%)	0	100	100
All	All	466/579 (80%)	439 (94%)	27 (6%)	20	17

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	ARG
1	A	39	LEU
1	A	57	LEU
1	A	73	GLU
1	A	79	LEU
1	A	85	GLU
1	A	86	ARG
1	A	93	LEU
1	A	98	GLU
1	A	225	ARG
1	A	237	ARG
1	A	254	LEU
1	A	262	THR
1	A	294	VAL
1	A	298	ARG
1	A	355	GLN
1	A	368	GLU
1	A	432	ARG
1	A	499	VAL
1	A	505	VAL
1	A	517	LEU
1	A	525	ARG
1	A	533	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	572	GLU
1	A	617	LEU
1	A	662	GLU
1	A	692	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	HIS
1	A	223	GLN
1	A	334	GLN
1	A	462	GLN
1	A	573	GLN
1	A	597	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NIY	S	19	2	13,15,16	4.55	2 (15%)	13,20,22	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NIY	S	19	2	-	2/7/10/12	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	S	19	NIY	O2-NN	13.05	1.44	1.22
2	S	19	NIY	CE1-NN	-9.68	1.28	1.45

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	S	19	NIY	CD1-CE1-NN-O2
2	S	19	NIY	CZ-CE1-NN-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	A	801	-	4,4,4	1.11	0	6,6,6	1.03	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	594/732 (81%)	0.16	55 (9%) 8 11	12, 24, 81, 99	0
2	S	4/5 (80%)	0.32	0 100 100	22, 27, 31, 41	0
All	All	598/737 (81%)	0.17	55 (9%) 9 11	12, 24, 81, 99	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	239	ALA	6.9
1	A	284	LEU	6.0
1	A	692	PHE	6.0
1	A	244	PRO	5.7
1	A	342	GLN	4.6
1	A	243	GLU	4.6
1	A	712	GLY	4.6
1	A	344	PRO	4.5
1	A	241	THR	4.1
1	A	285	VAL	4.0
1	A	103	GLU	4.0
1	A	225	ARG	3.9
1	A	108	LEU	3.7
1	A	100	HIS	3.5
1	A	237	ARG	3.4
1	A	218	PHE	3.4
1	A	236	ALA	3.4
1	A	341	PRO	3.3
1	A	238	TYR	3.2
1	A	219	GLN	3.1
1	A	107	GLY	3.1
1	A	230	ALA	3.1
1	A	105	VAL	3.0
1	A	106	GLU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	298	ARG	3.0
1	A	231	LEU	2.9
1	A	222	PRO	2.9
1	A	525	ARG	2.8
1	A	283	TYR	2.8
1	A	258	SER	2.8
1	A	232	ARG	2.7
1	A	227	TYR	2.7
1	A	691	GLY	2.7
1	A	693	ARG	2.7
1	A	259	SER	2.5
1	A	224	ALA	2.5
1	A	99	ILE	2.5
1	A	223	GLN	2.4
1	A	73	GLU	2.4
1	A	290	TRP	2.4
1	A	220	ARG	2.4
1	A	96	GLY	2.3
1	A	71	SER	2.3
1	A	15	GLU	2.3
1	A	249	GLN	2.3
1	A	260	SER	2.3
1	A	294	VAL	2.2
1	A	527	HIS	2.2
1	A	711	GLU	2.2
1	A	343	ALA	2.1
1	A	86	ARG	2.1
1	A	708	ARG	2.1
1	A	242	GLY	2.1
1	A	229	GLU	2.0
1	A	95	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
2	NIY	S	19	15/16	0.90	0.18	20,29,43,49	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	A	801	5/5	0.98	0.08	23,23,24,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.