



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 11, 2023 – 02:09 AM EDT

PDB ID : 7K7C  
Title : Crystal structure of diphtheria toxin from crystals obtained at pH 5.5  
Authors : Lovell, S.; Kashipathy, M.M.; Battaile, K.P.; Rodnin, M.V.; Ladokhin, A.S.  
Deposited on : 2020-09-22  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	538	
1	B	538	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7754 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Diphtheria toxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	502	3710	2360	625	712	13	0	0	0
1	B	500	3691	2339	622	718	12	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	51	GLU	LYS	engineered mutation	UNP Q5PY51
A	148	LYS	GLU	engineered mutation	UNP Q5PY51
A	536	MET	-	cloning artifact	UNP Q5PY51
A	537	ALA	-	cloning artifact	UNP Q5PY51
B	51	GLU	LYS	engineered mutation	UNP Q5PY51
B	148	LYS	GLU	engineered mutation	UNP Q5PY51
B	536	MET	-	cloning artifact	UNP Q5PY51
B	537	ALA	-	cloning artifact	UNP Q5PY51


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	189	Total	O	0	0
			189	189		
2	B	164	Total	O	0	0
			164	164		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Diphtheria toxin



- Molecule 1: Diphtheria toxin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.55Å 69.67Å 73.40Å 117.57° 93.48° 98.11°	Depositor
Resolution (Å)	35.02 – 2.05 46.77 – 2.05	Depositor EDS
% Data completeness (in resolution range)	96.9 (35.02-2.05) 96.9 (46.77-2.05)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.88 (at 2.05Å)	Xtrriage
Refinement program	PHENIX 1.14rc3_3208	Depositor
R, $R_{free}$	0.183 , 0.243 0.189 , 0.246	Depositor DCC
$R_{free}$ test set	3554 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.2	Xtrriage
Anisotropy	0.359	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 44.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7754	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/3782	0.58	0/5137
1	B	0.47	0/3762	0.59	1/5113 (0.0%)
All	All	0.47	0/7544	0.59	1/10250 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	455	ARG	NE-CZ-NH2	-5.27	117.67	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3710	0	3557	14	0
1	B	3691	0	3509	23	0
2	A	189	0	0	1	0
2	B	164	0	0	0	0
All	All	7754	0	7066	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:LYS:O	1:A:238:VAL:HG13	1.99	0.62
1:B:536:MET:O	1:B:537:ALA:HB2	2.02	0.59
1:B:398:TRP:HH2	1:B:509:ILE:HD12	1.68	0.59
1:B:225:PRO:O	1:B:229:LYS:N	2.36	0.59
1:A:226:ILE:O	1:A:229:LYS:N	2.38	0.55
1:B:108:LEU:HB3	1:B:117:GLN:OE1	2.06	0.55
1:A:216:LYS:O	1:A:220:LEU:HD12	2.07	0.55
1:A:186:CYS:O	1:A:187:ALA:HB3	2.08	0.52
1:B:406:ILE:HB	1:B:533:ILE:HD13	1.92	0.51
1:B:142:GLU:CD	1:B:142:GLU:H	2.13	0.50
1:B:398:TRP:CH2	1:B:509:ILE:HD12	2.46	0.50
1:A:5:VAL:HG21	1:A:105:GLU:CB	2.43	0.49
1:A:516:LYS:HG3	1:A:523:VAL:HG13	1.95	0.49
1:B:83:VAL:HG12	1:B:161:VAL:HG22	1.96	0.48
1:B:186:CYS:O	1:B:187:ALA:HB3	2.14	0.48
1:B:28:VAL:HG11	1:B:166:ASN:ND2	2.28	0.48
1:B:109:SER:O	1:B:117:GLN:NE2	2.47	0.47
1:B:504:ILE:O	1:B:504:ILE:CG2	2.63	0.46
1:B:451:SER:HA	1:B:455:ARG:O	2.16	0.45
1:A:225:PRO:O	1:A:229:LYS:N	2.50	0.44
1:B:504:ILE:O	1:B:504:ILE:HG23	2.17	0.44
1:B:536:MET:O	1:B:537:ALA:CB	2.66	0.43
1:B:90:LYS:NZ	1:B:127:PHE:O	2.49	0.43
1:A:102:ILE:O	1:A:106:LEU:HB2	2.19	0.42
1:A:373:ASN:O	1:A:377:ARG:HG3	2.19	0.42
1:B:5:VAL:CG2	1:B:106:LEU:HD13	2.49	0.42
1:B:523:VAL:HG22	1:B:524:ASN:N	2.33	0.42
1:B:18:SER:OG	1:B:82:LYS:HE2	2.20	0.42
1:B:450:ILE:HG23	1:B:457:ILE:HB	2.01	0.42
1:A:387:GLN:NE2	2:A:601:HOH:O	2.20	0.41
1:A:451:SER:HA	1:A:455:ARG:O	2.19	0.41
1:B:20:TYR:O	1:B:55:SER:HB2	2.20	0.41
1:B:220:LEU:HD13	1:B:338:LEU:CD1	2.51	0.41
1:A:486:ASN:ND2	1:B:142:GLU:HG2	2.36	0.40
1:A:83:VAL:HA	1:A:160:SER:O	2.21	0.40
1:A:136:LEU:O	1:A:149:TYR:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	488/538 (91%)	473 (97%)	12 (2%)	3 (1%)	25	15
1	B	486/538 (90%)	473 (97%)	11 (2%)	2 (0%)	34	24
All	All	974/1076 (90%)	946 (97%)	23 (2%)	5 (0%)	29	18

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	201	CYS
1	A	237	THR
1	B	107	GLY
1	A	68	ASP
1	B	347	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	381/454 (84%)	377 (99%)	4 (1%)	76	75
1	B	381/454 (84%)	374 (98%)	7 (2%)	59	55
All	All	762/908 (84%)	751 (99%)	11 (1%)	67	65

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	SER

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	A	226	ILE
1	A	291	SER
1	A	504	ILE
1	B	9	SER
1	B	30	SER
1	B	133	ARG
1	B	291	SER
1	B	475	SER
1	B	504	ILE
1	B	516	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	502/538 (93%)	0.11	25 (4%) 28 31	27, 45, 70, 88	0
1	B	500/538 (92%)	0.05	22 (4%) 34 37	27, 44, 70, 100	0
All	All	1002/1076 (93%)	0.08	47 (4%) 31 33	27, 45, 70, 100	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	123	PHE	4.0
1	A	111	THR	3.7
1	A	226	ILE	3.4
1	A	225	PRO	3.4
1	B	420	ILE	3.4
1	A	200	SER	3.3
1	A	347	VAL	3.3
1	B	124	ILE	3.1
1	B	110	LEU	3.0
1	B	234	PRO	3.0
1	B	233	SER	2.9
1	A	237	THR	2.9
1	B	125	LYS	2.8
1	A	124	ILE	2.7
1	B	111	THR	2.7
1	A	108	LEU	2.7
1	B	422	ALA	2.7
1	A	346	LEU	2.7
1	B	231	SER	2.6
1	B	126	ARG	2.6
1	A	234	PRO	2.6
1	B	230	MET	2.6
1	A	110	LEU	2.6
1	B	348	GLY	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	102	ILE	2.5
1	A	291	SER	2.5
1	A	348	GLY	2.5
1	B	395	ALA	2.5
1	B	421	THR	2.4
1	A	233	SER	2.4
1	A	231	SER	2.4
1	A	106	LEU	2.4
1	A	230	MET	2.3
1	B	238	VAL	2.3
1	B	479	VAL	2.3
1	B	396	VAL	2.2
1	B	108	LEU	2.2
1	A	107	GLY	2.2
1	B	346	LEU	2.2
1	A	420	ILE	2.2
1	A	485	ALA	2.1
1	A	235	ASN	2.1
1	A	114	LEU	2.1
1	A	126	ARG	2.1
1	B	109	SER	2.0
1	B	237	THR	2.0
1	A	390	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.